



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 06:38 am BST

PDB ID : 6CJ5
Title : Crystal Structure of Mnk2-D228G in Complex With Inhibitor
Authors : Han, Q.
Deposited on : 2018-02-26
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MAP kinase-interacting serine/threonine-protein kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	271	2163	1371	376	401	15	0	0	0


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	70	GLY	-	expression tag	UNP Q9HBH9
A	71	SER	-	expression tag	UNP Q9HBH9
A	228	GLY	ASP	engineered mutation	UNP Q9HBH9

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 3-(pyridin-3-yl)imidazo[1,2-a]pyridine-8-carboxamide (three-letter code: F4G) (formula: C₁₃H₁₀N₄O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	18	13	4	1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	8	8	8	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	105.37Å 105.37Å 72.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	56.70 – 2.80 56.72 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (56.70-2.80) 99.9 (56.72-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.204 , 0.280 0.218 , 0.293	Depositor DCC
R_{free} test set	569 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	70.3	Xtrriage
Anisotropy	0.014	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 56.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.038 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2190	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, F4G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.67	1/2209 (0.0%)	0.83	1/2978 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	367	TRP	CD2-CE2	5.81	1.48	1.41

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	122	ILE	N-CA-C	-7.13	91.74	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2163	0	2097	106	0
2	A	1	0	0	0	0
3	A	18	0	0	1	0
4	A	8	0	0	2	0
All	All	2190	0	2097	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:CYS:CB	1:A:313:ALA:HB1	1.53	1.35
1:A:122:ILE:HG23	1:A:125:ARG:HD2	1.27	1.14
1:A:299:CYS:HB3	1:A:313:ALA:CB	1.79	1.12
1:A:86:GLN:OE1	1:A:102:ILE:HD12	1.53	1.09
1:A:89:VAL:HG22	1:A:97:ARG:HH22	1.21	1.04
1:A:299:CYS:HB3	1:A:313:ALA:HB1	1.04	1.01
1:A:121:HIS:C	1:A:123:ARG:H	1.57	0.96
1:A:217:ASN:HD22	1:A:217:ASN:H	1.05	0.95
1:A:121:HIS:C	1:A:123:ARG:N	2.17	0.94
1:A:89:VAL:HG22	1:A:97:ARG:NH2	1.83	0.93
1:A:299:CYS:CB	1:A:313:ALA:CB	2.40	0.93
1:A:122:ILE:HG23	1:A:125:ARG:CD	2.00	0.92
1:A:365:HIS:HD2	1:A:367:TRP:H	1.19	0.87
1:A:89:VAL:CG2	1:A:97:ARG:HH22	1.89	0.84
1:A:150:GLU:HG3	1:A:155:PHE:CE2	2.13	0.83
1:A:86:GLN:OE1	1:A:102:ILE:CD1	2.30	0.78
1:A:365:HIS:CD2	1:A:367:TRP:H	2.02	0.78
1:A:299:CYS:CB	1:A:313:ALA:O	2.33	0.76
1:A:278:LEU:HG	1:A:357:LEU:O	1.86	0.75
1:A:217:ASN:N	1:A:217:ASN:HD22	1.79	0.75
1:A:371:CYS:SG	4:A:503:HOH:O	2.43	0.75
1:A:299:CYS:HB3	1:A:313:ALA:CA	2.18	0.74
1:A:203:HIS:CD2	1:A:227:PHE:HB3	2.24	0.73
1:A:347:LYS:HB3	1:A:357:LEU:CD1	2.19	0.72
1:A:139:HIS:HD2	1:A:141:ASN:H	1.36	0.72
1:A:299:CYS:SG	1:A:313:ALA:CB	2.77	0.71
1:A:312:PRO:C	1:A:314:CYS:H	1.91	0.71
1:A:299:CYS:HB2	1:A:313:ALA:HB1	1.64	0.71
1:A:347:LYS:HB3	1:A:357:LEU:HD11	1.72	0.71
1:A:108:GLN:NE2	1:A:109:GLU:H	1.88	0.70
1:A:204:ARG:NH1	1:A:273:ASP:OD1	2.27	0.67
1:A:149:PHE:HB2	1:A:156:TYR:HB2	1.77	0.67
1:A:275:ARG:O	1:A:278:LEU:HB2	1.95	0.67
1:A:263:GLU:O	1:A:266:SER:HB2	1.94	0.66
1:A:78:ARG:O	1:A:81:ASP:HB2	1.97	0.65
1:A:312:PRO:C	1:A:314:CYS:N	2.51	0.63
1:A:259:PRO:HA	1:A:262:VAL:HG13	1.81	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275:ARG:HA	1:A:278:LEU:HD23	1.81	0.62
1:A:299:CYS:SG	1:A:313:ALA:HB1	2.39	0.62
1:A:299:CYS:HB3	1:A:313:ALA:O	2.00	0.62
1:A:263:GLU:O	1:A:266:SER:CB	2.48	0.61
1:A:139:HIS:CD2	1:A:141:ASN:H	2.18	0.60
1:A:299:CYS:CA	1:A:313:ALA:O	2.50	0.60
1:A:298:ARG:O	1:A:317:MET:HG3	2.02	0.60
1:A:299:CYS:SG	1:A:313:ALA:HB3	2.41	0.60
1:A:103:ASN:HD21	1:A:105:ILE:HG22	1.67	0.58
1:A:162:MET:O	3:A:402:F4G:N18	2.37	0.58
1:A:117:LYS:HB2	1:A:153:ASP:O	2.04	0.57
1:A:278:LEU:HD12	1:A:348:LEU:CD2	2.35	0.57
1:A:122:ILE:O	1:A:125:ARG:HB3	2.05	0.56
1:A:327:TYR:OH	1:A:346:SER:HA	2.05	0.56
1:A:253:SER:O	1:A:257:MET:HG3	2.04	0.56
1:A:115:ILE:HB	1:A:155:PHE:HB2	1.88	0.56
1:A:311:CYS:O	1:A:314:CYS:HB2	2.06	0.55
1:A:103:ASN:HD21	1:A:105:ILE:CG2	2.20	0.55
1:A:285:LEU:HD12	1:A:285:LEU:O	2.08	0.53
1:A:217:ASN:H	1:A:217:ASN:ND2	1.87	0.53
1:A:186:VAL:HG21	1:A:285:LEU:HD22	1.90	0.53
1:A:197:HIS:CD2	1:A:274:LYS:HG2	2.44	0.52
1:A:299:CYS:SG	1:A:313:ALA:O	2.67	0.52
1:A:290:SER:HB3	1:A:334:TRP:CZ2	2.45	0.52
1:A:347:LYS:HD3	1:A:357:LEU:HD11	1.92	0.52
1:A:97:ARG:HD3	1:A:99:GLN:OE1	2.11	0.51
1:A:364:GLN:HA	1:A:364:GLN:NE2	2.25	0.51
1:A:347:LYS:HB3	1:A:357:LEU:HD13	1.92	0.51
1:A:122:ILE:CG2	1:A:125:ARG:HD2	2.19	0.50
1:A:139:HIS:HD2	1:A:141:ASN:N	2.08	0.50
1:A:93:GLY:C	1:A:95:HIS:H	2.15	0.50
1:A:108:GLN:HE21	1:A:109:GLU:H	1.57	0.50
1:A:136:CYS:O	1:A:142:VAL:HG11	2.12	0.49
1:A:299:CYS:HB3	1:A:313:ALA:C	2.31	0.49
1:A:103:ASN:OD1	1:A:106:THR:HG23	2.13	0.48
1:A:117:LYS:HB2	1:A:153:ASP:HA	1.95	0.48
1:A:122:ILE:HG22	1:A:122:ILE:O	2.12	0.48
1:A:337:ILE:O	1:A:342:LYS:NZ	2.46	0.48
1:A:283:VAL:HG13	1:A:294:PRO:HD2	1.96	0.48
1:A:121:HIS:O	1:A:123:ARG:N	2.46	0.48
1:A:347:LYS:O	1:A:357:LEU:HD13	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93:GLY:N	4:A:501:HOH:O	2.47	0.48
1:A:358:SER:O	1:A:362:VAL:HG23	2.15	0.47
1:A:214:GLU:OE2	1:A:221:PRO:HG2	2.15	0.47
1:A:267:GLU:O	1:A:271:ILE:HD12	2.16	0.46
1:A:116:GLU:OE2	1:A:154:ARG:NE	2.47	0.46
1:A:312:PRO:O	1:A:314:CYS:N	2.49	0.46
1:A:80:GLU:HA	1:A:83:TYR:O	2.17	0.45
1:A:344:LEU:HD11	1:A:362:VAL:HG13	1.99	0.44
1:A:278:LEU:HD12	1:A:348:LEU:HD22	1.97	0.44
1:A:285:LEU:HD12	1:A:285:LEU:C	2.38	0.44
1:A:108:GLN:NE2	1:A:109:GLU:N	2.63	0.44
1:A:82:VAL:HG13	1:A:83:TYR:CD1	2.53	0.43
1:A:275:ARG:O	1:A:278:LEU:N	2.45	0.43
1:A:94:ALA:O	1:A:95:HIS:HB2	2.18	0.43
1:A:334:TRP:HA	1:A:337:ILE:HD12	2.01	0.43
1:A:108:GLN:HE21	1:A:109:GLU:N	2.17	0.43
1:A:121:HIS:CA	1:A:123:ARG:H	2.29	0.42
1:A:299:CYS:SG	1:A:313:ALA:C	2.98	0.42
1:A:141:ASN:HB2	1:A:192:ALA:HB2	2.02	0.42
1:A:103:ASN:O	1:A:107:SER:N	2.53	0.41
1:A:259:PRO:O	1:A:263:GLU:HB2	2.20	0.41
1:A:100:THR:HG22	1:A:101:CYS:N	2.35	0.41
1:A:197:HIS:CG	1:A:274:LYS:HG2	2.56	0.41
1:A:299:CYS:N	1:A:313:ALA:O	2.53	0.41
1:A:352:ASP:HB3	1:A:355:GLN:HG2	2.01	0.41
1:A:213:CYS:HA	1:A:222:VAL:HG12	2.02	0.40
1:A:321:SER:O	1:A:324:GLU:HB3	2.21	0.40
1:A:117:LYS:N	1:A:153:ASP:O	2.45	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	263/316 (83%)	228 (87%)	29 (11%)	6 (2%)	6 21

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	95	HIS
1	A	96	ALA
1	A	119	PRO
1	A	253	SER
1	A	89	VAL
1	A	313	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	234/274 (85%)	201 (86%)	33 (14%)	3 10

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	SER
1	A	76	SER
1	A	80	GLU
1	A	84	GLN
1	A	87	GLU
1	A	104	LEU
1	A	107	SER
1	A	116	GLU
1	A	118	GLN
1	A	121	HIS
1	A	128	ARG
1	A	129	GLU
1	A	144	GLU
1	A	153	ASP
1	A	180	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	191	SER
1	A	209	GLU
1	A	212	LEU
1	A	217	ASN
1	A	219	VAL
1	A	255	GLU
1	A	262	VAL
1	A	263	GLU
1	A	273	ASP
1	A	274	LYS
1	A	285	LEU
1	A	298	ARG
1	A	299	CYS
1	A	311	CYS
1	A	323	GLN
1	A	332	LYS
1	A	357	LEU
1	A	369	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	108	GLN
1	A	139	HIS
1	A	217	ASN
1	A	323	GLN
1	A	364	GLN
1	A	365	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	F4G	A	402	-	17,20,20	2.07	3 (17%)	19,28,28	1.16	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	F4G	A	402	-	-	0/8/8/8	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	F4G	C1-C2	7.27	1.51	1.43
3	A	402	F4G	C10-C9	2.50	1.51	1.48
3	A	402	F4G	C4-C5	2.16	1.41	1.35

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	F4G	C5-C4-N3	2.66	124.19	120.78
3	A	402	F4G	C13-N12-C11	2.30	120.82	116.85

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	F4G	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	271/316 (85%)	0.30	14 (5%) 27 18	40, 61, 112, 163	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	71	SER	6.6
1	A	313	ALA	6.4
1	A	70	GLY	5.9
1	A	121	HIS	5.1
1	A	73	ASP	3.8
1	A	72	THR	3.7
1	A	298	ARG	3.7
1	A	310	ALA	3.0
1	A	122	ILE	2.9
1	A	227	PHE	2.4
1	A	94	ALA	2.2
1	A	120	GLY	2.2
1	A	93	GLY	2.1
1	A	118	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	F4G	A	402	18/18	0.84	0.29	76,97,106,113	0
2	ZN	A	401	1/1	0.96	0.07	163,163,163,163	0

6.5 Other polymers [i](#)

There are no such residues in this entry.