



Full wwPDB X-ray Structure Validation Report ⓘ

May 27, 2020 – 02:07 am BST

PDB ID : 6CLU
Title : Staphylococcus aureus Dihydropteroate Synthase (saDHPS) F17L E208K double mutant structure
Authors : Gajewski, S.; Griffith, E.C.; Wu, Y.; White, S.W.
Deposited on : 2018-03-02
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	291	 84% 13%
1	B	291	 71% 27%
1	C	291	 75% 21%
1	D	291	 83% 14%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 7252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydropteroate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	253	1909	1204	329	364	12	0	1	0
1	B	212	1557	989	262	297	9	0	1	0
1	C	230	1711	1087	287	327	10	0	0	0
1	D	251	1878	1181	326	359	12	0	1	0

There are 104 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MET	-	initiating methionine	UNP O05701
A	-22	GLY	-	expression tag	UNP O05701
A	-21	HIS	-	expression tag	UNP O05701
A	-20	HIS	-	expression tag	UNP O05701
A	-19	HIS	-	expression tag	UNP O05701
A	-18	HIS	-	expression tag	UNP O05701
A	-17	HIS	-	expression tag	UNP O05701
A	-16	HIS	-	expression tag	UNP O05701
A	-15	HIS	-	expression tag	UNP O05701
A	-14	HIS	-	expression tag	UNP O05701
A	-13	HIS	-	expression tag	UNP O05701
A	-12	HIS	-	expression tag	UNP O05701
A	-11	SER	-	expression tag	UNP O05701
A	-10	SER	-	expression tag	UNP O05701
A	-9	GLY	-	expression tag	UNP O05701
A	-8	HIS	-	expression tag	UNP O05701
A	-7	ILE	-	expression tag	UNP O05701
A	-6	GLU	-	expression tag	UNP O05701
A	-5	GLY	-	expression tag	UNP O05701
A	-4	ARG	-	expression tag	UNP O05701
A	-3	HIS	-	expression tag	UNP O05701

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP O05701
A	-1	LEU	-	expression tag	UNP O05701
A	0	GLU	-	expression tag	UNP O05701
A	17	LEU	PHE	engineered mutation	UNP O05701
A	208	LYS	GLU	engineered mutation	UNP O05701
B	-23	MET	-	initiating methionine	UNP O05701
B	-22	GLY	-	expression tag	UNP O05701
B	-21	HIS	-	expression tag	UNP O05701
B	-20	HIS	-	expression tag	UNP O05701
B	-19	HIS	-	expression tag	UNP O05701
B	-18	HIS	-	expression tag	UNP O05701
B	-17	HIS	-	expression tag	UNP O05701
B	-16	HIS	-	expression tag	UNP O05701
B	-15	HIS	-	expression tag	UNP O05701
B	-14	HIS	-	expression tag	UNP O05701
B	-13	HIS	-	expression tag	UNP O05701
B	-12	HIS	-	expression tag	UNP O05701
B	-11	SER	-	expression tag	UNP O05701
B	-10	SER	-	expression tag	UNP O05701
B	-9	GLY	-	expression tag	UNP O05701
B	-8	HIS	-	expression tag	UNP O05701
B	-7	ILE	-	expression tag	UNP O05701
B	-6	GLU	-	expression tag	UNP O05701
B	-5	GLY	-	expression tag	UNP O05701
B	-4	ARG	-	expression tag	UNP O05701
B	-3	HIS	-	expression tag	UNP O05701
B	-2	MET	-	expression tag	UNP O05701
B	-1	LEU	-	expression tag	UNP O05701
B	0	GLU	-	expression tag	UNP O05701
B	17	LEU	PHE	engineered mutation	UNP O05701
B	208	LYS	GLU	engineered mutation	UNP O05701
C	-23	MET	-	initiating methionine	UNP O05701
C	-22	GLY	-	expression tag	UNP O05701
C	-21	HIS	-	expression tag	UNP O05701
C	-20	HIS	-	expression tag	UNP O05701
C	-19	HIS	-	expression tag	UNP O05701
C	-18	HIS	-	expression tag	UNP O05701
C	-17	HIS	-	expression tag	UNP O05701
C	-16	HIS	-	expression tag	UNP O05701
C	-15	HIS	-	expression tag	UNP O05701
C	-14	HIS	-	expression tag	UNP O05701
C	-13	HIS	-	expression tag	UNP O05701

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-12	HIS	-	expression tag	UNP O05701
C	-11	SER	-	expression tag	UNP O05701
C	-10	SER	-	expression tag	UNP O05701
C	-9	GLY	-	expression tag	UNP O05701
C	-8	HIS	-	expression tag	UNP O05701
C	-7	ILE	-	expression tag	UNP O05701
C	-6	GLU	-	expression tag	UNP O05701
C	-5	GLY	-	expression tag	UNP O05701
C	-4	ARG	-	expression tag	UNP O05701
C	-3	HIS	-	expression tag	UNP O05701
C	-2	MET	-	expression tag	UNP O05701
C	-1	LEU	-	expression tag	UNP O05701
C	0	GLU	-	expression tag	UNP O05701
C	17	LEU	PHE	engineered mutation	UNP O05701
C	208	LYS	GLU	engineered mutation	UNP O05701
D	-23	MET	-	initiating methionine	UNP O05701
D	-22	GLY	-	expression tag	UNP O05701
D	-21	HIS	-	expression tag	UNP O05701
D	-20	HIS	-	expression tag	UNP O05701
D	-19	HIS	-	expression tag	UNP O05701
D	-18	HIS	-	expression tag	UNP O05701
D	-17	HIS	-	expression tag	UNP O05701
D	-16	HIS	-	expression tag	UNP O05701
D	-15	HIS	-	expression tag	UNP O05701
D	-14	HIS	-	expression tag	UNP O05701
D	-13	HIS	-	expression tag	UNP O05701
D	-12	HIS	-	expression tag	UNP O05701
D	-11	SER	-	expression tag	UNP O05701
D	-10	SER	-	expression tag	UNP O05701
D	-9	GLY	-	expression tag	UNP O05701
D	-8	HIS	-	expression tag	UNP O05701
D	-7	ILE	-	expression tag	UNP O05701
D	-6	GLU	-	expression tag	UNP O05701
D	-5	GLY	-	expression tag	UNP O05701
D	-4	ARG	-	expression tag	UNP O05701
D	-3	HIS	-	expression tag	UNP O05701
D	-2	MET	-	expression tag	UNP O05701
D	-1	LEU	-	expression tag	UNP O05701
D	0	GLU	-	expression tag	UNP O05701
D	17	LEU	PHE	engineered mutation	UNP O05701
D	208	LYS	GLU	engineered mutation	UNP O05701


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	73	Total O 73 73	0	0
2	B	26	Total O 26 26	0	0
2	C	50	Total O 50 50	0	0
2	D	48	Total O 48 48	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dihydropteroate synthase




- Molecule 1: Dihydropteroate synthase



- Molecule 1: Dihydropteroate synthase



- Molecule 1: Dihydropteroate synthase



10259	R263	HIS	ASN	PHE	SER
-------	------	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	76.64Å 76.64Å 176.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.62 – 1.95 46.62 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.62-1.95) 95.6 (46.62-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.77 (at 1.95Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.182 , 0.204 0.170 , 0.186	Depositor DCC
R_{free} test set	2006 reflections (2.72%)	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtriage
Anisotropy	0.180	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 26.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	0.327 for h,-k,-l	Xtriage
Reported twinning fraction	0.320 for h,-k,-l	Depositor
Outliers	0 of 73619 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7252	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/1937	0.40	0/2625
1	B	0.23	0/1575	0.38	0/2139
1	C	0.23	0/1730	0.39	0/2345
1	D	0.23	0/1906	0.39	0/2584
All	All	0.23	0/7148	0.39	0/9693

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1909	0	1901	5	0
1	B	1557	0	1516	3	0
1	C	1711	0	1692	6	0
1	D	1878	0	1846	6	0
2	A	73	0	0	0	0
2	B	26	0	0	0	0
2	C	50	0	0	0	0
2	D	48	0	0	0	0
All	All	7252	0	6955	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:224:ALA:HB1	1:C:249:LEU:HD22	1.82	0.60
1:D:224:ALA:HB1	1:D:249:LEU:HD22	1.83	0.60
1:A:89:GLU:OE2	1:A:113:ARG:NH2	2.38	0.56
1:C:88:SER:HB2	1:C:117:VAL:HG21	1.93	0.51
1:A:154:ALA:HB1	1:A:159:ILE:HD12	1.93	0.49
1:C:64:LEU:HD21	1:C:90:VAL:HA	1.93	0.49
1:C:201:SER:HB2	1:C:240:VAL:HA	1.95	0.48
1:C:33:VAL:HA	1:C:36:MET:HE3	1.97	0.47
1:A:201:SER:HB2	1:A:240:VAL:HA	1.95	0.47
1:D:14:PRO:HD3	1:D:22:LYS:HD3	1.97	0.46
1:B:126:VAL:HG22	1:B:165:TRP:HB2	1.99	0.43
1:D:201:SER:HB2	1:D:240:VAL:HA	2.01	0.43
1:A:236:ARG:HA	1:A:236:ARG:HH21	1.84	0.42
1:B:89:GLU:OE2	1:B:113:ARG:NH1	2.53	0.42
1:D:126:VAL:HG22	1:D:165:TRP:HB2	2.02	0.42
1:D:165:TRP:CE3	1:D:195:PRO:HB2	2.55	0.41
1:B:227:THR:HG23	1:B:238:VAL:HG11	2.03	0.41
1:A:41:ALA:HB2	1:A:243:VAL:HG11	2.03	0.41
1:C:184:ALA:HB2	1:D:259:ASN:ND2	2.36	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/291 (86%)	248 (99%)	2 (1%)	0	100	100
1	B	201/291 (69%)	200 (100%)	1 (0%)	0	100	100
1	C	220/291 (76%)	220 (100%)	0	0	100	100
1	D	248/291 (85%)	245 (99%)	3 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	919/1164 (79%)	913 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/244 (82%)	199 (100%)	0	100	100
1	B	155/244 (64%)	155 (100%)	0	100	100
1	C	175/244 (72%)	175 (100%)	0	100	100
1	D	192/244 (79%)	192 (100%)	0	100	100
All	All	721/976 (74%)	721 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	150	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	253/291 (86%)	0.11	2 (0%) 86 90	19, 28, 43, 66	0
1	B	212/291 (72%)	0.15	3 (1%) 75 82	22, 32, 45, 55	0
1	C	230/291 (79%)	0.14	3 (1%) 77 83	21, 31, 44, 51	0
1	D	251/291 (86%)	0.14	2 (0%) 86 90	21, 33, 47, 57	0
All	All	946/1164 (81%)	0.13	10 (1%) 80 85	19, 31, 45, 66	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	217	VAL	4.4
1	B	172	PHE	4.0
1	D	212	TYR	3.0
1	C	215	THR	2.4
1	B	220	ASP	2.4
1	C	262	ALA	2.3
1	B	221	GLU	2.2
1	D	2	THR	2.2
1	A	211	GLY	2.2
1	A	21	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.