

Full wwPDB X-ray Structure Validation Report (i)

Dec 17, 2023 – 07:21 PM EST

PDB ID	:	1CPU
Title	:	SUBSITE MAPPING OF THE ACTIVE SITE OF HUMAN PANCREATIC
		ALPHA-AMYLASE USING SUBSTRATES, THE PHARMACOLOGICAL
		INHIBITOR ACARBOSE, AND AN ACTIVE SITE VARIANT
Authors	:	Brayer, G.D.; Sidhu, G.; Maurus, R.; Rydberg, E.H.; Braun, C.; Wang, Y.;
		Nguyen, N.T.; Overall, C.M.; Withers, S.G.
Deposited on	:	1999-06-07
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# {\rm Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	А	496	85%	13%	•
2	В	2	100%		_
3	С	2	100%		



$1 \mathrm{CPU}$

2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 4277 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called PROTEIN (ALPHA-AMYLASE).


Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	496	$\begin{array}{c} \text{Total} \\ 3945 \end{array}$	C 2497	N 696	О 732	S 20	0	0	0

• Molecule 2 is an oligosaccharide called 4-amino-4,6-dideoxy-alpha-D-glucopyranose-(1-4)-al pha-D-glucopyranose.



Mol	Chain	Residues	A	Aton	ns		ZeroOcc	AltConf	Trace
2	В	2	Total 22	C 12	N 1	O 9	0	0	0


• Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	At	\mathbf{oms}		ZeroOcc	AltConf	Trace
3	С	2	Total 22	C 12	O 10	0	0	0

• Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).





Mol	Chain	Residues	A	Atoms			ZeroOcc	AltConf
4	Δ	1	Total	С	Ν	Ο	0	0
T	11	Ĩ	14	8	1	5	0	0

• Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	Total Ca 1 1	0	0

• Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	1	Total Cl 1 1	0	0

• Molecule 7 is 5-HYDROXYMETHYL-CHONDURITOL (three-letter code: HMC) (formula: $C_7H_{12}O_5$).





ſ	Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
	7	А	1	Total 11	$\begin{array}{c} \mathrm{C} \\ 7 \end{array}$	0 4	0	0

• Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	А	261	Total O 261 261	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Chain A:
 85%
 13%

 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
 0
- Molecule 1: PROTEIN (ALPHA-AMYLASE)

• Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain C:

100%





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	52.91Å 68.90Å 131.77Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 - 2.00	Depositor
% Data completeness	93.8 (8.00-2.00)	Depositor
(in resolution range)	55.0 (0.00-2.00)	Depositor
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.170 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4277	wwPDB-VP
Average B, all atoms $(Å^2)$	15.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, CA, PCA, NAG, GLC, AGL, HMC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	nd lengths	Bond angles	
Mol Ch	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.39	1/4053~(0.0%)	0.60	3/5506~(0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(Å)	Ideal(Å)
1	А	484	GLU	CB-CG	-6.46	1.39	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	А	26	LEU	CA-CB-CG	6.17	129.49	115.30
1	А	115	CYS	CA-CB-SG	5.85	124.54	114.00
1	А	329	LEU	CA-CB-CG	5.27	127.42	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3945	0	3717	60	0
2	В	22	0	20	0	0
3	С	22	0	19	0	0
4	А	14	0	13	0	0

Continued on next page...



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	А	1	0	0	0	0
6	А	1	0	0	0	0
7	А	11	0	9	1	0
8	А	261	0	0	5	0
All	All	4277	0	3778	60	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:A:142:LYS:H	1:A:142:LYS:HD2	1.13	1.06
1:A:142:LYS:HD2	1:A:142:LYS:N	1.72	1.03
1:A:231:TYR:HD2	1:A:293:LEU:HG	1.30	0.96
1:A:142:LYS:H	1:A:142:LYS:CD	1.85	0.90
1:A:314:THR:HG23	1:A:316:TRP:H	1.40	0.87
1:A:325:VAL:O	1:A:329:LEU:HD22	1.82	0.78
1:A:329:LEU:HD22	1:A:329:LEU:H	1.50	0.76
1:A:261:LYS:O	1:A:265:VAL:HG12	1.85	0.76
1:A:8:GLN:H	1:A:8:GLN:CD	1.93	0.72
1:A:231:TYR:CD2	1:A:293:LEU:HG	2.20	0.69
1:A:227:LYS:HA	1:A:227:LYS:HE2	1.76	0.67
1:A:275:SER:O	1:A:278:LYS:HE2	1.96	0.65
1:A:306:GLY:HA3	8:A:632:HOH:O	1.99	0.62
1:A:8:GLN:CD	1:A:8:GLN:N	2.55	0.59
1:A:314:THR:HG21	1:A:316:TRP:HD1	1.68	0.59
1:A:329:LEU:CD2	1:A:394:MET:HB3	2.32	0.58
1:A:329:LEU:HD21	1:A:394:MET:HB3	1.86	0.57
1:A:216:ASN:OD1	1:A:227:LYS:HE3	2.05	0.57
1:A:314:THR:HG22	8:A:677:HOH:O	2.05	0.55
1:A:142:LYS:CE	1:A:155:THR:HG22	2.36	0.55
1:A:369:GLU:HG2	8:A:703:HOH:O	2.06	0.54
1:A:211:LEU:HA	1:A:214:LEU:HD22	1.90	0.54
1:A:347:GLN:HB3	1:A:354:VAL:HG22	1.90	0.53
1:A:460:GLY:O	1:A:496:LEU:HD11	2.10	0.52
1:A:264:THR:HG23	1:A:269:TRP:HB2	1.92	0.51
1:A:23:ASP:HA	1:A:26:LEU:HD21	1.92	0.51
1:A:380:ASN:O	1:A:381:ASP:HB2	2.10	0.50
1:A:349:GLN:O	1:A:352:ASN:ND2	2.46	0.48
1:A:326:GLY:HA2	1:A:329:LEU:HD21	1.95	0.48

Continued on next page...



Continued from prev		Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:A:274:MET:HG3	8:A:573:HOH:O	2.14	0.48
1:A:415:ASN:HB3	1:A:431:ASN:HB3	1.96	0.48
1:A:231:TYR:H	1:A:231:TYR:HD1	1.62	0.47
1:A:142:LYS:HG2	8:A:701:HOH:O	2.15	0.47
1:A:331:HIS:CG	1:A:332:PRO:HD2	2.50	0.47
1:A:274:MET:H	1:A:415:ASN:HD22	1.64	0.46
1:A:240:GLU:HB3	1:A:241:PRO:HD2	1.97	0.46
1:A:3:SER:HA	1:A:4:PRO:HD3	1.86	0.45
1:A:23:ASP:O	1:A:26:LEU:CD2	2.65	0.45
1:A:23:ASP:HA	1:A:26:LEU:CD2	2.46	0.45
1:A:314:THR:HG23	1:A:316:TRP:N	2.20	0.44
1:A:4:PRO:O	1:A:5:ASN:HB2	2.17	0.44
1:A:35:LYS:HD3	1:A:35:LYS:HA	1.87	0.44
1:A:126:PHE:HB2	1:A:131:TYR:HB2	2.00	0.43
1:A:269:TRP:O	1:A:272:GLU:HG3	2.17	0.43
1:A:349:GLN:NE2	1:A:354:VAL:HG11	2.32	0.43
1:A:62:TYR:CD1	1:A:62:TYR:N	2.86	0.43
1:A:460:GLY:C	1:A:496:LEU:HD11	2.39	0.43
1:A:274:MET:H	1:A:415:ASN:ND2	2.16	0.43
1:A:194:PHE:O	1:A:231:TYR:HD1	2.02	0.42
1:A:325:VAL:O	1:A:329:LEU:CD2	2.60	0.42
1:A:101:HIS:CD2	1:A:165:LEU:HG	2.54	0.42
1:A:195:ARG:HA	1:A:231:TYR:CD1	2.55	0.41
1:A:438:LEU:HD23	1:A:440:LEU:CD1	2.50	0.41
1:A:33:ALA:HB3	1:A:34:PRO:HD3	2.03	0.41
1:A:54:PRO:HB2	1:A:357:TRP:CE3	2.55	0.41
1:A:430:ASN:O	1:A:486:PRO:HB2	2.20	0.41
1:A:26:LEU:H	1:A:26:LEU:HD22	1.85	0.41
1:A:62:TYR:CD2	7:A:503:HMC:H4	2.56	0.41
1:A:28:CYS:HA	1:A:32:LEU:HB2	2.03	0.40
1:A:218:ASN:OD1	1:A:220:ASN:HB2	2.20	0.40

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	494/496~(100%)	483 (98%)	11 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	418/418 (100%)	402~(96%)	16 (4%)	33 31

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	26	LEU
1	А	125	ASP
1	А	142	LYS
1	А	165	LEU
1	А	208	LYS
1	А	214	LEU
1	А	231	TYR
1	А	243	LYS
1	А	265	VAL
1	А	278	LYS
1	А	323	MET
1	А	329	LEU
1	А	338	VAL
1	А	341	SER
1	А	440	LEU
1	А	466	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:



Mol	Chain	Res	Type
1	А	15	HIS
1	А	105	ASN
1	А	185	HIS
1	А	349	GLN
1	А	350	ASN
1	А	415	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Γ	Mol	Type	Chain	Dog	Link	B	Bond lengths			Bond angles		
	WIOI	Type	Unam	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2	
	1	PCA	А	1	1	5,7,9	1.32	0	7,8,12	1.40	1 (14%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	А	1	1	-	0/0/9/13	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	1	PCA	O-C-CA	-2.17	119.09	124.78

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	$_{\rm ths}$	Bond angles		
	Type	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	В	1	2	12,12,12	0.79	0	$17,\!17,\!17$	0.95	1 (5%)
2	AGL	В	2	2,7	9,10,11	1.12	0	14,14,16	1.31	2 (14%)
3	GLC	С	1	3,7	11,11,12	1.00	1 (9%)	$15,\!15,\!17$	1.16	1 (6%)
3	GLC	С	2	3	11,11,12	0.96	0	$15,\!15,\!17$	0.91	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	В	1	2	-	0/2/22/22	0/1/1/1
2	AGL	В	2	2,7	-	-	0/1/1/1
3	GLC	С	1	3,7	-	0/2/19/22	0/1/1/1
3	GLC	С	2	3	-	1/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	С	1	GLC	C2-C3	2.10	1.55	1.52

All (6) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	С	1	GLC	C1-O5-C5	3.34	116.72	112.19
2	В	2	AGL	C1-C2-C3	2.65	112.93	109.67
3	С	2	GLC	C1-O5-C5	2.35	115.38	112.19
2	В	1	GLC	C1-O5-C5	2.12	117.66	113.66
2	В	2	AGL	C2-C3-C4	-2.06	108.72	110.85
3	С	2	GLC	C2-C3-C4	-2.05	107.35	110.89

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	2	GLC	O5-C5-C6-O6

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bond lengths			Bond angles		
IVIOI	Type	Unam	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
4	NAG	А	497	1	14,14,15	0.77	0	17,19,21	1.12	2 (11%)
7	HMC	А	503	2,3	11,11,12	2.78	4 (36%)	$9,\!15,\!17$	1.15	1 (11%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	А	497	1	-	0/6/23/26	0/1/1/1
7	HMC	А	503	2,3	-	1/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	А	503	HMC	C4-C5	7.41	1.57	1.51
7	А	503	HMC	C2-C3	3.04	1.57	1.52
7	А	503	HMC	C6-C5	2.63	1.38	1.33
7	А	503	HMC	C3-C4	2.14	1.56	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	А	497	NAG	C1-O5-C5	2.76	115.93	112.19
7	А	503	HMC	C1-C6-C5	2.48	128.90	123.20
4	А	497	NAG	C8-C7-N2	-2.02	112.69	116.10

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	А	503	HMC	C6-C5-C7-O7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	А	503	HMC	1	0

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

