

Full wwPDB X-ray Structure Validation Report (i)

May 16, 2020 – 12:39 pm BST

PDB ID : 4CQB

Title : The reaction mechanism of the N-isopropylammelide isopropylaminohydrolas

e AtzC: insights from structural and mutagenesis studies

Authors: Balotra, S.; Newman, J.; French, N.G.; Peat, T.S.; Scott, C.

Deposited on : 2014-02-13

Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.11

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove) roteins) : Engh & Huber (2001)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	4003 (1.86-1.82)
Clashscore	141614	4233 (1.86-1.82)
Ramachandran outliers	138981	4185 (1.86-1.82)
Sidechain outliers	138945	4186 (1.86-1.82)
RSRZ outliers	127900	3957 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	423	90%	5%	5%
1	В	423	88%	6%	5%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6923 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called N-ISOPROPYLAMMELIDE ISOPROPYL AMIDOHYDRO-LASE.

Mol	Chain	Residues	${f Atoms}$			ZeroOcc	AltConf	Trace		
1	A	402	${ m Total}\ 3205$	C 2031	N 541	O 620	S 13	0	6	0
1	В	400	Total 3187	C 2021	N 537	O 616	S 13	0	6	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP O52063
A	-18	GLY	-	expression tag	UNP O52063
A	-17	SER	-	expression tag	UNP O52063
A	-16	SER	-	expression tag	UNP O52063
A	-15	HIS	-	expression tag	UNP O52063
A	-14	HIS	-	expression tag	UNP O52063
A	-13	HIS	-	expression tag	UNP O52063
A	-12	HIS	-	expression tag	UNP O52063
A	-11	HIS	_	expression tag	UNP O52063
A	-10	HIS	-	expression tag	UNP O52063
A	-9	SER	-	expression tag	UNP O52063
A	-8	SER	_	expression tag	UNP O52063
A	-7	GLY	-	expression tag	UNP O52063
A	-6	LEU	_	expression tag	UNP O52063
A	-5	VAL	-	expression tag	UNP O52063
A	-4	PRO	-	expression tag	UNP O52063
A	-3	ARG	-	expression tag	UNP O52063
A	-2	GLY	-	expression tag	UNP O52063
A	-1	SER	-	expression tag	UNP O52063
A	0	HIS	-	expression tag	UNP O52063
В	-19	MET		expression tag	UNP O52063
В	-18	GLY	=	expression tag	UNP O52063
В	-17	SER	-	expression tag	UNP O52063
В	-16	SER	-	expression tag	UNP O52063

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	-15	HIS	=	expression tag	UNP O52063
В	-14	HIS	-	expression tag	UNP O52063
В	-13	HIS	_	expression tag	UNP O52063
В	-12	HIS	-	expression tag	UNP O52063
В	-11	HIS	-	expression tag	UNP O52063
В	-10	HIS	-	expression tag	UNP O52063
В	-9	SER	-	expression tag	UNP O52063
В	-8	SER	=	expression tag	UNP O52063
В	-7	GLY	-	expression tag	UNP O52063
В	-6	LEU	-	expression tag	UNP O52063
В	-5	VAL	-	expression tag	UNP O52063
В	-4	PRO	-	expression tag	UNP O52063
В	-3	ARG	=	expression tag	UNP O52063
В	-2	GLY	=	expression tag	UNP O52063
В	-1	SER	-	expression tag	UNP O52063
В	0	HIS	=	expression tag	UNP O52063

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
3	В	1	Total Cl 1 1	0	0
3	A	1	Total Cl 1 1	0	0


• Molecule 4 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O S 4 2 1 1	0	0
4	В	1	Total C O S 8 4 2 2	0	1
4	В	1	Total C O S 4 2 1 1	0	0

 \bullet Molecule 5 is MALONATE ION (three-letter code: MLI) (formula: $\mathrm{C_3H_2O_4}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 7 3 4	0	0

Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 7 3 4	0	0
5	В	1	Total C O 7 3 4	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	241	Total O 241 241	0	0
6	В	249	Total O 249 249	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: N-ISOPROPYLAMMELIDE ISOPROPYL AMIDOHYDROLASE



• Molecule 1: N-ISOPROPYLAMMELIDE ISOPROPYL AMIDOHYDROLASE





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	106.47Å 86.73Å 114.17Å	Danagitan
a, b, c, α , β , γ	90.00° 104.69° 90.00°	Depositor
Resolution (Å)	40.40 - 1.84	Depositor
Resolution (A)	40.37 - 1.84	EDS
% Data completeness	99.2 (40.40-1.84)	Depositor
(in resolution range)	99.2 (40.37-1.84)	EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.30 (at 1.84Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
D D.	0.191 , 0.229	Depositor
R, R_{free}	0.200 , 0.234	DCC
R_{free} test set	4338 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	18.2	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.39, 50.0	EDS
L-test for twinning ²	$ < L > = 0.45, < L^2> = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6923	wwPDB-VP
Average B, all atoms $(Å^2)$	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 11.44% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MLI, DMS, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	Bond lengths		nd angles
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5
1	A	0.93	0/3263	0.91	4/4414 (0.1%)
1	В	0.97	0/3245	0.93	$2/4391 \ (0.0\%)$
All	All	0.95	0/6508	0.92	6/8805 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
1	В	231	ARG	NE-CZ-NH1	6.40	123.50	120.30
1	A	366	ASP	CB-CG-OD1	5.98	123.68	118.30
1	В	366	ASP	CB-CG-OD1	5.55	123.30	118.30
1	A	231	ARG	NE-CZ-NH1	5.41	123.00	120.30
1	A	220	ASP	CB-CG-OD2	-5.13	113.68	118.30
1	A	183	LEU	CB-CG-CD2	5.08	119.64	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	82	TYR	Peptide



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Α	3205	0	3193	11	0
1	В	3187	0	3171	14	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
3	A	1	0	0	1	0
3	В	1	0	0	1	0
4	A	4	0	6	2	0
4	В	12	0	18	3	0
5	A	14	0	4	0	0
5	В	7	0	2	0	0
6	A	241	0	0	8	0
6	В	249	0	0	6	1
All	All	6923	0	6394	32	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} ({\rm \AA}) \end{array}$	Clash overlap (Å)
3:A:1405:CL:CL	6:A:2210:HOH:O	1.93	1.24
3:B:1405:CL:CL	6:B:2247:HOH:O	2.27	0.89
1:A:102:GLU:OE1	6:A:2064:HOH:O	1.99	0.80
1:B:4:ASP:OD2	6:B:2002:HOH:O	2.03	0.77
1:B:234:GLN:OE1	6:B:2163:HOH:O	2.06	0.72
1:A:259[B]:GLU:O	1:A:263[B]:GLU:HG3	1.95	0.66
1:B:102:GLU:OE1	6:B:2071:HOH:O	2.16	0.63
1:A:174:ARG:NH1	1:A:209:GLU:OE1	2.36	0.59
6:A:2215:HOH:O	4:B:1406[A]:DMS:C2	2.51	0.58
1:B:174:ARG:NH1	1:B:209:GLU:OE1	2.38	0.56
6:A:2215:HOH:O	4:B:1406[A]:DMS:H22	2.07	0.54
1:B:4:ASP:OD2	1:B:41:VAL:HG12	2.12	0.50
1:A:41:VAL:HG11	1:A:44:GLU:OE2	2.12	0.49
4:A:1406:DMS:C1	6:A:2181:HOH:O	2.60	0.49
1:B:293:GLU:OE2	1:B:334:ARG:NH2	2.45	0.49

Continued on next page...



 $Continued\ from\ previous\ page...$

Atom-1	Atom-2	$egin{array}{l} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	Clash overlap (Å)
1:A:171[C]:SER:HA	1:A:174:ARG:HD2	1.95	0.48
1:B:105:ARG:HG3	6:B:2078:HOH:O	2.11	0.48
1:A:171[A]:SER:HA	1:A:174:ARG:HD2	1.96	0.47
1:A:171[B]:SER:HA	1:A:174:ARG:HD2	1.96	0.47
1:B:170:GLU:O	1:B:174:ARG:HG3	2.14	0.47
1:A:105:ARG:HG3	6:A:2072:HOH:O	2.14	0.46
1:B:171:SER:HA	1:B:174:ARG:HD2	1.96	0.46
6:A:2215:HOH:O	4:B:1406[A]:DMS:H21	2.15	0.45
1:A:170:GLU:O	1:A:174:ARG:HG3	2.17	0.45
1:B:156:VAL:HG22	1:B:183:LEU:HB2	1.98	0.45
1:A:340:TRP:CG	1:A:375:PRO:HB3	2.53	0.44
1:B:397:LYS:HA	1:B:397:LYS:HD2	1.77	0.44
1:B:145:GLU:HG2	6:B:2105:HOH:O	2.19	0.43
1:B:342:MET:HA	1:B:346:GLU:HB2	2.02	0.41
1:A:156:VAL:HG22	1:A:183:LEU:HB2	2.03	0.41
4:A:1406:DMS:H12	6:A:2181:HOH:O	2.21	0.40
1:B:340:TRP:CG	1:B:375:PRO:HB3	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{aligned}$	$egin{array}{c} ext{Clash} \ ext{overlap } (ext{Å}) \end{array}$
6:B:2134:HOH:O	6:B:2134:HOH:O[2_658]	2.09	0.11

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	\mathbf{s}
1	A	407/423 (96%)	392 (96%)	14 (3%)	1 (0%)	47 33	
1	В	404/423 (96%)	390 (96%)	13 (3%)	1 (0%)	47 33	
All	All	811/846 (96%)	782 (96%)	27 (3%)	2 (0%)	41 33	



All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	249	HIS
1	В	249	HIS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	${f Analysed}$	Rotameric	Outliers	Percentil	$\mathbf{e}\mathbf{s}$
1	A	$351/362 \ (97\%)$	347 (99%)	4 (1%)	73 64	
1	В	$348/362 \ (96\%)$	345 (99%)	3 (1%)	78 71	
All	All	699/724~(96%)	692 (99%)	7 (1%)	76 68	

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	${f Res}$	\mathbf{Type}
1	A	80	ARG
1	A	216	TYR
1	A	241	TYR
1	A	349	ARG
1	В	80	ARG
1	В	216	TYR
1	В	241	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	GLN
1	A	194	ASN
1	В	114	GLN
1	В	234	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Chain	Dec	Link	Bond lengths			Bond angles			
MIGI	$egin{array}{c c c c c c c c c c c c c c c c c c c $	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
5	MLI	A	1407	_	0,6,6	0.00	-	0,7,7	0.00	-
4	DMS	В	1406[A]	-	3,3,3	0.65	0	3,3,3	0.49	0
4	DMS	В	1407	-	3,3,3	0.76	0	3,3,3	1.05	0
4	DMS	В	1406[B]	-	3,3,3	0.54	0	3,3,3	0.70	0
5	MLI	A	1408	-	0,6,6	0.00	-	0,7,7	0.00	-
4	DMS	A	1406	-	3,3,3	0.66	0	3,3,3	0.26	0
5	MLI	В	1408	-	0,6,6	0.00	-	0,7,7	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	${ m Res}$	Link	Chirals	Torsions	Rings
5	MLI	A	1408	-	-	0/0/4/4	-
5	MLI	A	1407	-	-	0/0/4/4	-
5	MLI	В	1408	-	=	0/0/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	1406[A]	DMS	3	0
4	A	1406	DMS	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$RZ> \hspace{0.2cm} \#RSRZ{>}2$		$OWAB(Å^2)$	Q < 0.9
1	A	$402/423 \ (95\%)$	-0.08	4 (0%)	82 82	10, 18, 34, 53	0
1	В	$400/423 \ (94\%)$	-0.19	2 (0%)	91 91	9, 16, 30, 42	0
All	All	802/846 (94%)	-0.13	6 (0%)	87 87	9, 17, 33, 53	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	403	ALA	4.5
1	В	36	LYS	3.0
1	A	398	ASP	2.6
1	A	36	LYS	2.2
1	A	35	ALA	2.1
1	В	39	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
5	MLI	A	1408	7/7	0.85	0.14	24,34,40,49	0
3	CL	В	1405	1/1	0.88	0.08	39,39,39,39	0
4	DMS	В	1407	4/4	0.91	0.12	26,35,40,45	0
4	DMS	В	1406[A]	4/4	0.95	0.12	16,18,20,23	4
4	DMS	В	1406[B]	4/4	0.95	0.12	27,28,33,38	4
4	DMS	A	1406	4/4	0.95	0.17	20,28,34,35	0
5	MLI	A	1407	7/7	0.96	0.08	9,11,13,14	0
5	MLI	В	1408	7/7	0.97	0.09	10,12,13,13	0
3	CL	A	1405	1/1	0.99	0.04	33,33,33,33	0
2	ZN	В	1404	1/1	1.00	0.04	16,16,16,16	0
2	ZN	A	1404	1/1	1.00	0.05	17,17,17,17	0

6.5 Other polymers (i)

There are no such residues in this entry.

