



Full wwPDB NMR Structure Validation Report ⓘ

Feb 8, 2022 – 03:09 PM EST

PDB ID : 1CR8
Title : LOW DENSITY LIPOPROTEIN RECEPTOR-RELATED PROTEIN COMPLEMENT REPEAT 8
Authors : Huang, W.; Dolmer, K.; Gettins, P.G.W.
Deposited on : 1998-12-14

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.26
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.26


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	42	

2 Ensemble composition and analysis i

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:6-A:39 (34)	0.27	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 4 single-model clusters were found.

Cluster number	Models
1	3, 8, 11, 12, 13
2	1, 4, 9, 10, 15
3	5, 6, 7, 14
4	16, 19
Single-model clusters	2; 17; 18; 20

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 595 atoms, of which 276 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN).

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	42	594	187	276	56	68	7	0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	
			Total	Ca
2	A	1	1	1

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.2 Score per residue for model 2

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.3 Score per residue for model 3

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.4 Score per residue for model 4

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.5 Score per residue for model 5

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.6 Score per residue for model 6

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.7 Score per residue for model 7

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.8 Score per residue for model 8 (medoid)

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.9 Score per residue for model 9

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.10 Score per residue for model 10

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.11 Score per residue for model 11

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.12 Score per residue for model 12

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.13 Score per residue for model 13

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.14 Score per residue for model 14

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.15 Score per residue for model 15

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.16 Score per residue for model 16

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.17 Score per residue for model 17

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.18 Score per residue for model 18

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)




4.2.19 Score per residue for model 19

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)



4.2.20 Score per residue for model 20

- Molecule 1: PROTEIN (LOW DENSITY LIPOPROTEIN RECEPTOR RELATED PROTEIN)



5 Refinement protocol and experimental data overview

The models were refined using the following method: *DYNAMICS ANNEALING*.

Of the 20 calculated structures, 20 were deposited, based on the following criterion: *LEAST RESTRAINT VIOLATION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	refinement	
DYANA	structure solution	

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.65±0.01	0±0/276 (0.0± 0.0%)	1.84±0.09	6±1/372 (1.5± 0.3%)
All	All	0.65	0/5520 (0.0%)	1.84	113/7440 (1.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.7±0.7
All	All	0	34

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	31	ASP	CB-CG-OD1	19.35	135.71	118.30	20	19
1	A	38	GLU	OE1-CD-OE2	-18.11	101.57	123.30	17	20
1	A	31	ASP	CB-CG-OD2	15.61	132.35	118.30	14	13
1	A	31	ASP	OD1-CG-OD2	-11.34	101.76	123.30	20	19
1	A	23	ARG	NE-CZ-NH1	10.40	125.50	120.30	20	4
1	A	14	ARG	NE-CZ-NH1	8.33	124.46	120.30	20	8
1	A	25	ARG	NE-CZ-NH1	7.69	124.14	120.30	11	9
1	A	38	GLU	CG-CD-OE2	7.20	132.71	118.30	2	3
1	A	27	ASP	CB-CG-OD1	-6.10	112.81	118.30	5	7
1	A	38	GLU	CG-CD-OE1	5.43	129.15	118.30	18	1
1	A	24	TRP	O-C-N	-5.35	114.14	122.70	3	2
1	A	6	CYS	C-N-CA	5.30	134.95	121.70	14	5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	27	ASP	CB-CG-OD2	-5.26	113.57	118.30	7	1
1	A	8	THR	C-N-CA	5.13	134.54	121.70	2	1
1	A	32	CYS	CA-CB-SG	5.05	123.09	114.00	14	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	36	SER	Peptide	13
1	A	29	ASP	Peptide	6
1	A	31	ASP	Peptide	4
1	A	22	LEU	Peptide	3
1	A	37	ASP	Peptide	2
1	A	23	ARG	Sidechain	2
1	A	24	TRP	Peptide	1
1	A	28	GLY	Peptide	1
1	A	25	ARG	Sidechain	1
1	A	10	GLU	Peptide	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	271	236	236	0±0
All	All	5440	4720	4720	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:23:ARG:HD2	1:A:23:ARG:H	0.48	1.67	20	1
1:A:11:PHE:CG	1:A:25:ARG:HD2	0.48	2.43	10	2
1:A:11:PHE:CG	1:A:25:ARG:HD3	0.44	2.48	19	4

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:34:ASP:OD1	1:A:36:SER:OG	0.41	2.38	12	1
1:A:32:CYS:SG	1:A:33:MET:N	0.40	2.94	5	1

6.3 Torsion angles [\(i\)](#)

6.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	34/42 (81%)	20±1 (60±4%)	9±2 (28±5%)	4±1 (13±4%)	1	6
All	All	680/840 (81%)	405 (60%)	189 (28%)	86 (13%)	1	6

All 9 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	35	SER	20
1	A	39	LYS	14
1	A	23	ARG	14
1	A	8	THR	13
1	A	26	CYS	12
1	A	33	MET	10
1	A	9	ASP	1
1	A	34	ASP	1
1	A	25	ARG	1

6.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	32/37 (86%)	24±3 (74±9%)	8±3 (26±9%)	2	24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	640/740 (86%)	476 (74%)	164 (26%)	2 24

All 22 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	10	GLU	20
1	A	7	HIS	13
1	A	15	LEU	13
1	A	27	ASP	12
1	A	38	GLU	12
1	A	14	ARG	11
1	A	18	LEU	11
1	A	32	CYS	11
1	A	39	LYS	10
1	A	16	ASP	7
1	A	26	CYS	7
1	A	34	ASP	6
1	A	13	CYS	6
1	A	31	ASP	5
1	A	23	ARG	5
1	A	33	MET	4
1	A	37	ASP	4
1	A	12	GLN	3
1	A	36	SER	1
1	A	20	ILE	1
1	A	25	ARG	1
1	A	22	LEU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided