

wwPDB X-ray Structure Validation Summary Report (i)

Jan 27, 2024 – 07:24 PM EST

PDB ID : 1CTS

Title: CRYSTALLOGRAPHIC REFINEMENT AND ATOMIC MODELS OF

TWO DIFFERENT FORMS OF CITRATE SYNTHASE AT 2.7 AND 1.7

ANGSTROMS RESOLUTION

Authors : Remington, S.; Wiegand, G.; Huber, R.

Deposited on : 1984-01-27

Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quali	ty of chain	
1	Δ	437	47%	37%	13% •



2 Entry composition (i)


There are 2 unique types of molecules in this entry. The entry contains 3457 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called CITRATE SYNTHASE.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	437	Total 3444	C 2200	N 591	O 634	S 19	132	0	0

• Molecule 2 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
2	A	1	Total 13	C 6	O 7	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: CITRATE SYNTHASE





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 41 21 2	Depositor	
Cell constants	77.40Å 77.40Å 196.40Å	Depositor	
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	5.00 - 2.70	Depositor	
% Data completeness	(Not available) (5.00-2.70)	Depositor	
(in resolution range)	(1101 available) (9.00 2.10)	Depositor	
R_{merge}	(Not available)	Depositor	
R_{sym}	(Not available)	Depositor	
Refinement program	EREF	Depositor	
R, R_{free}	0.183 , (Not available)	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	3457	wwPDB-VP	
Average B, all atoms (Å ²)	23.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Во	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5
1	A	1.12	$11/3528 \ (0.3\%)$	1.51	28/4786 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	73

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
1	A	411	TRP	NE1-CE2	-8.31	1.26	1.37
1	A	209	TRP	NE1-CE2	-7.88	1.27	1.37
1	A	306	TRP	NE1-CE2	-7.71	1.27	1.37
1	A	94	TRP	NE1-CE2	-7.71	1.27	1.37
1	A	114	TRP	NE1-CE2	-7.64	1.27	1.37

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathrm{Ideal}(^{o})$
1	A	156	ARG	NE-CZ-NH1	-10.42	115.09	120.30
1	A	195	ARG	NE-CZ-NH2	-8.60	116.00	120.30
1	A	25	ARG	NE-CZ-NH1	-8.18	116.21	120.30
1	A	103	THR	N-CA-CB	-7.88	95.32	110.30
1	A	156	ARG	NE-CZ-NH2	7.87	124.24	120.30

All (1) chirality outliers are listed below:



Mol	Chain	Res	Type	Atom
1	A	71	ILE	CA

5 of 73 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	27	GLN	Sidechain
1	A	28	HIS	Mainchain
1	A	31	THR	Mainchain
1	A	39	ASP	Sidechain
1	A	4	THR	Mainchain, Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3444	0	3436	133	5
2	A	13	0	5	2	0
All	All	3457	0	3441	134	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 134 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:A:125:VAL:HG13	1:A:188:LYS:HE2	1.46	0.93
1:A:305:ILE:HD11	1:A:353:LEU:HD12	1.54	0.90
1:A:305:ILE:HD13	1:A:357:VAL:HG22	1.54	0.87
1:A:21:ILE:HD12	1:A:415:LEU:HD22	1.55	0.85
1:A:86:GLU:HG2	1:A:230:LEU:HB2	1.56	0.85

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:31:THR:CG2	1:A:434:VAL:CG2[8_665]	1.46	0.74

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	Clash overlap (Å)
1:A:127:MET:CE	1:A:127:MET:CE[8_665]	1.73	0.47
1:A:54:GLU:OE2	1:A:427:THR:OG1[8_665]	1.83	0.37
1:A:139:SER:OG	1:A:149:ASN:ND2[8_665]	1.84	0.36
1:A:31:THR:CB	1:A:434:VAL:CG2[8_665]	1.86	0.34

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	435/437 (100%)	382 (88%)	42 (10%)	11 (2%)	5 14

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	6	LEU
1	A	46	ARG
1	A	67	ARG
1	A	71	ILE
1	A	164	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	371/371 (100%)	321 (86%)	50 (14%)	4 9	

5 of 50 residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	A	285	LEU
1	A	314	VAL
1	A	432	LYS
1	A	292	VAL
1	A	308	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	192	ASN
1	A	211	HIS
1	A	352	GLN
1	A	267	ASN
1	A	307	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link		nd leng	ths	В	ond ang	les
IVIOI	туре	Chain	rtes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CIT	A	438	-	12,12,12	0.99	0	17,17,17	1.69	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	A	438	-	-	3/16/16/16	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	438	CIT	O7-C3-C6	3.69	114.05	108.86
2	A	438	CIT	C3-C4-C5	-3.04	106.44	113.81
2	A	438	CIT	O6-C6-C3	2.56	117.50	113.05
2	A	438	CIT	O6-C6-O5	-2.27	116.59	123.82
2	A	438	CIT	O4-C5-O3	-2.25	117.68	123.30

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	438	CIT	C2-C3-C6-O5
2	A	438	CIT	O7-C3-C6-O5
2	A	438	CIT	C4-C3-C6-O5

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	438	CIT	2	0

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

