



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 3, 2024 – 07:16 AM EST

PDB ID : 6CUT
Title : Engineered Holo TrpB from *Pyrococcus furiosus*, PfTrpB7E6 with (2S,3S)-iso propylserine bound as the external aldimine
Authors : Boville, C.E.; Scheele, R.A.; Buller, A.R.; Arnold, F.H.
Deposited on : 2018-03-26
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	
1	B	388	
1	C	388	
1	D	388	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 11972 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophan synthase beta chain 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	385	2860	1822	492	534	12	0	0	0
1	B	385	2974	1897	517	546	14	0	12	0
1	C	377	2768	1772	476	509	11	0	0	0
1	D	385	2877	1833	496	536	12	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	16	VAL	ILE	engineered mutation	UNP Q8U093
A	17	GLY	GLU	engineered mutation	UNP Q8U093
A	91	PRO	LEU	engineered mutation	UNP Q8U093
A	95	LEU	PHE	engineered mutation	UNP Q8U093
A	161	ALA	LEU	engineered mutation	UNP Q8U093
A	173	GLU	VAL	engineered mutation	UNP Q8U093
A	274	LEU	PHE	engineered mutation	UNP Q8U093
A	292	SER	THR	engineered mutation	UNP Q8U093
A	384	ALA	VAL	engineered mutation	UNP Q8U093
B	16	VAL	ILE	engineered mutation	UNP Q8U093
B	17	GLY	GLU	engineered mutation	UNP Q8U093
B	91	PRO	LEU	engineered mutation	UNP Q8U093
B	95	LEU	PHE	engineered mutation	UNP Q8U093
B	161	ALA	LEU	engineered mutation	UNP Q8U093
B	173	GLU	VAL	engineered mutation	UNP Q8U093
B	274	LEU	PHE	engineered mutation	UNP Q8U093
B	292	SER	THR	engineered mutation	UNP Q8U093
B	384	ALA	VAL	engineered mutation	UNP Q8U093
C	16	VAL	ILE	engineered mutation	UNP Q8U093
C	17	GLY	GLU	engineered mutation	UNP Q8U093
C	91	PRO	LEU	engineered mutation	UNP Q8U093

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	D	1	Total	C	N	O	P	0	0
			25	14	2	8	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		
3	B	1	Total	Na	0	0
			1	1		
3	C	1	Total	Na	0	0
			1	1		
3	D	1	Total	Na	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	85	Total	O	0	0
			85	85		
4	B	112	Total	O	0	0
			112	112		
4	C	85	Total	O	0	0
			85	85		
4	D	107	Total	O	0	0
			107	107		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tryptophan synthase beta chain 1



- Molecule 1: Tryptophan synthase beta chain 1




- Molecule 1: Tryptophan synthase beta chain 1



- Molecule 1: Tryptophan synthase beta chain 1





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	82.22Å 107.43Å 159.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.77 39.16 – 1.77	Depositor EDS
% Data completeness (in resolution range)	99.5 (40.00-1.77) 99.5 (39.16-1.77)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 1.77Å)	Xtrriage
Refinement program	REFMAC 5.8.0218	Depositor
R, R_{free}	0.193 , 0.225 0.200 , 0.226	Depositor DCC
R_{free} test set	6812 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	35.7	Xtrriage
Anisotropy	0.255	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11972	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, FEJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/2920	0.50	0/3964
1	B	0.30	0/3035	0.51	0/4108
1	C	0.30	0/2827	0.50	0/3842
1	D	0.30	0/2937	0.50	0/3983
All	All	0.30	0/11719	0.51	0/15897

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2860	0	2778	5	0
1	B	2974	0	2931	13	0
1	C	2768	0	2671	3	0
1	D	2877	0	2810	2	0
2	A	25	0	0	0	0
2	B	25	0	0	0	0
2	C	25	0	0	0	0
2	D	25	0	0	0	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	85	0	0	0	0
4	B	112	0	0	0	0
4	C	85	0	0	0	0
4	D	107	0	0	0	0
All	All	11972	0	11190	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:137[B]:GLN:HB3	1:B:140:ASN:HD22	1.59	0.66
1:B:136[B]:ARG:O	1:B:137[B]:GLN:HG2	2.02	0.59
1:B:136[B]:ARG:C	1:B:137[B]:GLN:HG2	2.28	0.54
1:D:15:LEU:HD21	1:D:173:GLU:HA	1.90	0.54
1:B:131[B]:ALA:O	1:B:134[B]:VAL:HB	2.09	0.53
1:A:211:ILE:HG21	1:A:219:PRO:HD3	1.92	0.52
1:B:211:ILE:HG21	1:B:219:PRO:HD3	1.92	0.51
1:B:63:ILE:HD11	1:B:354:MET:HG2	1.94	0.48
1:B:134[B]:VAL:O	1:B:138[B]:LYS:HA	2.14	0.47
1:A:119:ALA:O	1:B:58:ARG:NH2	2.46	0.47
1:C:188:GLY:HA2	1:C:275:HIS:O	2.16	0.46
1:C:18:PRO:HG3	1:C:176:PHE:CD1	2.52	0.45
1:A:188:GLY:HA2	1:A:275:HIS:O	2.18	0.44
1:D:211:ILE:HG21	1:D:219:PRO:HD3	1.98	0.44
1:B:274:LEU:CD2	1:B:289:ILE:HD12	2.49	0.43
1:A:90:ALA:HB3	1:A:91:PRO:HD3	2.00	0.42
1:B:154:VAL:HG13	1:B:167:GLU:HG3	2.02	0.41
1:C:106:GLY:O	1:C:133:ASP:HB3	2.19	0.41
1:B:274:LEU:HD22	1:B:289:ILE:HD12	2.03	0.41
1:A:63:ILE:HD11	1:A:354:MET:HG2	2.03	0.40
1:B:110:HIS:CE1	1:B:184:GLY:HA2	2.57	0.40
1:B:289:ILE:HD11	1:B:302:PRO:HG2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	383/388 (99%)	377 (98%)	6 (2%)	0	100	100
1	B	395/388 (102%)	385 (98%)	9 (2%)	1 (0%)	41	25
1	C	373/388 (96%)	369 (99%)	3 (1%)	1 (0%)	41	25
1	D	383/388 (99%)	378 (99%)	4 (1%)	1 (0%)	41	25
All	All	1534/1552 (99%)	1509 (98%)	22 (1%)	3 (0%)	47	32

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	186	VAL
1	D	186	VAL
1	C	186	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	279/305 (92%)	279 (100%)	0	100	100
1	B	291/305 (95%)	288 (99%)	3 (1%)	76	68
1	C	263/305 (86%)	263 (100%)	0	100	100
1	D	282/305 (92%)	281 (100%)	1 (0%)	91	88
All	All	1115/1220 (91%)	1111 (100%)	4 (0%)	92	88

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	137[A]	GLN
1	B	137[B]	GLN
1	B	290	LYS
1	D	13	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	140	ASN
1	D	283	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FEJ	D	401	-	23,25,25	2.13	6 (26%)	32,36,36	1.39	3 (9%)
2	FEJ	A	401	-	23,25,25	2.06	6 (26%)	32,36,36	1.33	3 (9%)
2	FEJ	B	401	-	23,25,25	2.04	5 (21%)	32,36,36	1.40	4 (12%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FEJ	C	401	-	23,25,25	2.06	5 (21%)	32,36,36	1.36	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FEJ	D	401	-	-	5/23/23/23	0/1/1/1
2	FEJ	A	401	-	-	7/23/23/23	0/1/1/1
2	FEJ	B	401	-	-	6/23/23/23	0/1/1/1
2	FEJ	C	401	-	-	5/23/23/23	0/1/1/1

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	FEJ	CAA-CAS	-6.01	1.40	1.50
2	D	401	FEJ	CAA-CAS	-5.84	1.40	1.50
2	C	401	FEJ	CAA-CAS	-5.82	1.40	1.50
2	B	401	FEJ	CAA-CAS	-5.74	1.40	1.50
2	B	401	FEJ	CAL-N	4.43	1.35	1.27
2	C	401	FEJ	CAL-N	4.15	1.35	1.27
2	A	401	FEJ	CAL-N	3.92	1.34	1.27
2	C	401	FEJ	CAB-CAM	-3.72	1.40	1.50
2	D	401	FEJ	CAL-N	3.68	1.34	1.27
2	D	401	FEJ	CAB-CAM	-3.62	1.41	1.50
2	B	401	FEJ	CAB-CAM	-3.51	1.41	1.50
2	A	401	FEJ	CAB-CAM	-3.43	1.41	1.50
2	D	401	FEJ	PAP-OAQ	3.09	1.60	1.50
2	D	401	FEJ	CAN-CAL	-3.09	1.40	1.46
2	B	401	FEJ	CAO-NAR	2.87	1.40	1.34
2	C	401	FEJ	CAN-CAL	-2.86	1.41	1.46
2	C	401	FEJ	CAO-NAR	2.86	1.40	1.34
2	A	401	FEJ	CAO-NAR	2.84	1.40	1.34
2	D	401	FEJ	CAO-NAR	2.84	1.40	1.34
2	A	401	FEJ	CAN-CAL	-2.83	1.41	1.46
2	B	401	FEJ	CAN-CAL	-2.54	1.41	1.46
2	A	401	FEJ	CG-CB	2.27	1.56	1.53

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	FEJ	CAN-CAT-CAS	4.68	123.08	120.19
2	C	401	FEJ	CAN-CAT-CAS	4.55	123.00	120.19
2	A	401	FEJ	CAN-CAT-CAS	4.49	122.96	120.19
2	D	401	FEJ	CAN-CAT-CAS	4.45	122.94	120.19
2	D	401	FEJ	OAE-PAP-OAD	3.05	119.28	107.64
2	B	401	FEJ	CAT-CAN-CAM	-2.23	116.55	118.26
2	A	401	FEJ	OAE-PAP-OAQ	2.21	119.32	110.68
2	C	401	FEJ	OAE-PAP-OAQ	2.12	118.97	110.68
2	B	401	FEJ	OAD-PAP-OAC	-2.11	101.11	106.73
2	D	401	FEJ	CB-CA-N	2.07	115.74	109.27
2	B	401	FEJ	OAE-PAP-OAQ	2.06	118.75	110.68
2	A	401	FEJ	CA-N-CAL	2.00	123.29	118.47

There are no chirality outliers.


All (23) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	FEJ	C-CA-N-CAL
2	A	401	FEJ	CB-CA-N-CAL
2	A	401	FEJ	OXT-C-CA-CB
2	A	401	FEJ	O-C-CA-CB
2	B	401	FEJ	C-CA-N-CAL
2	B	401	FEJ	CB-CA-N-CAL
2	B	401	FEJ	N-CA-CB-OG
2	B	401	FEJ	N-CA-CB-CG
2	C	401	FEJ	C-CA-N-CAL
2	C	401	FEJ	CB-CA-N-CAL
2	C	401	FEJ	OXT-C-CA-CB
2	C	401	FEJ	N-CA-CB-OG
2	C	401	FEJ	N-CA-CB-CG
2	D	401	FEJ	C-CA-N-CAL
2	D	401	FEJ	CB-CA-N-CAL
2	D	401	FEJ	OXT-C-CA-CB
2	D	401	FEJ	O-C-CA-CB
2	A	401	FEJ	CA-CB-CG-CD1
2	A	401	FEJ	OG-CB-CG-CD1
2	B	401	FEJ	OXT-C-CA-N
2	B	401	FEJ	OXT-C-CA-CB
2	A	401	FEJ	CAB-OAC-PAP-OAQ
2	D	401	FEJ	N-CA-CB-CG


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	385/388 (99%)	0.50	33 (8%) 10 10	30, 44, 64, 80	0
1	B	385/388 (99%)	0.55	35 (9%) 9 8	26, 40, 65, 87	0
1	C	377/388 (97%)	0.76	56 (14%) 2 2	27, 44, 73, 85	0
1	D	385/388 (99%)	0.47	40 (10%) 6 6	29, 42, 61, 85	0
All	All	1532/1552 (98%)	0.57	164 (10%) 6 5	26, 42, 67, 87	0

All (164) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	286	GLU	6.4
1	C	176	PHE	6.3
1	C	174	ALA	6.2
1	A	384	ALA	6.2
1	B	289	ILE	6.1
1	D	285	GLU	5.6
1	C	178	TYR	5.5
1	C	268	ALA	5.4
1	B	284	ASP	4.7
1	C	153	PRO	4.7
1	B	158	SER	4.5
1	C	76	VAL	4.4
1	D	284	ASP	4.4
1	C	14	THR	4.4
1	B	285	GLU	4.3
1	C	291	PRO	4.3
1	C	75	LEU	4.2
1	D	76	VAL	4.2
1	D	142	PHE	4.2
1	B	157	GLY	4.2
1	B	75	LEU	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	291	PRO	4.1
1	D	176	PHE	4.1
1	D	258	GLU	4.1
1	B	260	GLY	4.0
1	A	157	GLY	4.0
1	D	158	SER	3.8
1	D	178	TYR	3.7
1	D	287	GLY	3.7
1	C	289	ILE	3.7
1	B	156	SER	3.6
1	B	286	GLU	3.6
1	C	160	THR	3.6
1	C	147	LEU	3.5
1	B	76	VAL	3.5
1	C	172	TRP	3.5
1	C	142	PHE	3.4
1	C	80	ALA	3.4
1	A	75	LEU	3.4
1	B	139[A]	MET	3.4
1	C	31	ASP	3.4
1	B	79	GLY	3.3
1	C	382	LEU	3.3
1	D	75	LEU	3.3
1	D	112	VAL	3.3
1	A	155	ASN	3.3
1	A	142	PHE	3.3
1	C	16	VAL	3.3
1	B	340	ILE	3.3
1	C	373	ARG	3.2
1	B	287	GLY	3.2
1	C	78	GLY	3.2
1	D	340	ILE	3.2
1	B	164	ALA	3.2
1	D	341	ILE	3.2
1	A	257	LEU	3.2
1	C	95	LEU	3.1
1	C	257	LEU	3.1
1	D	80	ALA	3.1
1	D	384	ALA	3.1
1	C	163	ASP	3.1
1	C	152	ILE	3.0
1	A	80	ALA	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	160	THR	3.0
1	C	79	GLY	3.0
1	D	79	GLY	3.0
1	B	80	ALA	3.0
1	C	173	GLU	3.0
1	B	78	GLY	2.9
1	D	372	GLY	2.9
1	A	76	VAL	2.9
1	D	288	GLN	2.9
1	C	96	MET	2.9
1	C	113	ALA	2.9
1	B	291	PRO	2.8
1	A	385	SER	2.8
1	D	170	ARG	2.8
1	C	151	VAL	2.8
1	D	291	PRO	2.8
1	D	156	SER	2.8
1	D	78	GLY	2.8
1	B	154	VAL	2.8
1	A	284	ASP	2.7
1	A	112	VAL	2.7
1	C	114	THR	2.7
1	A	78	GLY	2.6
1	C	21	GLU	2.6
1	B	374	GLY	2.6
1	D	157	GLY	2.6
1	A	293	HIS	2.6
1	A	79	GLY	2.6
1	B	341	ILE	2.6
1	D	169	LEU	2.6
1	D	373	ARG	2.5
1	D	154	VAL	2.5
1	A	33	GLU	2.5
1	A	325	GLU	2.5
1	B	372	GLY	2.5
1	C	258	GLU	2.5
1	A	372	GLY	2.5
1	B	174	ALA	2.5
1	A	156	SER	2.5
1	C	186	VAL	2.5
1	A	373	ARG	2.4
1	C	136	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	165	ILE	2.4
1	B	370	LEU	2.4
1	C	15	LEU	2.4
1	D	77	HIS	2.4
1	D	14	THR	2.4
1	A	187	VAL	2.4
1	A	83	THR	2.4
1	C	170	ARG	2.4
1	D	155	ASN	2.4
1	D	257	LEU	2.4
1	C	374	GLY	2.4
1	D	28	ARG	2.4
1	C	46	TRP	2.4
1	A	31	ASP	2.3
1	B	163	ASP	2.3
1	C	129	MET	2.3
1	C	77	HIS	2.3
1	A	113	ALA	2.3
1	C	83	THR	2.3
1	C	132	GLU	2.3
1	D	159	ARG	2.3
1	C	169	LEU	2.3
1	B	373	ARG	2.3
1	C	29	PHE	2.3
1	D	281	PHE	2.3
1	A	287	GLY	2.3
1	D	186	VAL	2.2
1	D	164	ALA	2.2
1	C	81	HIS	2.2
1	C	161	ALA	2.2
1	D	162	LYS	2.2
1	C	134	VAL	2.2
1	B	244	LYS	2.2
1	C	371	SER	2.2
1	B	136[A]	ARG	2.2
1	D	382	LEU	2.2
1	B	379	ASP	2.2
1	C	82	LYS	2.2
1	C	117	ALA	2.2
1	C	164	ALA	2.2
1	A	28	ARG	2.1
1	A	158	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	229	GLY	2.1
1	B	153	PRO	2.1
1	C	128	TYR	2.1
1	C	47	ALA	2.1
1	D	165	ILE	2.1
1	A	256	GLY	2.0
1	A	343	ALA	2.0
1	A	286	GLU	2.0
1	C	286	GLU	2.0
1	B	108	GLY	2.0
1	A	267	ASN	2.0
1	A	82	LYS	2.0
1	C	293	HIS	2.0
1	A	280	TYR	2.0
1	C	141	VAL	2.0
1	B	382	LEU	2.0
1	B	300	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FEJ	B	401	25/25	0.95	0.16	31,35,50,51	0
3	NA	A	402	1/1	0.96	0.25	45,45,45,45	0
2	FEJ	C	401	25/25	0.97	0.16	30,34,45,46	0
2	FEJ	A	401	25/25	0.97	0.16	31,34,42,42	0
3	NA	D	402	1/1	0.97	0.18	35,35,35,35	0
3	NA	B	402	1/1	0.98	0.20	34,34,34,34	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NA	C	402	1/1	0.98	0.07	39,39,39,39	0
2	FEJ	D	401	25/25	0.98	0.13	33,35,39,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FEJ B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
 and green (positive)






Electron density around FEJ C 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FEJ A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.