Full wwPDB EM Validation Report (i)

Mar 4, 2024 – 07:31 PM EST

PDB ID	:	6CVB
EMDB ID	:	EMD-7638
Title	:	CryoEM structure of human enterovirus D68 in complex with 6'-sialyl-N-acet yllactosamine
Authors	:	Liu, Y.; Rossmann, M.G.
Deposited on	:	2018-03-27
Resolution	:	2.43 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev70
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	6% 87% 11% •
2	B	247	• 84% 15% •
3	C	248	• 75% 20% •
4	D	68	7% 40% • 56%
5	E	3	33% 67% 33%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6660 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called viral protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	290	2273	1448	392	424	9	0	0

- Molecule 2 is a protein called viral protein 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	247	1900	1201	313	368	18	0	0

- Molecule 3 is a protein called viral protein 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	237	1864	1188	326	337	13	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	116	ARG	LYS	conflict	UNP A0A097ZN88

- Molecule 4 is a protein called viral protein 4.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
4	D	30	240	153	36	51	0	0

- Molecule 5 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galacto pyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
5	E	3	46	25	2	19	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		AltConf
			Total	O	
6	A	128	128	128	0
			Total	O	
6	B	92	92	92	0
			Total	O	
6	C	101	101	101	0
			Total	O	
6	D	16	16	16	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: viral protein 1

Chain A:



- Molecule 2: viral protein 3

Chain B:



- Molecule 3: viral protein 2

Chain C:



- Molecule 4: viral protein 4

Chain D:



• Molecule 5: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	5938	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION; On-the-fly CTF correction during 2D alignment and 3D reconstruction	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	25	Depositor
Minimum defocus (nm)	400	Depositor
Maximum defocus (nm)	3600	Depositor
Magnification	29000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	250.841	Depositor
Minimum map value	-193.535	Depositor
Average map value	0.078	Depositor
Map value standard deviation	14.182	Depositor
Recommended contour level	35.5	Depositor
Map size (Å)	464.0, 464.0, 464.0	wwPDB
Map dimensions	464, 464, 464	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.0, 1.0, 1.0	Depositor

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, NAG, SIA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.49	0/2323	0.60	0/3155
2	B	0.52	0/1945	0.67	0/2649
3	C	0.53	0/1922	0.63	0/2630
4	D	0.47	0/246	0.55	0/332
All	All	0.51	0/6436	0.63	0/8766

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2273	0	2279	32	0
2	B	1900	0	1850	29	0
3	C	1864	0	1793	32	0
4	D	240	0	219	3	0
5	E	46	0	40	2	0
6	A	128	0	0	0	0
6	B	92	0	0	1	0
6	C	101	0	0	1	0
6	D	16	0	0	0	0
All	All	6660	0	6181	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:23:VAL:HG11	1:A:26:LEU:HG	1.21	1.11
1:A:27:ASN:HD22	1:A:35:SER:HA	1.26	1.01
1:A:23:VAL:HG21	1:A:26:LEU:HD11	1.40	1.00
2:B:180:ASN:HD21	5:E:1:NAG:H81	1.36	0.91
1:A:23:VAL:CG1	1:A:26:LEU:HG	2.06	0.82
1:A:26:LEU:O	1:A:26:LEU:HD12	1.85	0.76
1:A:27:ASN:ND2	1:A:35:SER:HA	2.00	0.75
3:C:109:HIS:HD2	3:C:111:GLN:HE21	1.36	0.72
1:A:27:ASN:HD22	1:A:35:SER:CA	2.01	0.70
1:A:23:VAL:HG11	1:A:26:LEU:CG	2.14	0.67
1:A:56:GLU:HA	1:A:61:ASN:HD22	1.60	0.66
1:A:23:VAL:HG21	1:A:26:LEU:CD1	2.20	0.65
2:B:23:PRO:HG3	4:D:37:ALA:HB2	1.79	0.65
3:C:71:TRP:CE2	3:C:217:LEU:HB2	2.34	0.62
3:C:199:ASP:OD2	3:C:204:HIS:ND1	2.31	0.62
3:C:167:LEU:HD21	3:C:214:VAL:HG22	1.80	0.62
2:B:24:VAL:HG13	2:B:25:LEU:HG	1.83	0.61
3:C:116:ARG:NH1	6:C:301:HOH:O	2.33	0.61
2:B:180:ASN:ND2	5:E:1:NAG:H81	2.11	0.60
1:A:30:GLU:HA	3:C:177:TRP:HB2	1.83	0.59
2:B:47:VAL:HG22	2:B:99:VAL:HA	1.84	0.59
2:B:122:SER:O	3:C:181:ARG:NH1	2.37	0.57
3:C:71:TRP:NE1	3:C:217:LEU:HB2	2.20	0.57
3:C:122:LEU:HB2	3:C:178:ILE:HB	1.87	0.57
3:C:102:TYR:HH	3:C:104:SER:HG	1.53	0.56
1:A:280:ILE:O	2:B:95:ARG:NH2	2.39	0.56
1:A:75:GLU:OE1	1:A:128:ASN:ND2	2.39	0.56
1:A:100:PHE:HA	2:B:235:GLN:HE22	1.72	0.55
2:B:108:HIS:HB2	2:B:225:ARG:HB2	1.88	0.54
1:A:64:SER:OG	4:D:42:LYS:NZ	2.40	0.54
2:B:113:LEU:HD13	2:B:216:ILE:HD11	1.90	0.53
2:B:202:VAL:HG21	2:B:208:ASP:HA	1.90	0.53
3:C:12:ARG:HH21	3:C:27:GLU:HG2	1.75	0.52
3:C:198:MET:SD	3:C:240:ASN:ND2	2.78	0.51
1:A:183:THR:HG1	2:B:21:SER:HG	1.57	0.50
1:A:184:ILE:HG12	2:B:24:VAL:HG11	1.93	0.50
1:A:183:THR:OG1	2:B:21:SER:OG	2.23	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:158:GLY:O	3:C:181:ARG:NH2	2.44	0.50
3:C:108:ILE:HG12	3:C:232:ILE:HG12	1.92	0.50
3:C:15:GLN:HG3	3:C:24:VAL:HG12	1.93	0.49
2:B:47:VAL:HG13	2:B:218:ALA:HB3	1.94	0.49
3:C:199:ASP:OD2	3:C:206:GLN:NE2	2.45	0.49
3:C:66:LEU:HD22	3:C:150:GLU:HG2	1.96	0.48
1:A:112:TYR:HB2	1:A:249:TRP:HB2	1.96	0.47
1:A:63:LEU:HB2	1:A:246:ILE:HD13	1.96	0.47
3:C:99:HIS:CG	3:C:239:PHE:HB3	2.49	0.47
2:B:114:GLU:OE1	2:B:116:THR:OG1	2.33	0.47
3:C:125:VAL:HG13	3:C:210:ALA:HB3	1.96	0.47
2:B:205:GLU:HB3	3:C:221:THR:HB	1.97	0.46
2:B:50:GLU:HB2	2:B:215:PHE:HB3	1.97	0.46
3:C:109:HIS:CD2	3:C:111:GLN:HE21	2.25	0.46
2:B:74:LEU:HD12	2:B:196:MET:HG3	1.99	0.45
1:A:70:SER:HB3	1:A:239:VAL:HB	1.98	0.45
2:B:128:LYS:HE2	2:B:156:ASP:HB2	1.99	0.45
3:C:71:TRP:HE3	3:C:112:CYS:SG	2.39	0.45
1:A:95:ILE:HD11	1:A:220:LEU:HD13	1.99	0.44
2:B:199:ASN:ND2	6:B:405:HOH:O	2.45	0.44
1:A:25:SER:O	2:B:164:THR:N	2.39	0.44
1:A:8:ALA:HB1	2:B:221:ASP:HA	2.00	0.44
1:A:148:VAL:HB	1:A:219:ASN:HB2	2.00	0.44
3:C:190:LEU:HD13	3:C:207:TRP:CG	2.52	0.44
3:C:37:GLU:HB2	3:C:191:PRO:HG3	2.00	0.43
2:B:111:GLY:HA3	2:B:222:PHE:HA	2.00	0.43
3:C:71:TRP:CE3	3:C:112:CYS:SG	3.11	0.43
3:C:135:HIS:CE1	3:C:136:ASN:HB2	2.54	0.43
1:A:253:PRO:HB3	3:C:169:CYS:HB2	2.00	0.42
3:C:80:TRP:HZ3	3:C:211:ILE:HD12	1.84	0.42
1:A:26:LEU:HA	2:B:163:VAL:HA	2.00	0.42
1:A:77:LYS:H	1:A:89:ASN:HD21	1.68	0.42
3:C:161:LEU:HD11	3:C:170:ALA:HB2	2.01	0.42
3:C:105:GLY:N	3:C:235:MET:O	2.49	0.42
2:B:129:LEU:HG	2:B:157:PHE:HE2	1.85	0.41
1:A:127:VAL:HG22	1:A:233:PHE:HE1	1.85	0.41
1:A:105:ARG:HD2	2:B:233:ILE:HB	2.02	0.41
3:C:212:ILE:HA	3:C:213:PRO:HD2	1.90	0.41
1:A:27:ASN:ND2	1:A:35:SER:CA	2.73	0.41
2:B:144:ARG:O	2:B:148:MET:HG3	2.21	0.41
3:C:10:SER:OG	4:D:58:GLU:OE2	2.29	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:63:LEU:HD22	1:A:107:LEU:HD23	2.01	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/297 (96%)	272 (96%)	12 (4%)	0	100	100
2	B	245/247 (99%)	236 (96%)	9 (4%)	0	100	100
3	C	235/248 (95%)	223 (95%)	11 (5%)	1 (0%)	34	41
4	D	28/68 (41%)	25 (89%)	3 (11%)	0	100	100
All	All	792/860 (92%)	756 (96%)	35 (4%)	1 (0%)	54	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	30	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/257 (98%)	251 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	216/216 (100%)	212 (98%)	4 (2%)	57	69
3	C	200/208 (96%)	199 (100%)	1 (0%)	88	93
4	D	27/55 (49%)	27 (100%)	0	100	100
All	All	694/736 (94%)	689 (99%)	5 (1%)	84	90

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	41	ARG
2	B	47	VAL
2	B	148	MET
2	B	226	LEU
3	C	88	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	ASN
1	A	89	ASN
1	A	102	GLN
1	A	229	GLN
2	B	79	GLN
2	B	96	ASN
2	B	180	ASN
2	B	235	GLN
2	B	240	HIS
3	C	15	GLN
3	C	88	ASN
3	C	111	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	NAG	E	1	5	15,15,15	0.62	0	21,21,21	0.93	2 (9%)
5	GAL	E	2	5	11,11,12	0.46	0	15,15,17	0.81	1 (6%)
5	SIA	E	3	5	20,20,21	1.12	1 (5%)	24,28,31	1.24	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	E	1	5	-	4/6/26/26	0/1/1/1
5	GAL	E	2	5	-	1/2/19/22	0/1/1/1
5	SIA	E	3	5	-	0/18/34/38	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	3	SIA	C2-C1	4.10	1.56	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	E	3	SIA	O6-C2-C1	3.19	113.95	107.70
5	E	3	SIA	C6-O6-C2	2.42	116.51	111.34
5	E	3	SIA	C8-C7-C6	-2.30	108.68	113.03
5	E	2	GAL	C1-C2-C3	-2.25	106.91	109.67
5	E	1	NAG	C3-C4-C5	-2.08	106.52	110.24
5	E	1	NAG	C1-C2-N2	-2.02	108.39	110.73

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	E	1	NAG	O5-C5-C6-O6
5	E	1	NAG	C8-C7-N2-C2
5	E	1	NAG	O7-C7-N2-C2
5	E	2	GAL	O5-C5-C6-O6
5	E	1	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	1	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain E



5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-7638. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



X



Y



Z

6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map


X Index: 232


Y Index: 232


Z Index: 232

6.2.2 Raw map


X Index: 232


Y Index: 232


Z Index: 232

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map


X Index: 289


Y Index: 147


Z Index: 283

6.3.2 Raw map


X Index: 227


Y Index: 321


Z Index: 181

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map


X


Y


Z

6.4.2 Raw map


X


Y


Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map

X Y Z

The images above show the 3D surface view of the map at the recommended contour level 35.5. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 1689 nm^3; this corresponds to an approximate mass of 1526 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.412 Å^{-1}

8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.412 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.43	-	-
Author-provided FSC curve	2.43	2.74	2.47
Unmasked-calculated*	2.55	2.87	2.58

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-7638 and PDB model 6CVB. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlays

9.1.1 Map-model overlay (i)



X



Y



Z

9.1.2 Map-model assembly overlay (i)



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 35.5 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (35.5).

9.4 Atom inclusion (i)



At the recommended contour level, 91% of all backbone atoms, 89% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (35.5) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8930	0.6780
A	0.8630	0.6680
B	0.9300	0.6890
C	0.9220	0.6850
D	0.7350	0.6400
E	0.5650	0.6240