

# Full wwPDB X-ray Structure Validation Report (i)

#### May 25, 2020 – 02:27 pm BST

PDB ID	:	6CVU
Title	:	Crystal structure of Mycobacterium tuberculosis dethiobiotin synthetase in
		complex with cytidine
Authors	:	Thompson, A.P.; Wegener, K.L.; Bruning, J.B.; Polyak, S.W.
Deposited on	:	2018-03-28
$\operatorname{Resolution}$	:	2.43  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity		4.02b-467 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)		1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
$\operatorname{Refmac}$	:	5.8.0158
$\operatorname{CCP4}$	:	$7.0.044 (\mathrm{Gargrove})$
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
R <sub>free</sub>	130704	4647(2.44-2.40)
Clashscore	141614	5161(2.44-2.40)
Ramachandran outliers	138981	5073(2.44-2.40)
Sidechain outliers	138945	5074(2.44-2.40)
RSRZ outliers	127900	4543 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	235	86%	10% ••
1	В	235	78%	16% • •
1	С	235	% • 85%	11% ••
1	D	235	86%	10% •



#### $6\mathrm{CVU}$

# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6709 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	227	Total	С	Ν	Ο	S	0	0	0
	A	221	1562	976	282	297	7	0		0
1	В	226	Total	С	Ν	Ο	S	0	2	0
	I D	220	1585	988	289	301	7	0	2	U
1	C	228	Total	С	Ν	Ο	S	0	0	0
		220	1574	983	285	299	7	0	0	0
1	1 D	007	Total	С	Ν	Ο	S	0	0	0
	227	1567	979	282	299	7	0	0	U	

• Molecule 1 is a protein called ATP-dependent dethiobiotin synthetase BioD.

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	MET	-	initiating methionine	UNP A5U2S7
A	-7	GLY	-	expression tag	UNP A5U2S7
A	-6	HIS	-	expression tag	UNP A5U2S7
A	-5	HIS	-	expression tag	UNP A5U2S7
A	-4	HIS	-	expression tag	UNP A5U2S7
A	-3	HIS	-	expression tag	UNP A5U2S7
A	-2	HIS	-	expression tag	UNP A5U2S7
A	-1	HIS	-	expression tag	UNP A5U2S7
A	0	GLY	-	expression tag	UNP A5U2S7
A	1	GLY	-	expression tag	UNP A5U2S7
В	-8	MET	-	initiating methionine	UNP A5U2S7
В	-7	GLY	-	expression tag	UNP A5U2S7
В	-6	HIS	-	expression tag	UNP A5U2S7
В	-5	HIS	-	expression tag	UNP A5U2S7
В	-4	HIS	-	expression tag	UNP A5U2S7
В	-3	HIS	-	expression tag	UNP A5U2S7
В	-2	HIS	-	expression tag	UNP A5U2S7
В	-1	HIS	-	expression tag	UNP A5U2S7
В	0	GLY	-	expression tag	UNP A5U2S7
В	1	GLY	-	expression tag	UNP A5U2S7
С	-8	MET	-	initiating methionine	UNP A5U2S7



Chain	Residue	Modelled	Actual Comment		Reference
С	-7	GLY	-	expression tag	UNP A5U2S7
С	-6	HIS	-	expression tag	UNP A5U2S7
С	-5	HIS	-	expression tag	UNP A5U2S7
С	-4	HIS	-	expression tag	UNP A5U2S7
С	-3	HIS	-	expression tag	UNP A5U2S7
С	-2	HIS	-	expression tag	UNP A5U2S7
С	-1	HIS	-	expression tag	UNP A5U2S7
С	0	GLY	-	expression tag	UNP A5U2S7
С	1	GLY	-	expression tag	UNP A5U2S7
D	-8	MET	-	initiating methionine	UNP A5U2S7
D	-7	GLY	-	expression tag	UNP A5U2S7
D	-6	HIS	_	expression tag	UNP A5U2S7
D	-5	HIS	_	expression tag	UNP A5U2S7
D	-4	HIS	-	expression tag	UNP A5U2S7
D	-3	HIS	-	expression tag	UNP A5U2S7
D	-2	HIS	-	expression tag	UNP A5U2S7
D	-1	HIS	-	expression tag	UNP A5U2S7
D	0	GLY	-	expression tag	UNP A5U2S7
D	1	GLY	-	expression tag	UNP A5U2S7

Continued from previous page...

• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ) (labeled as "Ligand of Interest" by author).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	А	1	Total 5	0 4	${ m S}$	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2	С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2	С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 3 is 4-AMINO-1-BETA-D-RIBOFURANOSYL-2(1H)-PYRIMIDINONE (three-letter code: CTN) (formula:  $C_9H_{13}N_3O_5$ ) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total C N O 17 9 3 5	0	0
3	В	1	Total         C         N         O           17         9         3         5	0	0
3	С	1	Total         C         N         O           17         9         3         5	0	0
3	D	1	Total C N O 17 9 3 5	0	0



• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	89	Total O 89 89	0	0
4	В	87	Total O 87 87	0	0
4	С	68	Total O 68 68	0	0
4	D	69	Total O 69 69	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATP-dependent dethiobiotin synthetase BioD







## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	54.67Å 108.12Å 153.38Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	38.44 - 2.43	Depositor
Resolution (A)	46.49 - 2.43	EDS
% Data completeness	99.6 (38.44-2.43)	Depositor
(in resolution range)	99.6 (46.49-2.43)	EDS
R <sub>merge</sub>	0.07	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.40$ (at $2.42\text{\AA}$ )	Xtriage
Refinement program	PHENIX	Depositor
D D	0.197 , $0.272$	Depositor
$R, R_{free}$	0.197 , $0.272$	DCC
$R_{free}$ test set	1718 reflections $(4.89\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	54.0	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33, $39.4$	EDS
L-test for twinning <sup>2</sup>	$ \langle L  \rangle = 0.50, \langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6709	wwPDB-VP
Average B, all atoms $(Å^2)$	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.76% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CTN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	nd lengths	Bond angles		
	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.43	1/1581~(0.1%)	0.61	0/2164	
1	В	0.41	0/1604	0.59	1/2193~(0.0%)	
1	С	0.35	0/1593	0.54	0/2179	
1	D	0.40	0/1586	0.56	0/2171	
All	All	0.40	1/6364~(0.0%)	0.57	1/8707~(0.0%)	

All (1) bond length outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	А	19	CYS	CB-SG	-5.55	1.72	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	204	ASP	CB-CG-OD1	5.20	122.98	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1562	0	1608	15	0
1	В	1585	0	1636	34	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	С	1574	0	1621	18	0
1	D	1567	0	1611	12	0
2	А	10	0	0	0	0
2	В	10	0	0	0	0
2	С	10	0	0	0	0
2	D	10	0	0	1	0
3	А	17	0	13	0	0
3	В	17	0	13	0	0
3	С	17	0	13	0	0
3	D	17	0	13	0	0
4	А	89	0	0	5	0
4	В	87	0	0	15	0
4	С	68	0	0	5	0
4	D	69	0	0	4	0
All	All	6709	0	6528	77	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All $(77)$ close contacts	within the	e same	a symmetric	$\operatorname{unit}$	$\operatorname{are}$	listed	below,	sorted $\$	by	$\operatorname{their}$	$\operatorname{clash}$
magnitude.											

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:55[A]:ARG:NH2	4:B:402:HOH:O	1.93	1.01
1:B:177:LEU:O	4:B:401:HOH:O	1.85	0.94
1:B:48:ASP:OD1	4:B:403:HOH:O	1.96	0.83
1:C:163:CYS:SG	4:C:463:HOH:O	2.35	0.83
1:D:163:CYS:SG	4:D:467:HOH:O	2.38	0.79
1:B:163:CYS:SG	4:B:480:HOH:O	2.43	0.77
1:D:48:ASP:OD1	4:D:401:HOH:O	2.03	0.76
1:A:163:CYS:SG	4:A:484:HOH:O	2.45	0.74
1:B:19:CYS:HB3	4:B:476:HOH:O	1.86	0.74
1:B:181:SER:N	4:B:401:HOH:O	2.22	0.73
2:D:302:SO4:O4	4:D:402:HOH:O	2.06	0.73
1:A:87:ALA:HB3	1:A:90:GLN:HG3	1.72	0.72
1:B:40:GLN:HB2	4:B:417:HOH:O	1.92	0.69
1:B:124:LEU:HG	4:B:429:HOH:O	1.96	0.64
1:A:19:CYS:SG	4:A:481:HOH:O	2.54	0.64
1:C:55:ARG:NH1	4:C:405:HOH:O	2.30	0.63
1:B:116:GLU:HG3	1:B:121:GLY:HA2	1.82	0.61
1:B:93:ARG:NH2	4:B:410:HOH:O	2.33	0.59
1:B:43:THR:HA	1:B:47:ASP:HB2	1.84	0.59



Continued from previo	Atom-2	Interatomic	Clash
Atom-1	Atom-2	$distance ( m \AA)$	overlap (Å)
1:B:91:ILE:HD12	1:B:118:ALA:HB2	1.84	0.58
1:C:207:ASP:O	4:C:401:HOH:O	2.17	0.58
1:C:88:ARG:NH1	1:C:130:ASP:OD2	2.38	0.57
1:D:9:THR:HB	1:D:148:HIS:HB3	1.88	0.56
1:A:88:ARG:HG3	1:A:122:VAL:HG11	1.88	0.54
1:B:189:ILE:O	4:B:405:HOH:O	2.19	0.53
1:B:93:ARG:NH2	4:B:412:HOH:O	2.41	0.52
1:D:113:LEU:HD11	1:D:125:ARG:HB2	1.90	0.52
1:C:43:THR:OG1	1:C:69:PRO:HA	2.10	0.52
1:A:97:ASP:HB2	4:A:476:HOH:O	2.09	0.51
1:C:55:ARG:NH2	4:C:407:HOH:O	2.32	0.51
1:A:100:ARG:HG2	1:A:103:ARG:HD2	1.93	0.49
1:C:34:ALA:HB1	1:C:98:LEU:HD13	1.94	0.49
1:B:184[B]:SER:OG	1:B:188:ARG:NH1	2.45	0.49
1:B:120:PRO:O	1:C:90:GLN:NE2	2.46	0.49
1:C:86:PRO:HG2	1:C:91:ILE:HD11	1.95	0.49
1:C:172:PRO:HG2	1:C:175:PRO:HA	1.94	0.49
4:A:465:HOH:O	1:B:150:LYS:HE3	2.13	0.48
1:A:88:ARG:HD3	1:A:130:ASP:OD2	2.13	0.48
1:C:35:VAL:HG21	1:C:53:VAL:HG11	1.96	0.47
1:C:88:ARG:NH2	1:C:126:ASP:HB3	2.30	0.47
1:B:122:VAL:HG13	1:B:126:ASP:HB2	1.97	0.46
1:B:188:ARG:NH1	4:B:421:HOH:O	2.49	0.46
1:B:147:ASN:HB3	4:B:432:HOH:O	2.17	0.45
1:A:177:LEU:HG	1:B:72:MET:HE2	1.98	0.45
1:B:109:GLY:HA3	1:B:124:LEU:HD22	1.99	0.45
1:C:118:ALA:HB3	1:C:122:VAL:HB	1.99	0.45
1:B:217:ASP:O	1:B:221:VAL:HG23	2.17	0.45
1:C:35:VAL:HB	1:C:62:LEU:HD23	2.00	0.44
1:C:3:ILE:HD12	1:C:131:VAL:HB	1.98	0.44
1:C:65:LEU:HB2	1:C:94:LEU:HD22	1.99	0.44
1:A:63:ALA:CB	1:A:98:LEU:HD11	2.46	0.44
1:A:44:ALA:HB2	1:A:69:PRO:HB3	1.99	0.44
1:B:88:ARG:HG3	1:B:122:VAL:HG11	1.99	0.44
1:D:78:ALA:HB1	1:D:83:MET:O	2.18	0.44
1:D:88:ARG:HH21	1:D:130:ASP:CG	2.21	0.44
1:A:9:THR:HB	1:A:148:HIS:HB3	1.98	0.43
1:B:5:VAL:HG21	4:B:429:HOH:O	2.18	0.43
1:B:47:ASP:HB3	1:B:48:ASP:H	1.66	0.43
1:D:15:LYS:HB3	4:D:415:HOH:O	2.18	0.43
1:D:186:LEU:HA	1:D:189:ILE:HD12	2.01	0.43

Continued from previous page..



Atom-1	Atom-2	Interatomic	Clash
Atom-1	At0111-2	distance (Å)	overlap (Å)
1:B:5:VAL:HG22	1:B:135:ALA:HA	2.01	0.42
1:C:88:ARG:HG3	1:C:122:VAL:HG11	2.02	0.42
1:A:63:ALA:HB3	1:A:98:LEU:HD11	2.01	0.42
1:A:147:ASN:ND2	1:B:112:GLY:HA3	2.34	0.42
1:B:52:GLU:HG3	1:B:55[B]:ARG:HH21	1.85	0.42
1:B:40:GLN:HG3	1:B:41:THR:N	2.35	0.42
1:C:201:ALA:HB1	4:C:406:HOH:O	2.19	0.42
1:A:100:ARG:HD2	1:A:103:ARG:CZ	2.50	0.41
1:D:119:GLU:HG3	1:D:119:GLU:H	1.64	0.41
1:D:5:VAL:HG12	1:D:107:VAL:HB	2.03	0.41
1:B:112:GLY:O	1:B:115:VAL:HG22	2.21	0.41
1:D:173:ASP:HA	1:D:174:PRO:HA	1.80	0.41
1:D:23:ALA:HB1	1:D:33:VAL:HG11	2.03	0.41
1:A:15:LYS:NZ	4:A:406:HOH:O	2.44	0.41
1:B:180:ALA:HB3	4:B:401:HOH:O	2.20	0.41
1:B:35:VAL:HG21	1:B:53:VAL:HG11	2.02	0.41
1:B:88:ARG:NH2	1:B:89:ASP:OD1	2.54	0.40

Continued from previous page...

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	225/235~(96%)	220~(98%)	4 (2%)	1 (0%)	34 47
1	В	226/235~(96%)	217~(96%)	7(3%)	2(1%)	17 24
1	С	226/235~(96%)	216~(96%)	9~(4%)	1 (0%)	34 47
1	D	225/235~(96%)	219~(97%)	6 (3%)	0	100 100
All	All	902/940~(96%)	872 (97%)	26~(3%)	4 (0%)	34 47

All (4) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	В	48	ASP
1	А	120	PRO
1	С	65	LEU
1	В	46	GLY

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	hain Analysed Rotameric Outliers		Percentiles		
1	А	149/158~(94%)	143~(96%)	6~(4%)	31	48
1	В	153/158~(97%)	148~(97%)	5(3%)	38	56
1	С	150/158~(95%)	145~(97%)	5(3%)	38	56
1	D	150/158~(95%)	144~(96%)	6~(4%)	31	48
All	All	602/632~(95%)	580~(96%)	22~(4%)	34	51

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	61	GLN
1	А	88	ARG
1	А	119	GLU
1	А	181	SER
1	А	191	MET
1	А	218	ARG
1	В	4	LEU
1	В	45	ARG
1	В	88	ARG
1	В	177	LEU
1	В	189	ILE
1	С	60	THR
1	С	98	LEU
1	С	100	ARG
1	С	162	SER
1	С	204	ASP
1	D	4	LEU
1	D	11	THR



Continued from previous page...

Mol	Chain	Res	Type
1	D	40	GLN
1	D	55	ARG
1	D	119	GLU
1	D	143	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trees	Chain	Res	Link	Bo	ond leng	ths	B	ond ang	les
	Type	Chain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	CTN	С	303	-	15, 18, 18	1.32	2 (13%)	17,26,26	1.17	1(5%)
3	CTN	А	303	-	15,18,18	1.56	4 (26%)	17,26,26	1.55	<mark>3 (17%)</mark>
2	SO4	А	302	-	4,4,4	0.29	0	6,6,6	0.71	0
2	SO4	С	301	-	4,4,4	0.16	0	6,6,6	0.34	0
2	SO4	D	301	-	4,4,4	0.13	0	6,6,6	0.14	0
2	SO4	А	301	-	4,4,4	0.17	0	6,6,6	0.23	0



Mol	ol Type Chain Res		Link	Bo	ond leng	ths	B	ond ang	les	
	туре	Cham	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	#  Z  > 2
2	SO4	В	301	-	4,4,4	0.19	0	6,6,6	0.36	0
2	SO4	С	302	-	$4,\!4,\!4$	0.15	0	6,6,6	0.30	0
2	SO4	В	302	-	$4,\!4,\!4$	0.23	0	6,6,6	0.38	0
2	SO4	D	302	-	$4,\!4,\!4$	0.08	0	6,6,6	0.17	0
3	CTN	В	303	-	15, 18, 18	1.39	3 (20%)	17,26,26	1.16	2 (11%)
3	CTN	D	303	-	15, 18, 18	1.35	2 (13%)	17,26,26	1.20	1(5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	$\mathbf{Link}$	Chirals	Torsions	Rings
3	CTN	В	303	-	-	0/4/22/22	0/2/2/2
3	CTN	С	303	-	-	0/4/22/22	0/2/2/2
3	CTN	А	303	-	-	2/4/22/22	0/2/2/2
3	$\operatorname{CTN}$	D	303	-	-	2/4/22/22	0/2/2/2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
3	С	303	CTN	C4-N4	3.76	1.46	1.35
3	В	303	CTN	C4-N4	3.71	1.46	1.35
3	D	303	CTN	C4-N4	3.59	1.45	1.35
3	А	303	CTN	C4-N4	3.59	1.45	1.35
3	А	303	CTN	C6-N1	-2.84	1.32	1.35
3	А	303	CTN	C5-C4	-2.47	1.35	1.41
3	D	303	CTN	C4-N3	-2.20	1.32	1.35
3	В	303	CTN	C6-N1	-2.20	1.33	1.35
3	С	303	CTN	C6-N1	-2.14	1.33	1.35
3	А	303	$\operatorname{CTN}$	C4-N3	-2.13	1.32	1.35
3	В	303	CTN	C4-N3	-2.10	1.32	1.35

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	А	303	CTN	C2-N3-C4	4.49	120.89	116.34
3	С	303	CTN	C2-N3-C4	3.52	119.91	116.34
3	D	303	CTN	C2-N3-C4	2.97	119.35	116.34
3	А	303	CTN	N4-C4-N3	2.94	121.14	116.49
3	В	303	CTN	C2-N3-C4	2.92	119.30	116.34



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	А	303	CTN	C5-C4-N3	-2.27	119.10	121.72
3	В	303	CTN	C5-C4-N3	-2.11	119.29	121.72

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	303	CTN	C3'-C4'-C5'-O5'
3	D	303	CTN	O4'-C4'-C5'-O5'
3	А	303	CTN	C3'-C4'-C5'-O5'
3	А	303	CTN	O4'-C4'-C5'-O5'


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	302	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















































## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# RSRZ {>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	$Q{<}0.9$
1	А	227/235~(96%)	-0.35	0 100 100	35,  48,  68,  89	0
1	В	226/235~(96%)	-0.11	0 100 100	37, 52, 80, 121	0
1	С	228/235~(97%)	0.03	3 (1%) 77 75	42, 63, 92, 109	0
1	D	227/235~(96%)	-0.30	0 100 100	37, 54, 76, 115	1 (0%)
All	All	908/940~(96%)	-0.18	3 (0%) 94 93	35, 54, 83, 121	1 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	226	GLY	2.3
1	С	30	GLY	2.2
1	С	61	GLN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.


## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
3	CTN	С	303	17/17	0.86	0.26	$66,\!73,\!78,\!78$	17
3	CTN	D	303	17/17	0.90	0.15	$51,\!61,\!71,\!74$	0
3	CTN	А	303	17/17	0.91	0.15	$49,\!59,\!69,\!74$	17
2	SO4	А	302	5/5	0.92	0.24	$38,\!44,\!48,\!48$	5
3	CTN	В	303	17/17	0.94	0.17	42,51,62,68	0
2	SO4	С	302	5/5	0.97	0.29	$47,\!48,\!50,\!61$	5
2	SO4	С	301	5/5	0.98	0.13	$52,\!59,\!60,\!61$	0
2	SO4	В	302	5/5	0.99	0.17	$37,\!38,\!42,\!47$	5
2	SO4	D	302	5/5	0.99	0.15	$46,\!48,\!51,\!53$	5
2	SO4	А	301	5/5	0.99	0.14	$41,\!41,\!46,\!49$	0
2	SO4	D	301	5/5	0.99	0.14	43,47,52,55	0
2	SO4	В	301	5/5	1.00	0.15	40,43,44,48	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
























































## 6.5 Other polymers (i)

There are no such residues in this entry.

