



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 04:04 am BST

PDB ID : 5D1Z  
Title : IsdB NEAT1 bound by clone D4-10  
Authors : Deng, X.  
Deposited on : 2015-08-04  
Resolution : 3.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1467 (3.20-3.16)
Clashscore	141614	1599 (3.20-3.16)
Ramachandran outliers	138981	1574 (3.20-3.16)
Sidechain outliers	138945	1573 (3.20-3.16)
RSRZ outliers	127900	1423 (3.20-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	 92% 7%
1	H	214	 93% 7%
2	B	264	 75% 7% 17%
2	G	264	 73% 10% 16%
3	C	214	 81% 14%
3	E	214	 88% 11%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	D	270	 <p>3% 72% 13% 15%</p>
4	F	270	 <p>% 67% 13% 17%</p>
5	I	157	 <p>2% 91% 5%</p>
5	J	157	 <p>% 86% 8% 5%</p>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 15704 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Y10 Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	213	Total 1644	C 1033	N 275	O 331	S 5	0	0	0
1	H	214	Total 1650	C 1036	N 276	O 332	S 6	0	0	0

- Molecule 2 is a protein called Y10 Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	218	Total 1645	C 1041	N 279	O 319	S 6	0	0	0
2	G	222	Total 1668	C 1053	N 283	O 325	S 7	0	0	0

- Molecule 3 is a protein called D4-10 Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	208	Total 1601	C 1004	N 268	O 325	S 4	0	0	0
3	E	213	Total 1635	C 1025	N 274	O 332	S 4	0	0	0

- Molecule 4 is a protein called D4-10 Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	230	Total 1729	C 1090	N 289	O 343	S 7	0	0	0
4	F	223	Total 1680	C 1062	N 280	O 331	S 7	0	0	0


- Molecule 5 is a protein called Iron-regulated surface determinant protein B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	I	149	Total	C	N	O	S	0	0	0
			1226	787	204	233	2			
5	J	149	Total	C	N	O	S	0	0	0
			1226	787	204	233	2			



There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	113	GLY	-	expression tag	UNP Q6GA86
I	114	PRO	-	expression tag	UNP Q6GA86
J	113	GLY	-	expression tag	UNP Q6GA86
J	114	PRO	-	expression tag	UNP Q6GA86








• Molecule 3: D4-10 Light Chain





• Molecule 4: D4-10 Heavy Chain




• Molecule 4: D4-10 Heavy Chain




• Molecule 5: Iron-regulated surface determinant protein B



• Molecule 5: Iron-regulated surface determinant protein B







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	115.29Å 147.00Å 165.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.96 – 3.17 24.96 – 3.17	Depositor EDS
% Data completeness (in resolution range)	98.0 (24.96-3.17) 98.2 (24.96-3.17)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.73 (at 3.17Å)	Xtrriage
Refinement program	REFMAC 5.8.0123	Depositor
R, $R_{free}$	0.233 , 0.287 0.233 , 0.285	Depositor DCC
$R_{free}$ test set	2344 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.7	Xtrriage
Anisotropy	0.101	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 6.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	15704	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.53 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.5101e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.77	0/1682	0.84	1/2288 (0.0%)
1	H	0.80	0/1688	0.82	1/2296 (0.0%)
2	B	0.82	0/1686	0.88	0/2294
2	G	0.79	0/1709	0.87	0/2325
3	C	0.79	0/1635	0.84	0/2223
3	E	0.79	0/1670	0.86	2/2271 (0.1%)
4	D	0.82	0/1772	0.91	1/2415 (0.0%)
4	F	0.80	0/1722	0.92	1/2348 (0.0%)
5	I	0.80	0/1257	0.85	1/1695 (0.1%)
5	J	0.78	0/1257	0.81	0/1695
All	All	0.80	0/16078	0.86	7/21850 (0.0%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	73	LEU	CA-CB-CG	6.67	130.65	115.30
3	E	127	LEU	CA-CB-CG	6.50	130.24	115.30
3	E	144	ARG	NE-CZ-NH1	5.96	123.28	120.30
5	I	149	ARG	NE-CZ-NH1	5.41	123.00	120.30
4	F	45	ARG	NE-CZ-NH1	5.21	122.90	120.30
1	A	144	ARG	NE-CZ-NH1	5.16	122.88	120.30
1	H	156	LEU	CA-CB-CG	5.06	126.94	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1644	0	1592	5	0
1	H	1650	0	1596	10	0
2	B	1645	0	1599	9	0
2	G	1668	0	1621	13	0
3	C	1601	0	1553	14	0
3	E	1635	0	1593	7	0
4	D	1729	0	1700	12	0
4	F	1680	0	1646	11	0
5	I	1226	0	1204	1	0
5	J	1226	0	1204	8	0
All	All	15704	0	15308	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:87:THR:HG22	1:H:105:ARG:HG3	1.82	0.61
4:F:58:THR:HG21	5:J:169:SER:HB2	1.83	0.60
4:D:170:ASN:O	4:D:212:ASN:ND2	2.36	0.59
4:F:99:LYS:HE3	4:F:117:VAL:HG22	1.85	0.59
1:A:165:VAL:HG22	1:A:177:LEU:HD12	1.84	0.59
1:A:147:LYS:HE2	1:A:149:GLN:OE1	2.03	0.58
4:D:35:TYR:HB3	4:D:54:TYR:CD1	2.39	0.58
4:F:11:LEU:CD2	4:F:131:THR:HB	2.34	0.57
1:A:93:LEU:HD13	2:B:108:TYR:CE1	2.40	0.57
3:C:9:PRO:HG2	3:C:12:LEU:HD13	1.87	0.57
4:D:99:LYS:HE3	4:D:117:VAL:HG12	1.88	0.55
2:G:89:ALA:O	2:G:92:THR:HG23	2.07	0.55
2:B:31:ASN:O	2:B:54:VAL:HG13	2.06	0.55
4:D:226:VAL:O	4:D:226:VAL:HG12	2.07	0.55
4:D:142:SER:OG	4:D:147:SER:OG	2.19	0.53
4:D:136:VAL:HG21	4:D:213:VAL:HG11	1.91	0.51
5:J:185:ILE:HD13	5:J:256:PHE:CZ	2.46	0.51
2:G:12:VAL:HG11	2:G:155:PHE:HE1	1.76	0.50
2:B:170:SER:HB2	4:F:104:ASP:HB3	1.94	0.50
4:F:58:THR:HG23	5:J:191:SER:OG	2.11	0.50
4:D:146:THR:HG22	4:D:146:THR:O	2.11	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:65:VAL:HG13	2:B:69:PHE:HB2	1.93	0.49
3:C:211:ASN:N	3:C:211:ASN:OD1	2.46	0.49
3:E:13:LEU:CD2	3:E:21:VAL:HG13	2.43	0.49
1:A:40:GLN:HE22	2:B:40:GLN:HE22	1.59	0.48
2:G:223:LYS:NZ	1:H:124:ASP:OD2	2.47	0.48
2:B:33:PHE:O	2:B:73:ARG:NH2	2.48	0.47
4:D:141:PRO:HG3	4:D:153:LEU:CB	2.45	0.47
3:E:13:LEU:HD21	3:E:21:VAL:HG13	1.96	0.46
2:B:3:VAL:HG13	2:B:28:PHE:CD1	2.51	0.46
1:H:87:THR:HG22	1:H:105:ARG:CG	2.45	0.46
5:J:200:VAL:HG22	5:J:236:ILE:HG12	1.98	0.46
1:A:94:ASN:HD22	1:A:95:THR:HG23	1.81	0.46
2:B:35:MET:CB	2:B:80:LEU:HD13	2.46	0.46
4:D:18:LEU:CD1	4:D:124:VAL:HG11	2.45	0.46
2:B:89:ALA:O	2:B:92:THR:HG23	2.16	0.45
3:C:124:GLU:N	3:C:124:GLU:OE2	2.50	0.45
2:G:33:PHE:O	2:G:73:ARG:NH2	2.50	0.45
5:J:182:LYS:HG2	5:J:183:PRO:HD2	1.99	0.45
1:H:93:LEU:HA	1:H:98:TYR:CD1	2.52	0.45
1:H:94:ASN:OD1	5:J:139:LYS:HE3	2.17	0.45
3:E:160:ASN:HB2	3:E:181:LEU:HD12	1.98	0.44
4:F:99:LYS:NZ	4:F:100:PRO:O	2.49	0.44
3:E:144:ARG:NH2	3:E:165:VAL:HG13	2.33	0.44
2:G:23:CYS:HB3	2:G:80:LEU:HB3	1.99	0.44
2:G:65:VAL:HG13	2:G:69:PHE:HB2	1.99	0.44
3:E:137:LEU:HD21	4:F:196:VAL:HG21	2.00	0.43
3:C:159:ASN:HD22	3:C:159:ASN:N	2.16	0.43
4:F:29:ILE:O	4:F:31:SER:N	2.52	0.43
2:G:12:VAL:HG11	2:G:155:PHE:CE1	2.53	0.43
4:F:226:VAL:O	4:F:226:VAL:HG23	2.18	0.43
4:F:168:SER:O	4:F:212:ASN:N	2.51	0.43
2:G:107:GLY:HA3	1:H:93:LEU:CD2	2.49	0.43
2:G:40:GLN:HE22	1:H:40:GLN:HE22	1.66	0.42
3:C:181:THR:O	3:C:182:LEU:HD12	2.19	0.42
3:C:151:VAL:HG22	3:C:190:HIS:CD2	2.54	0.42
3:C:92:LEU:HA	3:C:97:PHE:CD1	2.54	0.42
3:C:32:SER:OG	3:C:68:PRO:HB3	2.20	0.42
2:G:107:GLY:HA3	1:H:93:LEU:HD22	2.02	0.42
3:E:66:GLY:HA2	3:E:74:THR:O	2.19	0.42
1:H:151:LYS:HB2	1:H:195:ALA:HB3	2.01	0.42
2:G:106:SER:HB2	5:J:137:ALA:HB2	2.01	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:18:LEU:HD12	4:D:124:VAL:HG11	2.00	0.42
3:C:192:VAL:HA	3:C:211:ASN:HB3	2.01	0.41
4:D:168:SER:O	4:D:212:ASN:HB2	2.20	0.41
3:C:35:ALA:HA	3:C:49:ILE:O	2.20	0.41
4:D:13:LYS:HB2	4:D:16:GLU:HG3	2.02	0.41
3:C:114:PRO:HB3	3:C:140:PHE:HB3	2.03	0.41
5:I:200:VAL:HG23	5:I:209:ILE:HD11	2.03	0.41
3:C:12:LEU:HD21	3:C:20:VAL:HG13	2.03	0.41
4:F:58:THR:HG21	5:J:169:SER:CB	2.49	0.41
3:C:7:GLN:HE21	3:C:89:CYS:HB2	1.85	0.40
3:C:194:ALA:HA	3:C:209:SER:HB3	2.02	0.40
3:E:93:LEU:HA	3:E:98:PHE:CD1	2.56	0.40
2:G:108:TYR:CE1	1:H:93:LEU:HD13	2.56	0.40
2:G:73:ARG:NH1	2:G:75:ASN:OD1	2.53	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/214 (99%)	199 (94%)	12 (6%)	0	100	100
1	H	212/214 (99%)	199 (94%)	12 (6%)	1 (0%)	29	66
2	B	214/264 (81%)	202 (94%)	10 (5%)	2 (1%)	17	54
2	G	218/264 (83%)	205 (94%)	10 (5%)	3 (1%)	11	44
3	C	204/214 (95%)	186 (91%)	15 (7%)	3 (2%)	10	43
3	E	211/214 (99%)	196 (93%)	13 (6%)	2 (1%)	17	54
4	D	228/270 (84%)	198 (87%)	24 (10%)	6 (3%)	5	29
4	F	219/270 (81%)	190 (87%)	18 (8%)	11 (5%)	2	14
5	I	147/157 (94%)	134 (91%)	13 (9%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	J	147/157 (94%)	139 (95%)	8 (5%)	0	100	100
All	All	2011/2238 (90%)	1848 (92%)	135 (7%)	28 (1%)	11	44

All (28) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	144	THR
4	F	30	SER
4	F	130	SER
2	B	54	VAL
4	D	142	SER
4	F	129	ALA
4	F	150	THR
4	F	219	ASN
2	G	125	THR
2	G	224	SER
1	H	145	GLU
4	D	143	SER
4	F	29	ILE
4	F	151	ALA
3	C	155	LEU
4	D	66	GLN
4	D	144	LYS
3	E	129	SER
4	F	15	SER
4	D	30	SER
3	E	213	ARG
4	F	75	ARG
4	F	26	GLY
4	F	227	GLU
2	G	211	PRO
3	C	84	VAL
3	C	69	GLY
4	D	162	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/187 (100%)	177 (95%)	9 (5%)	25	60
1	H	187/187 (100%)	182 (97%)	5 (3%)	44	74
2	B	182/220 (83%)	176 (97%)	6 (3%)	38	70
2	G	185/220 (84%)	175 (95%)	10 (5%)	22	55
3	C	183/188 (97%)	172 (94%)	11 (6%)	19	51
3	E	187/188 (100%)	176 (94%)	11 (6%)	19	52
4	D	199/232 (86%)	188 (94%)	11 (6%)	21	55
4	F	192/232 (83%)	171 (89%)	21 (11%)	6	24
5	I	135/140 (96%)	132 (98%)	3 (2%)	52	78
5	J	135/140 (96%)	130 (96%)	5 (4%)	34	67
All	All	1771/1934 (92%)	1679 (95%)	92 (5%)	23	56

All (92) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	THR
1	A	58	SER
1	A	65	SER
1	A	72	GLU
1	A	128	LYS
1	A	147	LYS
1	A	161	SER
1	A	169	ASP
1	A	204	SER
2	B	64	SER
2	B	78	ASN
2	B	92	THR
2	B	103	ASP
2	B	122	SER
2	B	218	LYS
3	C	34	LEU
3	C	54	THR
3	C	61	SER
3	C	82	GLU
3	C	110	THR
3	C	124	GLU
3	C	133	VAL
3	C	157	SER

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
3	C	169	SER
3	C	190	HIS
3	C	211	ASN
4	D	7	SER
4	D	87	VAL
4	D	125	THR
4	D	135	SER
4	D	171	SER
4	D	194	SER
4	D	201	SER
4	D	208	THR
4	D	220	THR
4	D	223	ASP
4	D	225	LYS
5	I	120	ASN
5	I	158	LYS
5	I	263	SER
3	E	31	VAL
3	E	52	SER
3	E	65	SER
3	E	95	THR
3	E	111	THR
3	E	116	SER
3	E	131	THR
3	E	133	SER
3	E	157	GLN
3	E	166	THR
3	E	199	THR
4	F	7	SER
4	F	19	SER
4	F	23	THR
4	F	30	SER
4	F	60	TYR
4	F	76	SER
4	F	78	ASN
4	F	85	SER
4	F	97	CYS
4	F	104	ASP
4	F	105	ARG
4	F	126	VAL
4	F	130	SER
4	F	168	SER

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
4	F	171	SER
4	F	184	VAL
4	F	188	SER
4	F	198	THR
4	F	203	SER
4	F	218	SER
4	F	220	THR
2	G	29	THR
2	G	64	SER
2	G	92	THR
2	G	126	LYS
2	G	178	VAL
2	G	182	SER
2	G	187	LEU
2	G	197	SER
2	G	212	SER
2	G	214	THR
1	H	9	SER
1	H	32	THR
1	H	62	PRO
1	H	133	SER
1	H	176	SER
5	J	121	THR
5	J	161	THR
5	J	182	LYS
5	J	227	VAL
5	J	252	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	94	ASN
2	B	40	GLN
2	B	78	ASN
2	B	208	ASN
3	C	148	GLN
3	C	159	ASN
3	C	161	GLN
3	C	190	HIS
5	I	262	ASN
3	E	149	GLN
4	F	207	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	G	53	GLN
1	H	40	GLN
1	H	162	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	213/214 (99%)	0.00	1 (0%) 91 86	34, 53, 66, 72	0
1	H	214/214 (100%)	-0.01	2 (0%) 84 75	35, 51, 67, 75	0
2	B	218/264 (82%)	-0.12	2 (0%) 84 75	30, 43, 62, 89	0
2	G	222/264 (84%)	-0.05	7 (3%) 47 31	32, 42, 60, 84	2 (0%)
3	C	208/214 (97%)	-0.10	2 (0%) 82 72	26, 41, 93, 108	0
3	E	213/214 (99%)	0.07	5 (2%) 60 46	28, 49, 92, 102	0
4	D	230/270 (85%)	0.16	8 (3%) 44 28	27, 50, 87, 96	6 (2%)
4	F	223/270 (82%)	0.10	2 (0%) 84 75	30, 55, 83, 95	0
5	I	149/157 (94%)	-0.23	3 (2%) 65 50	26, 38, 61, 78	0
5	J	149/157 (94%)	-0.16	1 (0%) 87 81	29, 40, 66, 80	0
All	All	2039/2238 (91%)	-0.02	33 (1%) 72 59	26, 47, 82, 108	8 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	216	CYS	5.1
2	G	1	ALA	4.8
2	G	225	CYS	3.9
4	D	143	SER	3.6
2	G	142	GLY	3.3
2	G	224	SER	3.2
3	C	123	ASP	3.0
4	F	205	GLY	2.9
4	D	145	SER	2.8
2	G	210	LYS	2.8
4	D	230	SER	2.7
1	H	215	GLU	2.6
3	E	129	SER	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	G	141	SER	2.5
5	I	119	ASN	2.5
2	B	213	ASN	2.4
4	D	146	THR	2.3
2	B	222	PRO	2.3
5	I	267	PHE	2.3
4	D	28	SER	2.3
4	D	202	SER	2.3
5	I	180	ASP	2.2
4	D	105	ARG	2.2
4	F	148	GLY	2.2
3	E	59	GLY	2.2
2	G	200	THR	2.2
4	D	206	THR	2.2
1	A	214	GLY	2.1
5	J	267	PHE	2.1
3	E	204	SER	2.1
3	C	203	SER	2.1
3	E	215	GLU	2.1
3	E	153	ASP	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.