



# Full wwPDB X-ray Structure Validation Report i

Dec 12, 2023 – 07:30 pm GMT

PDB ID : 4D2B  
Title : Structure of a ligand free POT family peptide transporter  
Authors : Lyons, J.A.; Parker, J.L.; Solcan, N.; Brinth, A.; Li, D.; Shah, S.T.A.; Caffrey, M.; Newstead, S.  
Deposited on : 2014-05-09  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	491	7%	83%	8%	9%

## 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 3697 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DI-OR TRIPEPTIDE\H+ SYMPORTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	445	3440	2319	526	578	17	0	1	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	484	GLY	-	expression tag	UNP Q5M4H8
A	485	SER	-	expression tag	UNP Q5M4H8
A	486	GLU	-	expression tag	UNP Q5M4H8
A	487	ASN	-	expression tag	UNP Q5M4H8
A	488	LEU	-	expression tag	UNP Q5M4H8
A	489	TYR	-	expression tag	UNP Q5M4H8
A	490	PHE	-	expression tag	UNP Q5M4H8
A	491	GLN	-	expression tag	UNP Q5M4H8

- Molecule 2 is (2R)-2,3-DIHYDROXYPROPYL(7Z)-PENTADEC-7-ENOATE (three-letter code: 78N) (formula: C<sub>18</sub>H<sub>34</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 22 18 4	0	0
2	A	1	Total C O 22 18 4	0	0
2	A	1	Total C O 22 18 4	0	0
2	A	1	Total C O 22 18 4	0	0
2	A	1	Total C O 22 18 4	0	0

- Molecule 3 is (2S)-2,3-DIHYDROXYPROPYL(7Z)-PENTADEC-7-ENOATE (three-letter code: 78M) (formula: C<sub>18</sub>H<sub>34</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 22 18 4	0	0
3	A	1	Total C O 22 18 4	0	0
3	A	1	Total C O 22 18 4	0	0
3	A	1	Total C O 22 18 4	0	0

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total    O    P 5    4    1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	54	Total    O 54    54	0	0

### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DI-OR TRIPEPTIDE\H+ SYMPORTER



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.19Å    110.19Å    110.99Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	39.10 – 2.35 39.10 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.2 (39.10-2.35) 99.2 (39.10-2.35)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.91 (at 2.34Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
$R$ , $R_{free}$	0.213 , 0.241 0.213 , 0.241	Depositor DCC
$R_{free}$ test set	1331 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.9	Xtriage
Anisotropy	0.170	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 50.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3697	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 78N, 78M, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/3542	0.40	0/4833

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3440	0	3515	17	0
2	A	110	0	170	1	0
3	A	88	0	136	3	0
4	A	5	0	0	0	0
5	A	54	0	0	0	0
All	All	3697	0	3821	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:GLU:OE1	1:A:126:LYS:NZ	2.25	0.69
1:A:22:GLU:HG2	1:A:129:VAL:HG11	1.86	0.57
1:A:406:VAL:O	1:A:410:VAL:HG23	2.16	0.46
1:A:97:LEU:HD23	1:A:100:ILE:HD12	1.98	0.46
1:A:49:LEU:HG	1:A:51:ILE:HG12	1.98	0.45
1:A:288:LEU:HD13	1:A:410:VAL:HG11	1.98	0.45
1:A:74:GLY:HA3	1:A:123:GLY:O	2.16	0.44
1:A:286:ILE:O	1:A:290:ILE:HG12	2.18	0.44
1:A:425:SER:HA	1:A:428:PHE:HD2	1.83	0.44
1:A:394:ALA:HB1	3:A:1478:78M:H153	2.00	0.43
3:A:1475:78M:H31C	3:A:1475:78M:H18	1.99	0.43
1:A:162:ALA:HB3	1:A:163:PRO:HD3	2.01	0.43
1:A:260:ILE:HB	1:A:261:PRO:HD3	2.01	0.42
1:A:423:MET:O	1:A:426:MET:HB2	2.19	0.42
2:A:1473:78N:H122	2:A:1479:78N:H52C	2.02	0.42
1:A:59:ILE:HD11	1:A:246:LEU:HD21	2.02	0.42
3:A:1474:78M:H8	3:A:1474:78M:H112	1.85	0.41
1:A:258:ILE:O	1:A:261:PRO:HD2	2.21	0.41
1:A:284:SER:O	1:A:287:PRO:HD2	2.21	0.40
1:A:356:LYS:CD	1:A:406:VAL:HG21	2.51	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	438/491 (89%)	434 (99%)	4 (1%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	359/398 (90%)	351 (98%)	8 (2%)	52 63

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	TYR
1	A	135	THR
1	A	141	ASP
1	A	244	ASN
1	A	344	THR
1	A	353	SER
1	A	393	TRP
1	A	404	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	78N	A	1477	-	21,21,21	0.89	1 (4%)	22,22,22	1.08	2 (9%)
3	78M	A	1476	-	21,21,21	0.89	1 (4%)	22,22,22	1.22	2 (9%)
2	78N	A	1473	-	21,21,21	0.90	1 (4%)	22,22,22	1.11	2 (9%)
2	78N	A	1480	-	21,21,21	0.90	1 (4%)	22,22,22	1.16	2 (9%)
4	PO4	A	1482	-	4,4,4	0.98	0	6,6,6	0.53	0
3	78M	A	1474	-	21,21,21	0.90	1 (4%)	22,22,22	1.18	2 (9%)
3	78M	A	1475	-	21,21,21	0.93	1 (4%)	22,22,22	1.22	2 (9%)
3	78M	A	1478	-	21,21,21	0.89	1 (4%)	22,22,22	1.08	2 (9%)
2	78N	A	1481	-	21,21,21	0.91	1 (4%)	22,22,22	1.08	2 (9%)
2	78N	A	1479	-	21,21,21	0.89	1 (4%)	22,22,22	1.18	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	78N	A	1477	-	-	7/21/21/21	-
3	78M	A	1476	-	-	13/21/21/21	-
2	78N	A	1473	-	-	12/21/21/21	-
2	78N	A	1480	-	-	12/21/21/21	-
3	78M	A	1474	-	-	12/21/21/21	-
3	78M	A	1475	-	-	10/21/21/21	-
3	78M	A	1478	-	-	12/21/21/21	-
2	78N	A	1481	-	-	11/21/21/21	-
2	78N	A	1479	-	-	10/21/21/21	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1481	78N	O16-C17	-2.74	1.38	1.45

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1473	78N	O16-C17	-2.73	1.38	1.45
2	A	1480	78N	O16-C17	-2.71	1.39	1.45
2	A	1477	78N	O16-C17	-2.69	1.39	1.45
3	A	1475	78M	O2-C17	-2.67	1.39	1.45
3	A	1476	78M	O2-C17	-2.61	1.39	1.45
3	A	1474	78M	O2-C17	-2.59	1.39	1.45
3	A	1478	78M	O2-C17	-2.59	1.39	1.45
2	A	1479	78N	O16-C17	-2.53	1.39	1.45

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1475	78M	O2-C1-C2	3.05	121.48	111.91
3	A	1476	78M	O2-C1-C2	2.95	121.16	111.91
2	A	1473	78N	O16-C8-C7	2.82	120.75	111.91
3	A	1474	78M	O2-C1-C2	2.81	120.72	111.91
2	A	1480	78N	O16-C8-C7	2.73	120.47	111.91
3	A	1476	78M	O2-C17-C18	2.70	118.83	105.77
2	A	1479	78N	O16-C17-C18	2.70	118.82	105.77
2	A	1479	78N	O16-C8-C7	2.70	120.37	111.91
3	A	1474	78M	O2-C17-C18	2.67	118.68	105.77
2	A	1481	78N	O16-C8-C7	2.62	120.13	111.91
2	A	1480	78N	O16-C17-C18	2.60	118.34	105.77
2	A	1473	78N	O16-C17-C18	2.59	118.29	105.77
2	A	1477	78N	O16-C17-C18	2.54	118.05	105.77
3	A	1478	78M	O2-C17-C18	2.54	118.01	105.77
3	A	1478	78M	O2-C1-C2	2.47	119.66	111.91
2	A	1477	78N	O16-C8-C7	2.34	119.25	111.91
2	A	1481	78N	O16-C17-C18	2.33	117.03	105.77
3	A	1475	78M	O2-C17-C18	2.20	116.39	105.77

There are no chirality outliers.

All (99) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1473	78N	O16-C17-C18-C19
2	A	1479	78N	O16-C17-C18-C19
2	A	1479	78N	O16-C17-C18-O20
2	A	1480	78N	O16-C17-C18-C19
3	A	1474	78M	C17-C18-C20-O21
3	A	1474	78M	C11-C10-C9-C8
3	A	1475	78M	C17-C18-C20-O21

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	A	1475	78M	C2-C1-O2-C17
3	A	1476	78M	C17-C18-C20-O21
3	A	1475	78M	O1-C1-O2-C17
2	A	1480	78N	C7-C8-O16-C17
2	A	1480	78N	O16-C17-C18-O20
2	A	1473	78N	C11-C10-C9-C1
2	A	1480	78N	O15-C8-O16-C17
3	A	1478	78M	O2-C17-C18-C20
2	A	1473	78N	C4-C5-C6-C7
2	A	1473	78N	O16-C17-C18-O20
2	A	1479	78N	C5-C6-C7-C8
3	A	1478	78M	O2-C17-C18-O19
3	A	1475	78M	C1-C2-C3-C4
2	A	1479	78N	C9-C10-C11-C12
2	A	1480	78N	C10-C11-C12-C13
2	A	1481	78N	C9-C10-C11-C12
2	A	1481	78N	C11-C12-C13-C14
3	A	1478	78M	C9-C10-C11-C12
2	A	1477	78N	C9-C10-C11-C12
3	A	1475	78M	C3-C4-C5-C6
2	A	1477	78N	C17-C18-C19-O22
2	A	1480	78N	C17-C18-C19-O22
2	A	1481	78N	C17-C18-C19-O22
3	A	1474	78M	C4-C5-C6-C7
3	A	1476	78M	C10-C11-C12-C13
3	A	1478	78M	C3-C4-C5-C6
3	A	1478	78M	C10-C11-C12-C13
2	A	1480	78N	C9-C10-C11-C12
3	A	1476	78M	C11-C12-C13-C14
2	A	1479	78N	C3-C4-C5-C6
3	A	1476	78M	C3-C4-C5-C6
3	A	1474	78M	O19-C18-C20-O21
3	A	1476	78M	O19-C18-C20-O21
2	A	1481	78N	C10-C11-C12-C13
3	A	1476	78M	C2-C3-C4-C5
2	A	1479	78N	C11-C10-C9-C1
2	A	1473	78N	C10-C11-C12-C13
3	A	1476	78M	C9-C10-C11-C12
3	A	1474	78M	C3-C4-C5-C6
3	A	1474	78M	C2-C3-C4-C5
3	A	1478	78M	C11-C10-C9-C8
3	A	1476	78M	C1-C2-C3-C4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	A	1478	78M	C4-C5-C6-C7
3	A	1478	78M	C5-C6-C7-C8
3	A	1475	78M	O19-C18-C20-O21
2	A	1473	78N	C5-C6-C7-C8
2	A	1481	78N	O16-C17-C18-C19
3	A	1478	78M	C7-C8-C9-C10
2	A	1481	78N	O16-C17-C18-O20
2	A	1477	78N	C11-C12-C13-C14
2	A	1481	78N	O20-C18-C19-O22
2	A	1477	78N	C3-C4-C5-C6
2	A	1473	78N	C2-C3-C4-C5
2	A	1477	78N	C10-C11-C12-C13
3	A	1476	78M	C2-C1-O2-C17
3	A	1476	78M	C12-C13-C14-C15
2	A	1473	78N	O20-C18-C19-O22
3	A	1476	78M	O1-C1-O2-C17
3	A	1475	78M	C9-C10-C11-C12
3	A	1474	78M	C11-C12-C13-C14
3	A	1478	78M	C1-C2-C3-C4
2	A	1477	78N	C6-C7-C8-O16
2	A	1480	78N	C3-C4-C5-C6
3	A	1474	78M	O2-C17-C18-O19
2	A	1479	78N	C1-C2-C3-C4
2	A	1479	78N	C11-C12-C13-C14
2	A	1479	78N	C4-C5-C6-C7
2	A	1481	78N	C1-C2-C3-C4
2	A	1473	78N	C11-C12-C13-C14
3	A	1476	78M	C11-C10-C9-C8
2	A	1481	78N	C2-C1-C9-C10
3	A	1478	78M	C12-C13-C14-C15
3	A	1474	78M	C10-C11-C12-C13
2	A	1481	78N	O15-C8-O16-C17
2	A	1473	78N	C17-C18-C19-O22
2	A	1473	78N	C1-C2-C3-C4
3	A	1475	78M	C5-C6-C7-C8
3	A	1476	78M	C5-C6-C7-C8
2	A	1481	78N	C7-C8-O16-C17
2	A	1477	78N	O20-C18-C19-O22
3	A	1475	78M	C7-C8-C9-C10
2	A	1473	78N	C2-C1-C9-C10
2	A	1479	78N	C2-C1-C9-C10
2	A	1480	78N	C6-C7-C8-O16

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	1480	78N	C18-C17-O16-C8
3	A	1474	78M	C7-C8-C9-C10
3	A	1474	78M	C9-C10-C11-C12
3	A	1474	78M	C1-C2-C3-C4
2	A	1480	78N	C6-C7-C8-O15
3	A	1475	78M	C2-C3-C4-C5
3	A	1478	78M	C2-C3-C4-C5
2	A	1480	78N	C12-C13-C14-C15


There are no ring outliers.


5 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1473	78N	1	0
3	A	1474	78M	1	0
3	A	1475	78M	1	0
3	A	1478	78M	1	0
2	A	1479	78N	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	445/491 (90%)	0.35	36 (8%) 12   17	19, 32, 72, 90	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	411	THR	8.1
1	A	423	MET	7.0
1	A	349	ASN	5.4
1	A	268	MET	5.1
1	A	140	HIS	5.1
1	A	284	SER	4.8
1	A	426	MET	4.8
1	A	267	TRP	4.5
1	A	269	ILE	4.4
1	A	412	THR	4.3
1	A	142	ARG	4.3
1	A	341	TRP	4.2
1	A	408	LEU	4.1
1	A	281	ARG	3.7
1	A	138	ASP	3.5
1	A	139	GLU	3.3
1	A	424	MET	3.2
1	A	409	SER	3.2
1	A	145	ASP	3.1
1	A	350	GLN	3.0
1	A	285	TYR	2.9
1	A	340	ALA	2.7
1	A	50	HIS	2.6
1	A	137	TYR	2.6
1	A	344	THR	2.5
1	A	287	PRO	2.5
1	A	338	PHE	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	410	VAL	2.5
1	A	265	PHE	2.4
1	A	315	ASP	2.3
1	A	270	SER	2.3
1	A	282	VAL	2.2
1	A	407	GLY	2.1
1	A	141	ASP	2.1
1	A	264	TYR	2.0
1	A	283	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	78M	A	1475	22/22	0.61	0.25	31,66,85,88	0
3	78M	A	1474	22/22	0.78	0.33	49,60,67,80	0
3	78M	A	1478	22/22	0.82	0.20	36,53,61,67	0
2	78N	A	1480	22/22	0.84	0.25	41,52,62,70	0
2	78N	A	1481	22/22	0.85	0.17	34,48,59,62	0
2	78N	A	1477	22/22	0.87	0.29	30,49,71,74	0
2	78N	A	1479	22/22	0.88	0.23	45,56,62,67	0
3	78M	A	1476	22/22	0.88	0.25	34,47,55,64	0
2	78N	A	1473	22/22	0.88	0.23	41,53,64,67	0
4	PO4	A	1482	5/5	0.98	0.16	47,51,62,63	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.











## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.