



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 20, 2023 – 05:21 PM JST

PDB ID : 7D2P  
Title : Crystal structure of MazE-MazF (Form-II) from *Deinococcus radiodurans*  
Authors : Dhanasingh, I.; Lee, S.H.  
Deposited on : 2020-09-17  
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	117	 2% (poor fit), 85% (0-1 outliers), 9% (2 outliers), 5% (3+ outliers)
1	B	117	 3% (poor fit), 83% (0-1 outliers), 12% (2 outliers), 5% (3+ outliers)
1	C	117	 5% (poor fit), 77% (0-1 outliers), 12% (2 outliers), 11% (3+ outliers)
1	D	117	 3% (poor fit), 82% (0-1 outliers), 5% (2 outliers), 12% (3+ outliers)
2	E	80	 35% (0-1 outliers), 62% (2+ outliers)
2	F	80	 34% (0-1 outliers), 62% (2+ outliers)

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3902 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endoribonuclease MazF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	111	Total 826	C 520	N 152	O 150	S 4	0	0	0
1	B	111	Total 826	C 520	N 152	O 150	S 4	0	0	0
1	C	104	Total 769	C 484	N 142	O 139	S 4	0	0	0
1	D	103	Total 764	C 481	N 141	O 138	S 4	0	0	0

- Molecule 2 is a protein called AbrB/MazE/SpoVT family DNA-binding domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	30	Total 240	C 151	N 38	O 50	S 1	0	0	0
2	F	30	Total 240	C 151	N 38	O 50	S 1	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	52	Total 52	O 52	0	0
3	B	41	Total 41	O 41	0	0
3	C	43	Total 43	O 43	0	0
3	D	49	Total 49	O 49	0	0
3	E	29	Total 29	O 29	0	0
3	F	23	Total 23	O 23	0	0





MET	THR	SER	GLN	ILE	GLN	LYS	TRP	GLY	ASN	SER	LEU	ALA	LEU	ARG	ILE	PRO	LYS	ALA	LEU	ALA	GLN	GLN	VAL	GLY	LEU	THR	GLN	SER	SER	GLU	VAL	GLU	LEU	LEU	LEU	LEU	GLN	ASP	GLY	GLN	ILE	VAL	ILE	ARG	PRO	VAL	PRO	PRO	ALA	ARG	GLN	Y51	D52	W80
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	39.82Å 42.33Å 65.00Å 96.51° 97.82° 86.71°	Depositor
Resolution (Å)	29.40 – 2.07 29.38 – 2.07	Depositor EDS
% Data completeness (in resolution range)	93.8 (29.40-2.07) 93.9 (29.38-2.07)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.46 (at 2.08Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.168 , 0.231 0.175 , 0.235	Depositor DCC
$R_{free}$ test set	1962 reflections (8.26%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.3	Xtrriage
Anisotropy	0.072	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 49.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3902	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.69	0/846	0.82	0/1153
1	B	0.70	0/846	0.88	0/1153
1	C	0.71	0/785	0.80	0/1068
1	D	0.74	0/780	0.84	0/1061
2	E	0.69	0/246	0.87	0/335
2	F	0.70	0/246	0.83	0/335
All	All	0.71	0/3749	0.84	0/5105

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	69	GLY	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	826	0	816	7	0
1	B	826	0	816	9	0
1	C	769	0	765	8	0
1	D	764	0	763	8	0
2	E	240	0	212	6	0
2	F	240	0	212	1	0
3	A	52	0	0	1	0
3	B	41	0	0	3	0
3	C	43	0	0	0	0
3	D	49	0	0	0	0
3	E	29	0	0	1	0
3	F	23	0	0	0	0
All	All	3902	0	3584	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:26:GLY:O	3:B:201:HOH:O	2.05	0.74
1:D:43:THR:HG22	2:E:60:MET:HE3	1.77	0.66
1:B:15:LEU:HD11	1:B:46:MET:CE	2.29	0.63
1:B:104:ARG:HD3	1:C:111:LEU:O	2.01	0.60
1:A:14:TRP:CZ3	1:A:28:ARG:NH2	2.74	0.55
1:A:24:GLU:OE2	3:A:201:HOH:O	2.18	0.54
1:C:14:TRP:HB3	1:C:28:ARG:HG2	1.88	0.54
1:C:55:ALA:HB2	1:C:73:VAL:HG22	1.90	0.54
1:A:58:TYR:CD1	1:A:59:PRO:HD2	2.43	0.53
1:D:43:THR:CG2	2:E:60:MET:HE3	2.37	0.53
1:A:15:LEU:HD11	1:A:46:MET:CE	2.39	0.52
1:A:13:VAL:O	1:A:30:PRO:HA	2.13	0.49
1:D:43:THR:CG2	2:E:60:MET:CE	2.92	0.48
1:D:43:THR:HG22	2:E:60:MET:CE	2.43	0.46
1:B:80:ARG:NH2	3:B:204:HOH:O	2.29	0.45
1:A:15:LEU:HD11	1:A:46:MET:HE1	1.98	0.45
1:D:57:GLY:H	1:D:61:GLU:CD	2.19	0.45
1:B:14:TRP:O	1:B:89:ALA:HA	2.19	0.43
1:B:97:ALA:HB1	3:B:206:HOH:O	2.18	0.43
1:D:68:LEU:HD23	1:D:68:LEU:HA	1.84	0.43
2:E:80:TRP:O	3:E:101:HOH:O	2.21	0.43
1:C:70:VAL:HB	1:C:92:LEU:HD21	2.01	0.42

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:57:GLY:N	1:D:61:GLU:OE1	2.46	0.42
1:A:16:ASN:OD1	1:A:28:ARG:HD3	2.19	0.42
1:C:55:ALA:HB1	1:C:73:VAL:CG1	2.49	0.42
1:B:19:PRO:HD3	1:B:26:GLY:CA	2.50	0.42
1:C:58:TYR:CD1	1:C:59:PRO:HD2	2.55	0.42
1:B:58:TYR:HA	1:B:59:PRO:HD3	1.96	0.41
2:E:60:MET:HE2	2:E:60:MET:HB3	1.92	0.41
1:C:50:PRO:HD3	2:F:80:TRP:CZ2	2.56	0.41
1:B:13:VAL:O	1:B:30:PRO:HA	2.22	0.40
1:D:14:TRP:CD1	1:D:28:ARG:NH1	2.90	0.40
1:C:63:THR:HA	1:C:73:VAL:HG12	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	109/117 (93%)	109 (100%)	0	0	100	100
1	B	109/117 (93%)	104 (95%)	4 (4%)	1 (1%)	17	8
1	C	100/117 (86%)	98 (98%)	2 (2%)	0	100	100
1	D	99/117 (85%)	97 (98%)	1 (1%)	1 (1%)	15	6
2	E	28/80 (35%)	28 (100%)	0	0	100	100
2	F	28/80 (35%)	27 (96%)	0	1 (4%)	3	0
All	All	473/628 (75%)	463 (98%)	7 (2%)	3 (1%)	25	15

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	27	GLY

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
2	F	52	ASP
1	B	98	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	84/90 (93%)	83 (99%)	1 (1%)	71	69
1	B	84/90 (93%)	84 (100%)	0	100	100
1	C	78/90 (87%)	75 (96%)	3 (4%)	33	26
1	D	78/90 (87%)	77 (99%)	1 (1%)	69	67
2	E	23/66 (35%)	22 (96%)	1 (4%)	29	22
2	F	23/66 (35%)	20 (87%)	3 (13%)	4	1
All	All	370/492 (75%)	361 (98%)	9 (2%)	49	43

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	TRP
1	C	29	ARG
1	C	56	LYS
1	C	98	ASP
1	D	68	LEU
2	E	80	TRP
2	F	51	TYR
2	F	52	ASP
2	F	80	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	111/117 (94%)	-0.21	2 (1%) 68 70	15, 22, 42, 58	0
1	B	111/117 (94%)	-0.01	4 (3%) 42 45	15, 24, 52, 67	0
1	C	104/117 (88%)	-0.06	6 (5%) 23 24	14, 25, 54, 75	0
1	D	103/117 (88%)	-0.02	3 (2%) 51 54	15, 24, 48, 73	0
2	E	30/80 (37%)	-0.26	0 100 100	21, 25, 35, 38	0
2	F	30/80 (37%)	-0.07	1 (3%) 46 49	18, 27, 38, 60	0
All	All	489/628 (77%)	-0.09	16 (3%) 46 49	14, 25, 51, 75	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	26	GLY	5.7
2	F	51	TYR	5.2
1	B	114	SER	4.8
1	A	114	SER	4.4
1	B	54	ARG	4.4
1	D	17	PHE	4.3
1	C	114	SER	3.9
1	D	27	GLY	3.7
1	C	56	LYS	3.5
1	C	55	ALA	3.4
1	A	54	ARG	3.3
1	C	18	THR	3.2
1	C	4	ASP	2.6
1	B	4	ASP	2.6
1	B	25	GLN	2.4
1	C	26	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.