Full wwPDB X-ray Structure Validation Report (i)

Jun 18, 2024 – 08:33 AM EDT

PDB ID	:	4D39
Title	:	Structure of bovine endothelial nitric oxide synthase heme domain in complex with2-(2-(1H-imidazol-1-yl)pyrimidin-4-yl)-N-(3-cyanobenzyl) ethan-1-amine
Authors	:	Li, H.; Poulos, T.L.
Deposited on	:	2014-10-20
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	4% 81% 9% • 9%
1	B	443	3% 80% 10% • 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	GOL	B	880	-	-	X	-

2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 7162 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, ENDOTHELIAL.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	As	C	N	O	S			
1	A	405	3231	1	2053	569	592	16	0	1	0
1	B	404	3221	1	2048	568	588	16	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	SEE REMARK 999	UNP P29473
B	100	ARG	CYS	SEE REMARK 999	UNP P29473

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	B	1	43	34	1	4	4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O			
3	A	1	17	9	5	3		0	0
3	B	1	17	9	5	3		0	0

- Molecule 4 is 3-[({2-[2-(1H-imidazol-1-yl)pyrimidin-4-yl]ethyl}amino)methyl]benzonitrile (three-letter code: OLW) (formula: $C_{17}H_{16}N_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	N		
4	A	1	23	17	6	0	0
4	B	1	23	17	6	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	4	2	2	0	0
5	B	1	4	2	2	0	0

- Molecule 6 is D-MANNITOL (three-letter code: MTL) (formula: $C_6H_{14}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	A	1	12	6	6	0	0
			Total	C	O		
6	B	1	12	6	6	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	A	1	6	3	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	B	1	6	3	3	0	0

- Molecule 8 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
8	A	1	1	1	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
9	A	287	287	287	0	0
			Total	O		
9	B	212	212	212	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL

Chain A:



• Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	57.76Å 106.12Å 156.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.08 – 2.00 39.08 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (39.08-2.00) 94.6 (39.08-1.90)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.29 (at 1.91Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.172 , 0.222 0.170 , 0.221	Depositor DCC
R_{free} test set	3586 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å^2)	36.9	Xtriage
Anisotropy	0.200	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 46.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	7162	wwPDB-VP
Average B, all atoms (Å^2)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, ZN, OLW, MTL, GOL, CAS, HEM, H4B

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.39	0/3311	0.53	0/4508
1	B	0.40	0/3300	0.53	0/4491
All	All	0.39	0/6611	0.53	0/8999

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3231	0	3130	30	0
1	B	3221	0	3130	29	0
2	A	43	0	30	3	0
2	B	43	0	30	2	0
3	A	17	0	15	2	0
3	B	17	0	15	1	0
4	A	23	0	16	2	0
4	B	23	0	16	5	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	A	12	0	14	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	12	0	14	0	0
7	A	6	0	8	2	0
7	B	6	0	8	4	0
8	A	1	0	0	0	0
9	A	287	0	0	3	0
9	B	212	0	0	2	0
All	All	7162	0	6432	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:281:TRP:HB2	1:B:304:LEU:HD21	1.77	0.67
4:B:800:OLW:H5'	7:B:880:GOL:H32	1.78	0.65
3:A:600:H4B:O4	7:A:880:GOL:O2	2.14	0.65
1:A:72:ARG:NH1	1:A:83:TYR:OH	2.31	0.64
1:A:106:VAL:HG12	1:A:107:LEU:HD12	1.82	0.62
1:B:154:GLU:OE1	1:B:168:ARG:NH2	2.33	0.60
1:B:141:LYS:O	1:B:142:ARG:NH1	2.34	0.60
2:B:500:HEM:HBB2	2:B:500:HEM:HHC	1.84	0.60
1:A:105:LEU:HD22	1:B:465:GLU:HB3	1.85	0.58
1:B:236:ARG:HD2	1:B:242:ASP:OD1	2.04	0.57
1:A:271:GLU:OE2	1:A:482:TRP:NE1	2.36	0.56
1:B:218:LYS:HG3	1:B:311:LEU:HD22	1.88	0.56
1:A:99:ARG:NH1	1:B:93:ASP:OD1	2.43	0.52
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.91	0.52
1:A:465:GLU:HB3	1:B:105:LEU:HD22	1.93	0.51
1:B:236:ARG:HD3	1:B:351:SER:HB3	1.93	0.51
1:B:378:LEU:HB2	9:B:2170:HOH:O	2.12	0.50
1:A:246:TRP:HB2	1:A:294:LEU:HB3	1.93	0.50
2:A:500:HEM:HBB2	2:A:500:HEM:HHC	1.94	0.49
1:B:175:GLY:HA3	1:B:345:ILE:HD13	1.94	0.49
1:A:93:ASP:OD1	1:B:99:ARG:NH1	2.43	0.49
1:A:100:ARG:NH1	1:A:102:LEU:HD22	2.28	0.48
1:A:367:ARG:HH22	7:A:880:GOL:H2	1.78	0.48
1:A:449:TRP:HA	3:A:600:H4B:N1	2.28	0.48
1:A:154:GLU:OE1	1:A:168:ARG:NH2	2.47	0.48
1:A:378:LEU:HB2	9:A:2224:HOH:O	2.13	0.47
1:B:167:LEU:HG	1:B:348:LEU:HD12	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:CYS:HB2	2:A:500:HEM:ND	2.28	0.47
1:B:121:PRO:HG2	1:B:123:GLU:OE1	2.13	0.47
1:A:252:ARG:HD3	1:A:269:ASN:HD21	1.80	0.47
1:A:72:ARG:HD2	9:A:2018:HOH:O	2.14	0.46
1:B:126:LEU:O	1:B:130:ARG:HG3	2.15	0.46
1:B:366:THR:O	1:B:370:CYS:HB2	2.15	0.46
1:A:340:ASN:HD21	6:A:870:MTL:H62	1.81	0.45
1:B:252:ARG:HD3	1:B:269:ASN:HD21	1.81	0.45
4:B:800:OLW:H5'	7:B:880:GOL:C3	2.44	0.45
1:A:455:SER:O	1:A:458:LEU:HB2	2.17	0.44
1:A:252:ARG:HD3	1:A:269:ASN:ND2	2.32	0.44
1:A:249:GLN:NE2	4:A:800:OLW:H15	2.33	0.44
1:B:252:ARG:HD3	1:B:269:ASN:ND2	2.33	0.44
1:A:70:PHE:CD1	1:A:85:THR:HG22	2.52	0.43
1:B:121:PRO:HB2	1:B:122:ALA:H	1.67	0.43
4:B:800:OLW:C5'	7:B:880:GOL:H32	2.47	0.43
1:A:152:LEU:O	1:A:156:GLU:HG3	2.19	0.43
1:B:449:TRP:HA	3:B:600:H4B:N1	2.33	0.43
1:B:180:TRP:CE3	1:B:192:TRP:HA	2.54	0.42
2:A:500:HEM:C1A	4:A:800:OLW:H02	2.54	0.42
1:A:244:ARG:NH2	1:A:481:PRO:HD3	2.34	0.42
9:A:2276:HOH:O	7:B:880:GOL:H2	2.20	0.42
1:B:249:GLN:NE2	4:B:800:OLW:H15	2.35	0.42
1:B:231:THR:O	1:B:354:PRO:HD2	2.19	0.42
1:A:472:SER:HA	1:A:473:PRO:C	2.40	0.42
1:B:319:HIS:CG	1:B:320:PRO:HD2	2.55	0.42
1:A:392:THR:HB	1:B:423:HIS:HB2	2.02	0.41
1:B:330:LEU:HB2	9:B:2151:HOH:O	2.18	0.41
1:A:126:LEU:HD12	1:A:126:LEU:HA	1.73	0.41
1:A:389:THR:HA	1:A:396:TRP:CD1	2.55	0.41
2:B:500:HEM:C1A	4:B:800:OLW:H02	2.56	0.41
1:A:240:ARG:HD3	1:A:298:PRO:CB	2.50	0.41
1:A:344:GLU:OE1	1:A:476:ARG:NH2	2.54	0.41
1:B:322:LEU:HD13	1:B:324:TRP:CZ2	2.56	0.40
1:B:240:ARG:HD2	1:B:241:GLY:O	2.21	0.40
1:B:370:CYS:SG	1:B:378:LEU:HD13	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	401/443 (90%)	392 (98%)	9 (2%)	0	100	100
1	B	399/443 (90%)	391 (98%)	8 (2%)	0	100	100
All	All	800/886 (90%)	783 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	344/375 (92%)	336 (98%)	8 (2%)	50	53
1	B	343/375 (92%)	332 (97%)	11 (3%)	39	38
All	All	687/750 (92%)	668 (97%)	19 (3%)	43	44

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	ARG
1	A	102	LEU
1	A	105	LEU
1	A	125	LEU
1	A	252	ARG
1	A	293	LEU
1	A	294	LEU
1	A	328	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	69	LYS
1	B	100	ARG
1	B	102	LEU
1	B	124	GLN
1	B	141	LYS
1	B	156	GLU
1	B	240	ARG
1	B	259	GLN
1	B	330	LEU
1	B	331	ARG
1	B	378	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	CAS	A	384	1	5,8,9	1.04	0	1,9,11	0.74	0
1	CAS	B	384	1	5,8,9	1.13	0	1,9,11	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	384	1	-	0/0/7/9	-
1	CAS	B	384	1	-	0/0/7/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	OLW	B	800	2	25,25,25	1.84	4 (16%)	29,32,32	2.39	12 (41%)
3	H4B	B	600	-	16,18,18	0.90	0	14,26,26	2.30	5 (35%)
6	MTL	B	870	-	11,11,11	0.46	0	14,14,14	0.63	0
5	ACT	A	860	-	3,3,3	0.77	0	3,3,3	0.84	0
2	HEM	B	500	1,4	42,50,50	1.84	6 (14%)	46,82,82	1.90	12 (26%)
6	MTL	A	870	-	11,11,11	0.42	0	14,14,14	0.72	0
2	HEM	A	500	1,4	42,50,50	1.89	5 (11%)	46,82,82	1.56	6 (13%)
3	H4B	A	600	-	16,18,18	0.90	0	14,26,26	2.35	5 (35%)
7	GOL	B	880	-	5,5,5	0.32	0	5,5,5	0.44	0
4	OLW	A	800	2	25,25,25	1.84	4 (16%)	29,32,32	2.21	7 (24%)
5	ACT	B	860	-	3,3,3	0.82	0	3,3,3	0.76	0
7	GOL	A	880	-	5,5,5	0.35	0	5,5,5	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OLW	B	800	2	-	0/9/13/13	0/3/3/3
3	H4B	B	600	-	-	0/8/17/17	0/2/2/2
6	MTL	B	870	-	-	2/16/16/16	-
2	HEM	B	500	1,4	-	1/12/54/54	-
6	MTL	A	870	-	-	3/16/16/16	-
2	HEM	A	500	1,4	-	1/12/54/54	-
3	H4B	A	600	-	-	0/8/17/17	0/2/2/2
7	GOL	B	880	-	-	0/4/4/4	-
4	OLW	A	800	2	-	0/9/13/13	0/3/3/3
7	GOL	A	880	-	-	2/4/4/4	-

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	HEM	C3D-C2D	7.71	1.53	1.36
2	B	500	HEM	C3D-C2D	7.54	1.53	1.36
4	B	800	OLW	C3'-C7'	-7.11	1.29	1.44
4	A	800	OLW	C3'-C7'	-6.92	1.29	1.44
2	A	500	HEM	C3C-C2C	-3.70	1.35	1.40
2	B	500	HEM	C3C-C2C	-3.69	1.35	1.40
2	B	500	HEM	C3C-CAC	3.63	1.55	1.47
2	A	500	HEM	C3C-CAC	3.44	1.55	1.47
4	B	800	OLW	C02-N03	-3.39	1.33	1.36
4	A	800	OLW	C04-N03	-3.28	1.34	1.39
4	B	800	OLW	C04-N03	-3.24	1.34	1.39
2	A	500	HEM	CAB-C3B	3.05	1.55	1.47
4	A	800	OLW	C12-N11	3.02	1.36	1.31
4	A	800	OLW	C02-N03	-2.89	1.33	1.36
2	B	500	HEM	CAB-C3B	2.87	1.55	1.47
2	B	500	HEM	C3C-C4C	2.77	1.45	1.41
2	A	500	HEM	C3C-C4C	2.60	1.45	1.41
4	B	800	OLW	C12-N11	2.51	1.35	1.31
2	B	500	HEM	CMB-C2B	2.06	1.55	1.50

All (47) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	A	800	OLW	C16-N11-C12	5.91	121.52	114.04

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	B	600	H4B	C8A-C4A-C4	5.69	119.68	114.50
4	B	800	OLW	C16-N11-C12	5.58	121.10	114.04
2	B	500	HEM	C4D-ND-C1D	5.24	111.42	105.21
3	A	600	H4B	C8A-C4A-C4	5.16	119.20	114.50
2	A	500	HEM	C4D-ND-C1D	5.01	111.14	105.21
2	B	500	HEM	CBA-CAA-C2A	-4.82	104.43	112.54
4	A	800	OLW	N11-C12-N13	-4.40	121.26	126.18
4	A	800	OLW	C15-C16-N11	-4.37	118.62	123.97
4	B	800	OLW	C15-C16-N11	-4.34	118.65	123.97
4	B	800	OLW	C17-C14-N13	4.05	122.20	116.06
4	B	800	OLW	C02-N03-C04	3.99	115.73	108.50
4	B	800	OLW	N11-C12-N13	-3.95	121.77	126.18
3	A	600	H4B	C2-N3-C4	3.72	121.14	115.96
3	A	600	H4B	N1-C2-N3	-3.68	119.84	125.48
4	A	800	OLW	C02-N03-C04	3.53	114.91	108.50
4	A	800	OLW	C17-C14-N13	3.46	121.30	116.06
2	B	500	HEM	CBD-CAD-C3D	-3.28	103.46	112.53
4	B	800	OLW	C16-C15-C14	3.25	119.97	116.68
4	A	800	OLW	C04-N03-C12	-3.23	121.86	125.50
3	B	600	H4B	N1-C2-N3	-3.21	120.56	125.48
2	A	500	HEM	C3B-C2B-C1B	3.12	108.76	106.41
2	B	500	HEM	C3B-C4B-NB	-3.11	107.23	109.47
4	A	800	OLW	C16-C15-C14	3.08	119.80	116.68
3	B	600	H4B	C2-N3-C4	3.04	120.19	115.96
2	B	500	HEM	C4B-CHC-C1C	2.87	126.34	122.56
2	B	500	HEM	C4C-CHD-C1D	2.86	126.33	122.56
4	B	800	OLW	C2'-C3'-C7'	2.85	123.40	119.55
2	B	500	HEM	C1B-NB-C4B	2.83	108.56	105.21
3	A	600	H4B	C2-N1-C8A	2.71	121.03	114.59
4	B	800	OLW	C04-N03-C12	-2.69	122.47	125.50
3	B	600	H4B	C2-N1-C8A	2.65	120.90	114.59
2	B	500	HEM	C3B-C2B-C1B	2.53	108.31	106.41
2	A	500	HEM	CBD-CAD-C3D	-2.51	105.59	112.53
2	B	500	HEM	CHD-C1D-ND	2.49	127.12	124.44
4	B	800	OLW	C4'-C3'-C7'	-2.43	116.01	119.97
2	A	500	HEM	CAD-C3D-C4D	2.36	128.81	124.70
3	B	600	H4B	N2-C2-N3	2.20	120.52	117.22
2	B	500	HEM	CMC-C2C-C3C	2.18	129.04	124.68
4	B	800	OLW	C02-N03-C12	-2.16	121.79	126.02
3	A	600	H4B	N2-C2-N3	2.13	120.42	117.22
4	B	800	OLW	C1'-C20-N19	-2.13	106.10	112.79
2	B	500	HEM	C2B-C1B-NB	-2.06	107.47	109.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	500	HEM	CHA-C4D-ND	2.05	126.92	124.37
4	B	800	OLW	C3'-C7'-N8'	-2.04	172.83	177.87
2	A	500	HEM	C4C-CHD-C1D	2.04	125.25	122.56
2	A	500	HEM	C4B-CHC-C1C	2.02	125.23	122.56

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	880	GOL	C1-C2-C3-O3
7	A	880	GOL	O2-C2-C3-O3
6	A	870	MTL	O5-C5-C6-O6
6	A	870	MTL	C4-C5-C6-O6
6	A	870	MTL	O1-C1-C2-O2
6	B	870	MTL	C4-C5-C6-O6
6	B	870	MTL	O5-C5-C6-O6
2	B	500	HEM	C4B-C3B-CAB-CBB
2	A	500	HEM	CAD-CBD-CGD-O2D

There are no ring outliers.

9 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	800	OLW	5	0
3	B	600	H4B	1	0
2	B	500	HEM	2	0
6	A	870	MTL	1	0
2	A	500	HEM	3	0
3	A	600	H4B	2	0
7	B	880	GOL	4	0
4	A	800	OLW	2	0
7	A	880	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand OLW B 800



Bond lengths



Bond angles



Torsions



Rings

Ligand HEM B 500



Bond lengths



Bond angles



Torsions



Rings

Ligand HEM A 500



Bond lengths



Bond angles



Torsions



Rings

Ligand OLW A 800



Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	404/443 (91%)	0.07	19 (4%)	31	30	26, 37, 67, 95	0
1	B	403/443 (90%)	-0.00	12 (2%)	50	49	26, 39, 67, 102	0
All	All	807/886 (91%)	0.03	31 (3%)	40	39	26, 38, 67, 102	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	122	ALA	5.1
1	B	259	GLN	4.9
1	A	160	ALA	4.4
1	A	259	GLN	4.3
1	B	121	PRO	4.3
1	B	110	LYS	4.1
1	A	109	ARG	4.0
1	A	121	PRO	4.0
1	A	239	GLY	3.9
1	A	126	LEU	3.9
1	A	67	GLY	3.8
1	A	338	VAL	3.3
1	A	124	GLN	3.0
1	A	123	GLU	3.0
1	A	69	LYS	2.9
1	A	127	SER	2.8
1	B	146	GLN	2.8
1	B	122	ALA	2.8
1	B	123	GLU	2.7
1	B	141	LYS	2.7
1	A	260	ASP	2.5
1	A	390	ARG	2.4
1	B	142	ARG	2.3
1	A	261	GLY	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	239	GLY	2.3
1	B	451	VAL	2.2
1	B	99	ARG	2.2
1	A	355	PHE	2.1
1	A	156	GLU	2.1
1	B	147	ALA	2.1
1	A	299	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	CAS	A	384	9/10	0.96	0.08	30,37,80,87	0
1	CAS	B	384	9/10	0.96	0.10	44,49,85,90	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	MTL	A	870	12/12	0.87	0.19	35,54,65,65	0
6	MTL	B	870	12/12	0.90	0.13	38,49,52,52	0
7	GOL	A	880	6/6	0.93	0.26	66,68,75,78	0
3	H4B	B	600	17/17	0.94	0.14	33,39,42,44	0
7	GOL	B	880	6/6	0.95	0.21	51,55,57,61	0
4	OLW	A	800	23/23	0.96	0.20	28,34,52,60	0
5	ACT	B	860	4/4	0.96	0.22	59,60,62,63	0
3	H4B	A	600	17/17	0.96	0.12	31,36,44,49	0
5	ACT	A	860	4/4	0.97	0.12	39,48,51,57	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	OLW	B	800	23/23	0.97	0.21	23,34,50,60	0
2	HEM	A	500	43/43	0.98	0.19	21,29,41,47	0
2	HEM	B	500	43/43	0.98	0.14	21,30,47,64	0
8	ZN	A	1483	1/1	1.00	0.03	39,39,39,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OLW A 800:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around OLW B 800:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 500:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 500:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.