



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 4, 2024 – 07:42 pm GMT

PDB ID : 5D6G  
Title : CRYSTAL STRUCTURE OF FRAGMENT OF RIBOSOMAL PROTEIN P0  
IN COMPLEX WITH 74NT 23S RNA FROM METHANOCOCCUS JAN-  
NASCHII  
Authors : Gabdulkhakov, A.G.; Mitroshin, I.V.; Garber, M.B.  
Deposited on : 2015-08-12  
Resolution : 3.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)
RNA backbone	3102	1117 (3.70-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	213	 4% 41% 48% 6%
2	0	74	 % 41% 54% 5%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3167 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 50S ribosomal protein L10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	205	1572	1014	266	286	6	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	MET	-	initiating methionine	UNP P54049

- Molecule 2 is a RNA chain called 23S ribosomal RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	0	74	1588	708	293	513	74	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
0	1224	C	-	expression tag	REF 470491724

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	0	2	Total	Mg	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	0	5	Total	O	0	0
			5	5		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 50S ribosomal protein L10



- Molecule 2: 23S ribosomal RNA



## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.23Å 108.31Å 112.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.83 – 3.30 29.64 – 3.20	Depositor EDS
% Data completeness (in resolution range)	98.7 (24.83-3.30) 98.0 (29.64-3.20)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.25 (at 3.18Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.245 , 0.298 0.251 , 0.305	Depositor DCC
$R_{free}$ test set	532 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	70.1	Xtrriage
Anisotropy	1.079	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.22 , 56.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.34$ , $\langle L^2 \rangle = 0.17$	Xtrriage
Estimated twinning fraction	0.098 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	3167	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	139.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	0/1595	0.87	3/2154 (0.1%)
2	0	0.59	1/1777 (0.1%)	1.12	1/2768 (0.0%)
All	All	0.57	1/3372 (0.0%)	1.02	4/4922 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	0	1151	G	OP3-P	-10.47	1.48	1.61

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	83	LEU	CA-CB-CG	5.95	128.99	115.30
1	A	177	LEU	CA-CB-CG	5.53	128.02	115.30
1	A	35	MET	CG-SD-CE	-5.47	91.45	100.20
2	0	1166	G	C8-N9-C4	-5.19	104.33	106.40

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	162	GLU	Peptide
1	A	175	PRO	Peptide
1	A	176	LYS	Peptide
1	A	179	ALA	Peptide
1	A	198	VAL	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1572	0	1705	126	0
2	0	1588	0	802	24	0
3	0	2	0	0	0	0
4	0	5	0	0	0	0
All	All	3167	0	2507	146	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

All (146) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:LYS:HG2	1:A:158:ILE:H	1.38	0.88
1:A:54:VAL:HB	1:A:94:LEU:HD11	1.56	0.85
1:A:13:LYS:NZ	2:0:1156:A:O2'	2.10	0.83
1:A:130:LYS:NZ	1:A:162:GLU:O	2.12	0.83
1:A:176:LYS:HD2	1:A:178:ALA:HB3	1.62	0.80
1:A:31:ALA:HA	1:A:198:VAL:HG22	1.62	0.79
1:A:100:PRO:HG3	1:A:199:TYR:HE1	1.46	0.79
1:A:32:ILE:HG22	1:A:196:LEU:HD12	1.64	0.78
1:A:141:LEU:HA	1:A:144:LEU:HD12	1.65	0.77
2:0:1178:G:O6	2:0:1179:A:N6	2.20	0.74
1:A:34:ASP:HB2	1:A:196:LEU:HD21	1.71	0.72
1:A:66:ARG:O	1:A:69:LYS:N	2.21	0.72
1:A:179:ALA:HB3	1:A:183:ARG:NH1	2.05	0.71
1:A:160:ILE:HD11	1:A:162:GLU:HB3	1.72	0.71
2:0:1163:G:O6	2:0:1216:C:N4	2.18	0.70

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:THR:HG23	1:A:98:MET:H	1.58	0.69
1:A:106:LEU:O	1:A:110:ASN:ND2	2.26	0.69
1:A:176:LYS:HG3	1:A:179:ALA:H	1.59	0.68
1:A:180:VAL:O	1:A:184:LEU:N	2.27	0.68
1:A:21:LYS:HD2	1:A:71:ALA:HB2	1.77	0.67
1:A:107:LEU:HD12	1:A:195:ILE:HD11	1.78	0.66
1:A:66:ARG:HG3	1:A:67:ALA:H	1.60	0.66
2:0:1176:U:O2'	2:0:1178:G:N7	2.24	0.65
2:0:1174:C:H2'	2:0:1175:U:O4'	1.98	0.64
1:A:157:LYS:CG	1:A:158:ILE:H	2.11	0.63
1:A:199:TYR:CD2	1:A:203:ILE:HG22	2.34	0.63
1:A:80:LEU:O	1:A:83:LEU:HG	2.00	0.60
1:A:21:LYS:HA	1:A:24:ILE:HD12	1.83	0.60
1:A:100:PRO:HG3	1:A:199:TYR:CE1	2.34	0.60
1:A:127:LYS:HE3	1:A:128:VAL:H	1.67	0.60
1:A:62:THR:O	1:A:65:ILE:HG22	2.00	0.60
1:A:64:ILE:HA	1:A:67:ALA:HB2	1.82	0.60
1:A:179:ALA:HB1	1:A:182:ASP:H	1.66	0.59
2:0:1155:A:OP1	2:0:1156:A:O2'	2.20	0.59
1:A:144:LEU:O	1:A:149:ILE:HB	2.03	0.58
1:A:143:GLU:N	1:A:143:GLU:OE1	2.36	0.58
1:A:127:LYS:NZ	1:A:128:VAL:HG12	2.18	0.58
1:A:172:VAL:HG22	1:A:173:VAL:H	1.68	0.58
1:A:133:THR:OG1	1:A:157:LYS:NZ	2.21	0.57
1:A:164:LYS:HG3	1:A:165:VAL:N	2.19	0.57
1:A:39:PRO:HB2	1:A:42:GLN:OE1	2.04	0.57
1:A:135:MET:HB2	1:A:157:LYS:HE2	1.87	0.57
1:A:28:PRO:HG2	1:A:97:ASP:HA	1.87	0.56
1:A:157:LYS:HG2	1:A:158:ILE:N	2.14	0.56
1:A:99:ASN:HB2	1:A:100:PRO:HD2	1.89	0.55
1:A:179:ALA:HA	1:A:181:LEU:H	1.71	0.55
1:A:208:ASP:OD1	1:A:209:VAL:N	2.40	0.55
1:A:58:MET:HE1	2:0:1216:C:H4'	1.89	0.54
1:A:209:VAL:O	1:A:212:VAL:HG22	2.08	0.54
1:A:98:MET:HE2	1:A:102:LYS:HB2	1.91	0.53
1:A:135:MET:N	1:A:157:LYS:HE2	2.24	0.53
1:A:127:LYS:HD3	1:A:164:LYS:O	2.09	0.52
1:A:126:ILE:HD11	1:A:167:VAL:HB	1.91	0.52
2:0:1167:A:C6	2:0:1196:A:C2	2.97	0.52
1:A:199:TYR:HD2	1:A:203:ILE:HG22	1.74	0.52
1:A:50:ILE:HB	1:A:54:VAL:HG22	1.92	0.52

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:VAL:O	1:A:130:LYS:HE2	2.10	0.52
1:A:127:LYS:HZ1	1:A:128:VAL:HG12	1.75	0.52
1:A:144:LEU:HD21	1:A:180:VAL:HG11	1.92	0.52
1:A:130:LYS:N	1:A:130:LYS:HD3	2.26	0.51
1:A:121:ILE:HG22	1:A:122:ALA:H	1.76	0.51
1:A:199:TYR:HB3	1:A:203:ILE:O	2.11	0.51
1:A:30:VAL:H	1:A:200:GLU:CD	2.14	0.51
1:A:189:ILE:O	1:A:189:ILE:HD12	2.11	0.51
1:A:205:TYR:CE1	1:A:210:LEU:HD11	2.46	0.51
1:A:52:ASP:OD1	1:A:52:ASP:N	2.34	0.50
1:A:147:VAL:HG21	1:A:177:LEU:HD13	1.92	0.50
1:A:147:VAL:O	1:A:147:VAL:HG23	2.12	0.50
1:A:205:TYR:HE1	1:A:210:LEU:HD11	1.76	0.50
2:0:1169:G:H1	2:0:1189:C:H42	1.61	0.49
1:A:135:MET:H	1:A:157:LYS:HE2	1.78	0.49
2:0:1165:G:H1	2:0:1214:C:H42	1.60	0.49
1:A:96:THR:HG23	1:A:98:MET:N	2.26	0.48
2:0:1213:A:H8	2:0:1213:A:O5'	1.96	0.48
1:A:23:LEU:O	1:A:26:SER:OG	2.30	0.48
1:A:33:VAL:HG12	1:A:195:ILE:HG13	1.93	0.48
1:A:56:LEU:HG	1:A:94:LEU:HD13	1.95	0.48
1:A:211:LYS:O	1:A:211:LYS:HD2	2.14	0.48
1:A:35:MET:HE1	1:A:58:MET:HB2	1.96	0.48
1:A:135:MET:CB	1:A:157:LYS:HE2	2.44	0.47
2:0:1190:C:H2'	2:0:1191:U:O4'	2.14	0.47
1:A:135:MET:H	1:A:157:LYS:CE	2.28	0.47
2:0:1161:G:H2'	2:0:1162:C:H6	1.79	0.47
2:0:1210:C:H2'	2:0:1211:U:O4'	2.14	0.47
1:A:127:LYS:HA	1:A:127:LYS:HD2	1.67	0.47
1:A:68:LEU:HD21	1:A:81:ALA:HA	1.95	0.47
1:A:98:MET:HG3	1:A:99:ASN:O	2.14	0.47
1:A:135:MET:CA	1:A:157:LYS:HE2	2.45	0.47
1:A:111:LYS:HG2	1:A:192:GLY:HA3	1.97	0.47
1:A:130:LYS:HA	1:A:160:ILE:HG23	1.97	0.46
1:A:54:VAL:CB	1:A:94:LEU:HD11	2.37	0.46
1:A:212:VAL:HG23	1:A:213:ASP:N	2.31	0.46
2:0:1163:G:C6	2:0:1164:G:C5	3.03	0.46
1:A:54:VAL:HG12	1:A:96:THR:HB	1.98	0.46
1:A:127:LYS:HD2	1:A:166:VAL:HG23	1.99	0.45
1:A:147:VAL:HG21	1:A:177:LEU:CD1	2.47	0.45
1:A:193:LEU:N	1:A:193:LEU:HD12	2.31	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:LYS:NZ	1:A:162:GLU:OE2	2.50	0.45
1:A:162:GLU:OE2	1:A:164:LYS:N	2.48	0.45
1:A:128:VAL:HG13	1:A:130:LYS:HD2	1.99	0.45
1:A:186:ILE:HG22	1:A:188:PRO:HD3	1.99	0.44
1:A:206:THR:OG1	1:A:207:PRO:HD2	2.18	0.44
1:A:23:LEU:O	1:A:95:VAL:HG11	2.18	0.44
1:A:66:ARG:O	1:A:67:ALA:C	2.56	0.44
2:0:1161:G:C5	2:0:1162:C:C5	3.05	0.44
2:0:1177:A:N6	2:0:1178:G:N1	2.66	0.44
1:A:58:MET:CE	2:0:1216:C:H4'	2.47	0.44
1:A:130:LYS:HB3	1:A:160:ILE:O	2.18	0.43
1:A:206:THR:HG23	1:A:208:ASP:OD1	2.17	0.43
1:A:107:LEU:HD13	1:A:193:LEU:HB2	2.00	0.43
1:A:136:PRO:O	1:A:157:LYS:HD3	2.19	0.43
1:A:193:LEU:HD12	1:A:193:LEU:H	1.83	0.43
1:A:32:ILE:HG13	1:A:83:LEU:HD13	2.00	0.43
1:A:129:GLU:HG3	1:A:186:ILE:HD11	2.00	0.43
1:A:16:GLU:O	1:A:19:THR:N	2.52	0.43
1:A:174:SER:HB3	1:A:177:LEU:CB	2.49	0.43
1:A:199:TYR:CE2	1:A:203:ILE:HG22	2.52	0.43
2:0:1151:G:H1	2:0:1224:C:H42	1.66	0.43
2:0:1218:U:H2'	2:0:1219:C:O4'	2.19	0.43
1:A:130:LYS:HZ1	1:A:163:ASP:HA	1.84	0.42
1:A:124:CYS:C	1:A:169:LYS:HE2	2.40	0.42
1:A:68:LEU:HD13	1:A:83:LEU:HD11	2.01	0.42
1:A:125:ASP:OD1	1:A:126:ILE:N	2.53	0.42
1:A:29:VAL:HA	1:A:200:GLU:CD	2.40	0.42
1:A:179:ALA:HB3	1:A:183:ARG:HH12	1.80	0.42
1:A:61:ASN:O	1:A:65:ILE:N	2.53	0.42
2:0:1201:G:C6	2:0:1211:U:O2	2.72	0.41
2:0:1205:A:C6	2:0:1206:A:C6	3.08	0.41
1:A:29:VAL:HA	1:A:200:GLU:HG3	2.03	0.41
1:A:47:ARG:HD2	2:0:1192:U:O2'	2.19	0.41
1:A:156:GLY:O	1:A:157:LYS:HB2	2.20	0.41
1:A:168:LYS:HD2	1:A:168:LYS:HA	1.67	0.41
2:0:1186:C:H2'	2:0:1187:A:O4'	2.21	0.41
1:A:21:LYS:O	1:A:24:ILE:HB	2.21	0.41
1:A:157:LYS:HG2	1:A:158:ILE:HD12	2.03	0.41
1:A:161:LYS:H	1:A:161:LYS:HG2	1.54	0.41
1:A:46:ILE:HG21	1:A:193:LEU:HD11	2.02	0.41
1:A:176:LYS:CD	1:A:178:ALA:HB3	2.41	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:MET:HB3	1:A:136:PRO:HD2	2.03	0.40
1:A:143:GLU:HA	1:A:146:SER:HB3	2.02	0.40
1:A:154:GLU:CG	1:A:155:LYS:H	2.34	0.40
1:A:66:ARG:HG3	1:A:67:ALA:N	2.34	0.40
1:A:178:ALA:O	1:A:181:LEU:HD13	2.21	0.40
1:A:199:TYR:HD1	1:A:199:TYR:HA	1.79	0.40
1:A:31:ALA:HB3	1:A:94:LEU:HB3	2.03	0.40
1:A:68:LEU:O	1:A:72:ALA:N	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	203/213 (95%)	181 (89%)	22 (11%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	173/180 (96%)	163 (94%)	10 (6%)	20 50

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	MET
1	A	73	GLU
1	A	160	ILE
1	A	172	VAL
1	A	177	LEU
1	A	183	ARG
1	A	187	LYS
1	A	199	TYR
1	A	209	VAL
1	A	212	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	0	73/74 (98%)	12 (16%)	0

All (12) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	0	1155	A
2	0	1157	G
2	0	1171	U
2	0	1172	G
2	0	1180	A
2	0	1198	A
2	0	1199	G
2	0	1205	A
2	0	1207	C
2	0	1211	U
2	0	1221	A
2	0	1222	G

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	205/213 (96%)	0.06	9 (4%) 34 33	84, 129, 174, 229	1 (0%)
2	0	74/74 (100%)	0.35	1 (1%) 75 75	103, 145, 187, 217	0
All	All	279/287 (97%)	0.13	10 (3%) 42 40	84, 133, 177, 229	1 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	163	ASP	6.2
1	A	129	GLU	4.4
1	A	213	ASP	4.3
1	A	89	ARG	4.1
1	A	162	GLU	2.9
1	A	90	GLY	2.6
1	A	205	TYR	2.5
1	A	165	VAL	2.5
1	A	125	ASP	2.4
2	0	1151	G	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	0	1302	1/1	0.91	0.28	98,98,98,98	0
3	MG	0	1301	1/1	0.95	0.07	119,119,119,119	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.