



Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 10:56 pm BST

PDB ID : 5DEM
Title : Structure of Pseudomonas aeruginosa LpxA
Authors : Smith, E.W.; Chen, Y.
Deposited on : 2015-08-25
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	 3% 94% 6%
1	B	258	 93% 7%
1	C	258	 3% 95% . .
1	D	258	 98% .
1	E	258	 94% 5%
1	F	258	 3% 94% 5%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 12653 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acyl-[acyl-carrier-protein]--UDP-N-acetylglucosamine O-acyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	257	Total 1985	C 1241	N 368	O 370	S 6	0	2	0
1	B	257	Total 1973	C 1234	N 364	O 369	S 6	0	1	0
1	C	256	Total 1960	C 1227	N 362	O 365	S 6	0	0	0
1	D	257	Total 1975	C 1235	N 365	O 369	S 6	0	1	0
1	E	257	Total 1966	C 1230	N 363	O 367	S 6	0	0	0
1	F	257	Total 1973	C 1234	N 364	O 369	S 6	0	1	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	133	Total	O	0	0
			133	133		
3	B	171	Total	O	0	0
			171	171		
3	C	74	Total	O	0	0
			74	74		
3	D	152	Total	O	0	0
			152	152		
3	E	137	Total	O	0	0
			137	137		
3	F	149	Total	O	0	0
			149	149		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acyl-[acyl-carrier-protein]--UDP-N-acetylglucosamine O-acyltransferase



- Molecule 1: Acyl-[acyl-carrier-protein]--UDP-N-acetylglucosamine O-acyltransferase



- Molecule 1: Acyl-[acyl-carrier-protein]--UDP-N-acetylglucosamine O-acyltransferase





- Molecule 1: Acyl-[acyl-carrier-protein]--UDP-N-acetylglucosamine O-acyltransferase



- Molecule 1: Acyl-[acyl-carrier-protein]--UDP-N-acetylglucosamine O-acyltransferase



- Molecule 1: Acyl-[acyl-carrier-protein]--UDP-N-acetylglucosamine O-acyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.03Å 82.18Å 220.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	110.26 – 1.81 40.39 – 1.81	Depositor EDS
% Data completeness (in resolution range)	97.9 (110.26-1.81) 97.9 (40.39-1.81)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.72 (at 1.81Å)	Xtrriage
Refinement program	REFMAC 5.8.0124	Depositor
R, R_{free}	0.184 , 0.211 0.193 , 0.215	Depositor DCC
R_{free} test set	6531 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtrriage
Anisotropy	0.097	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.040 for k,h,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12653	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	0/2028	0.74	0/2751
1	B	0.62	0/2015	0.76	1/2734 (0.0%)
1	C	0.58	0/2002	0.74	0/2716
1	D	0.59	0/2017	0.76	0/2736
1	E	0.59	0/2008	0.77	2/2724 (0.1%)
1	F	0.66	0/2015	0.74	0/2734
All	All	0.61	0/12085	0.75	3/16395 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	176	ASP	CB-CG-OD2	5.86	123.57	118.30
1	E	199	ARG	NE-CZ-NH2	-5.71	117.45	120.30
1	E	199	ARG	NE-CZ-NH1	5.64	123.12	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1985	0	1948	14	0
1	B	1973	0	1941	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1960	0	1930	8	0
1	D	1975	0	1942	4	0
1	E	1966	0	1935	15	0
1	F	1973	0	1941	11	0
2	B	5	0	0	0	0
3	A	133	0	0	2	0
3	B	171	0	0	3	0
3	C	74	0	0	0	0
3	D	152	0	0	0	0
3	E	137	0	0	3	0
3	F	149	0	0	2	0
All	All	12653	0	11637	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100[B]:GLN:H	1:A:100[B]:GLN:HE21	1.12	0.97
1:A:100[B]:GLN:H	1:A:100[B]:GLN:NE2	1.78	0.82
1:A:209:HIS:CD2	1:A:213:ARG:NH1	2.48	0.80
1:A:209:HIS:CD2	1:A:213:ARG:HH11	2.05	0.75
1:E:102:ARG:NH1	3:E:301:HOH:O	2.25	0.69
1:D:156:HIS:HE1	3:E:415:HOH:O	1.77	0.68
1:C:15:ARG:NH2	1:C:33:GLU:OE1	2.28	0.65
1:F:176:ASP:OD2	1:F:252:ALA:HA	1.97	0.64
1:B:209:HIS:CE1	3:B:422:HOH:O	2.56	0.58
1:E:156:HIS:CE1	1:F:169:MET:HE2	2.41	0.56
1:E:213:ARG:HG2	1:E:213:ARG:HH11	1.71	0.55
1:E:15:ARG:C	1:E:16:LEU:HD12	2.27	0.55
1:B:240:PRO:HG3	1:D:72:LYS:HD2	1.88	0.55
1:A:100[B]:GLN:N	1:A:100[B]:GLN:HE21	1.95	0.54
1:A:101:ASP:OD2	1:A:140:HIS:ND1	2.31	0.53
1:D:113:ILE:HD13	1:D:119:ILE:HD11	1.90	0.53
1:E:189:GLU:HG2	3:E:356:HOH:O	2.07	0.53
1:F:71:LEU:HD22	1:F:71:LEU:N	2.24	0.53
1:E:70:ASP:OD1	1:E:72:LYS:HG2	2.08	0.53
1:C:164:HIS:CE1	1:C:201:ARG:HG3	2.44	0.52
1:B:77:PRO:HA	3:B:512:HOH:O	2.09	0.52
1:B:70:ASP:OD1	1:B:72:LYS:HG2	2.09	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:ASP:OD2	1:A:74:LYS:HD3	2.11	0.51
1:C:164:HIS:HE1	1:C:201:ARG:HD2	1.75	0.51
1:C:70:ASP:OD1	1:C:72:LYS:HG2	2.10	0.51
1:F:166:PHE:HB3	1:F:182[B]:THR:HG22	1.94	0.50
1:F:191:ARG:HD2	3:F:526:HOH:O	2.12	0.50
1:B:202:GLY:HA3	1:D:76:GLU:O	2.13	0.49
1:C:113:ILE:HD13	1:C:119:ILE:HD11	1.94	0.49
1:E:253:THR:HG23	1:E:254:ARG:HG3	1.94	0.48
1:A:71:LEU:HD23	1:A:71:LEU:N	2.28	0.48
1:A:113:ILE:HD13	1:A:119:ILE:HD11	1.96	0.48
1:A:156[A]:HIS:CE1	1:C:169:MET:HG2	2.49	0.48
1:B:237:ALA:O	1:E:200:ARG:HD3	2.15	0.46
1:A:121:HIS:HE1	1:A:140:HIS:NE2	2.14	0.46
1:B:113:ILE:HD13	1:B:119:ILE:HD11	1.96	0.46
1:B:157:GLN:HG3	3:B:556:HOH:O	2.16	0.46
1:B:76:GLU:HB3	1:B:77:PRO:HD2	1.98	0.46
1:F:113:ILE:HD13	1:F:119:ILE:HD11	1.98	0.45
1:F:76:GLU:HB3	1:F:77:PRO:HD2	1.98	0.44
1:A:254:ARG:HB2	3:A:305:HOH:O	2.16	0.44
1:A:156[A]:HIS:NE2	1:C:169:MET:HG2	2.33	0.44
1:B:234:GLU:HG3	1:E:199:ARG:O	2.18	0.44
1:E:76:GLU:HB3	1:E:77:PRO:HD2	2.00	0.44
1:B:192:SER:HA	1:B:215:TYR:CZ	2.54	0.43
1:B:238:GLN:HG2	1:E:200:ARG:HG2	2.00	0.43
1:B:233:ALA:HB1	1:E:199:ARG:HD3	2.01	0.43
1:F:71:LEU:HD22	1:F:71:LEU:H	1.85	0.42
1:F:149:LEU:HD23	3:F:503:HOH:O	2.20	0.42
1:C:36:GLU:HG2	1:C:54:LYS:HD2	2.02	0.41
1:E:16:LEU:HD12	1:E:16:LEU:N	2.35	0.41
1:E:213:ARG:HG2	1:E:213:ARG:NH1	2.33	0.41
3:A:428:HOH:O	1:B:71:LEU:HD21	2.20	0.41
1:A:57:ARG:HG2	1:A:59:TYR:CE1	2.56	0.41
1:E:156:HIS:CE1	1:F:169:MET:CE	3.04	0.40
1:F:195:PHE:CZ	1:F:212:ARG:HG2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/258 (100%)	251 (98%)	6 (2%)	0	100	100
1	B	256/258 (99%)	247 (96%)	9 (4%)	0	100	100
1	C	254/258 (98%)	244 (96%)	10 (4%)	0	100	100
1	D	256/258 (99%)	250 (98%)	6 (2%)	0	100	100
1	E	255/258 (99%)	249 (98%)	6 (2%)	0	100	100
1	F	256/258 (99%)	246 (96%)	10 (4%)	0	100	100
All	All	1534/1548 (99%)	1487 (97%)	47 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/206 (100%)	207 (100%)	0	100	100
1	B	206/206 (100%)	206 (100%)	0	100	100
1	C	204/206 (99%)	204 (100%)	0	100	100
1	D	206/206 (100%)	206 (100%)	0	100	100
1	E	205/206 (100%)	205 (100%)	0	100	100
1	F	206/206 (100%)	206 (100%)	0	100	100
All	All	1234/1236 (100%)	1234 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	121	HIS
1	A	209	HIS
1	B	223	HIS
1	B	238	GLN
1	C	164	HIS
1	C	223	HIS
1	D	156	HIS
1	D	223	HIS
1	E	223	HIS
1	F	157	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	B	301	-	4,4,4	0.95	0	6,6,6	1.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/258 (99%)	0.00	8 (3%) 49 43	12, 21, 49, 62	0
1	B	257/258 (99%)	-0.17	0 100 100	12, 19, 34, 51	0
1	C	256/258 (99%)	0.32	7 (2%) 54 49	13, 28, 49, 67	0
1	D	257/258 (99%)	-0.08	3 (1%) 79 76	10, 19, 41, 67	0
1	E	257/258 (99%)	-0.13	3 (1%) 79 76	10, 20, 40, 55	0
1	F	257/258 (99%)	-0.04	7 (2%) 54 49	11, 18, 37, 64	0
All	All	1541/1548 (99%)	-0.02	28 (1%) 68 64	10, 21, 44, 67	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	71	LEU	4.4
1	F	74	LYS	4.2
1	F	73	TYR	3.9
1	A	73	TYR	3.7
1	C	230	ALA	3.5
1	D	253	THR	3.5
1	D	254	ARG	3.2
1	D	222	GLY	3.1
1	C	253	THR	3.1
1	F	253	THR	3.1
1	A	224	THR	3.0
1	F	18	ALA	2.9
1	F	72	LYS	2.8
1	A	253	THR	2.7
1	A	71	LEU	2.7
1	C	233	ALA	2.6
1	A	222	GLY	2.6
1	C	226	GLU	2.5
1	C	234	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	F	75	GLY	2.3
1	E	2	SER	2.3
1	E	4	ILE	2.2
1	A	223	HIS	2.2
1	C	102	ARG	2.2
1	A	221	GLN	2.1
1	C	18	ALA	2.1
1	A	229	LEU	2.1
1	E	209	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	B	301	5/5	0.81	0.31	29,34,44,51	5

6.5 Other polymers [i](#)

There are no such residues in this entry.