



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 7, 2023 – 01:57 PM EDT

PDB ID : 6DER
Title : Crystal structure of *Candida albicans* acetohydroxyacid synthase in complex with the herbicide metosulam
Authors : Garcia, M.D.; Guddat, L.W.
Deposited on : 2018-05-12
Resolution : 2.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6241 (2.14-2.10)
Clashscore	141614	6778 (2.14-2.10)
Ramachandran outliers	138981	6705 (2.14-2.10)
Sidechain outliers	138945	6706 (2.14-2.10)
RSRZ outliers	127900	6112 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	682	

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5283 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetolactate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	600	4620	2933	786	880	21	0	12	0

There are 45 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	MET	-	initiating methionine	UNP A0A1D8PJF9
A	3	HIS	-	expression tag	UNP A0A1D8PJF9
A	4	HIS	-	expression tag	UNP A0A1D8PJF9
A	5	HIS	-	expression tag	UNP A0A1D8PJF9
A	6	HIS	-	expression tag	UNP A0A1D8PJF9
A	7	HIS	-	expression tag	UNP A0A1D8PJF9
A	8	HIS	-	expression tag	UNP A0A1D8PJF9
A	9	SER	-	expression tag	UNP A0A1D8PJF9
A	10	SER	-	expression tag	UNP A0A1D8PJF9
A	11	GLY	-	expression tag	UNP A0A1D8PJF9
A	12	LEU	-	expression tag	UNP A0A1D8PJF9
A	13	VAL	-	expression tag	UNP A0A1D8PJF9
A	14	PRO	-	expression tag	UNP A0A1D8PJF9
A	15	ARG	-	expression tag	UNP A0A1D8PJF9
A	16	GLY	-	expression tag	UNP A0A1D8PJF9
A	17	SER	-	expression tag	UNP A0A1D8PJF9
A	18	GLY	-	expression tag	UNP A0A1D8PJF9
A	19	MET	-	expression tag	UNP A0A1D8PJF9
A	20	LYS	-	expression tag	UNP A0A1D8PJF9
A	21	GLU	-	expression tag	UNP A0A1D8PJF9
A	22	THR	-	expression tag	UNP A0A1D8PJF9
A	23	ALA	-	expression tag	UNP A0A1D8PJF9
A	24	ALA	-	expression tag	UNP A0A1D8PJF9
A	25	ALA	-	expression tag	UNP A0A1D8PJF9
A	26	LYS	-	expression tag	UNP A0A1D8PJF9
A	27	PHE	-	expression tag	UNP A0A1D8PJF9
A	28	GLU	-	expression tag	UNP A0A1D8PJF9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	29	ARG	-	expression tag	UNP A0A1D8PJF9
A	30	GLN	-	expression tag	UNP A0A1D8PJF9
A	31	HIS	-	expression tag	UNP A0A1D8PJF9
A	32	MET	-	expression tag	UNP A0A1D8PJF9
A	33	ASP	-	expression tag	UNP A0A1D8PJF9
A	34	SER	-	expression tag	UNP A0A1D8PJF9
A	35	PRO	-	expression tag	UNP A0A1D8PJF9
A	36	ASP	-	expression tag	UNP A0A1D8PJF9
A	37	LEU	-	expression tag	UNP A0A1D8PJF9
A	38	GLY	-	expression tag	UNP A0A1D8PJF9
A	39	THR	-	expression tag	UNP A0A1D8PJF9
A	40	ASP	-	expression tag	UNP A0A1D8PJF9
A	41	ASP	-	expression tag	UNP A0A1D8PJF9
A	42	ASP	-	expression tag	UNP A0A1D8PJF9
A	43	ASP	-	expression tag	UNP A0A1D8PJF9
A	44	LYS	-	expression tag	UNP A0A1D8PJF9
A	45	ALA	-	expression tag	UNP A0A1D8PJF9
A	46	MET	-	expression tag	UNP A0A1D8PJF9

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	53	27	9	15	2	0	0


- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total K 1 1	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0

- Molecule 5 is Metosulam (three-letter code: G87) (formula: C₁₄H₁₃Cl₂N₅O₄S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
5	A	1	Total	C	Cl	N	O	S	0	0
			26	14	2	5	4	1		

- Molecule 6 is 2-{3-[(4-AMINO-2-METHYLPYRIMIDIN-5-YL)METHYL]-4-METHYL-2-OXO-2,3-DIHYDRO-1,3-THIAZOL-5-YL}ETHYL TRIHYDROGEN DIPHOSPHATE (three-letter code: TZD) (formula: C₁₂H₁₈N₄O₈P₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
6	A	1	27	12	4	8	2	1	0	1

- Molecule 7 is (3Z)-4-[[[(4-AMINO-2-METHYLPYRIMIDIN-5-YL)METHYL]AMINO]-3-MERCAPTOPENT-3-EN-1-YL TRIHYDROGEN DIPHOSPHATE (three-letter code: TP9) (formula: C₁₁H₁₈N₄O₇P₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
7	A	1	25	11	4	7	2	1	0	1


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	530	Total 530	O 530	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetolactate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	173.81Å 173.81Å 176.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.82 – 2.13 47.82 – 2.13	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.82-2.13) 99.6 (47.82-2.13)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 2.12Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.148 , 0.166 0.152 , 0.168	Depositor DCC
R_{free} test set	1998 reflections (2.27%)	wwPDB-VP
Wilson B-factor (Å ²)	36.5	Xtrriage
Anisotropy	0.378	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5283	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, TP9, TZD, K, G87, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/4734	0.46	0/6426

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4620	0	4631	15	0
2	A	53	0	30	1	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	26	0	0	1	0
6	A	27	0	15	2	0
7	A	25	0	17	0	0
8	A	530	0	0	3	0
All	All	5283	0	4693	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:704:G87:O22	8:A:801:HOH:O	2.14	0.66
1:A:166[B]:VAL:HG23	1:A:170:MET:HE2	1.77	0.65
1:A:479:GLN:O	1:A:482:THR:HB	2.02	0.60
1:A:132:PRO:HG3	1:A:138:ALA:HB2	1.84	0.59
1:A:643:LYS:NZ	8:A:806:HOH:O	2.34	0.50
1:A:279:GLU:HA	1:A:282:SER:HB3	1.93	0.50
1:A:300:ILE:HG13	1:A:325:ILE:HG21	1.97	0.46
6:A:705[A]:TZD:OC2	8:A:802:HOH:O	2.20	0.46
1:A:567:LYS:HB3	1:A:628:PHE:CZ	2.52	0.45
1:A:99[B]:LEU:HD22	1:A:126:LYS:HB3	1.99	0.45
1:A:167:ILE:HD13	1:A:170:MET:HE1	1.98	0.45
1:A:472:LEU:HD23	1:A:571:LEU:HG	1.99	0.45
1:A:189:THR:HA	1:A:192:ILE:HD12	2.00	0.44
1:A:490:THR:HG22	1:A:513:ILE:HB	1.99	0.43
6:A:705[A]:TZD:C2	6:A:705[A]:TZD:H4'2	2.31	0.43
1:A:168:THR:HB	1:A:169:PRO:HD3	2.01	0.42
1:A:163:ALA:HB1	1:A:202:ILE:HG12	2.02	0.42
2:A:701:FAD:H9	2:A:701:FAD:H1'1	1.81	0.41
1:A:490:THR:HA	1:A:513:ILE:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	608/682 (89%)	597 (98%)	11 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	495/563 (88%)	492 (99%)	3 (1%)	86 90

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	482	THR
1	A	512	MET
1	A	551	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	TP9	A	706[B]	4	21,25,25	2.15	2 (9%)	28,36,36	1.82	6 (21%)
6	TZD	A	705[A]	4	23,28,28	3.39	6 (26%)	28,42,42	1.50	6 (21%)
2	FAD	A	701	-	53,58,58	1.62	9 (16%)	68,89,89	1.32	11 (16%)
5	G87	A	704	-	24,28,28	2.02	6 (25%)	27,42,42	3.17	8 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	TP9	A	706[B]	4	-	2/17/22/22	0/1/1/1
6	TZD	A	705[A]	4	-	1/16/17/17	0/2/2/2
2	FAD	A	701	-	-	2/30/50/50	0/6/6/6
5	G87	A	704	-	-	1/9/15/15	0/3/3/3

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	705[A]	TZD	C5-S1	-11.40	1.52	1.74
7	A	706[B]	TP9	C4-N3	8.62	1.43	1.32
6	A	705[A]	TZD	C4-N3	7.52	1.56	1.39
6	A	705[A]	TZD	C5A-C5	6.28	1.53	1.50
5	A	704	G87	O22-C18	5.15	1.43	1.35
2	A	701	FAD	C10-N1	4.74	1.42	1.33
2	A	701	FAD	C4X-N5	4.53	1.39	1.30
2	A	701	FAD	C2B-C1B	-4.34	1.47	1.53
5	A	704	G87	C20-N21	4.21	1.41	1.35
5	A	704	G87	O24-C16	3.95	1.41	1.34
7	A	706[B]	TP9	C4'-N4'	3.52	1.43	1.34
5	A	704	G87	C17-C18	3.19	1.43	1.38
6	A	705[A]	TZD	C4'-N4'	3.17	1.42	1.34
2	A	701	FAD	O3'-C3'	-2.74	1.36	1.43
2	A	701	FAD	O4B-C1B	2.55	1.44	1.41
2	A	701	FAD	C1'-N10	-2.40	1.42	1.48
6	A	705[A]	TZD	C5'-C4'	-2.38	1.38	1.42
2	A	701	FAD	C2'-C3'	2.32	1.57	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	FAD	C2-N1	2.27	1.42	1.36
6	A	705[A]	TZD	C35-N3	-2.24	1.44	1.48
5	A	704	G87	C18-N19	2.19	1.33	1.30
5	A	704	G87	C20-N19	2.16	1.43	1.33
2	A	701	FAD	C1'-C2'	2.03	1.55	1.52

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	704	G87	O12-S10-O11	-13.02	103.55	119.55
7	A	706[B]	TP9	PA-O3A-PB	-5.02	115.60	132.83
5	A	704	G87	C18-C17-C16	4.79	118.53	115.51
2	A	701	FAD	N3A-C2A-N1A	-4.11	122.26	128.68
5	A	704	G87	C23-O22-C18	-3.95	111.16	117.36
7	A	706[B]	TP9	C7'-N3-C4	-3.65	120.94	125.97
5	A	704	G87	C18-N19-C20	3.34	119.41	116.58
6	A	705[A]	TZD	N1'-C2'-N3'	-3.21	120.02	125.54
7	A	706[B]	TP9	C6'-N1'-C2'	3.20	121.40	115.96
7	A	706[B]	TP9	C5'-C6'-N1'	-3.19	118.50	123.82
6	A	705[A]	TZD	C2A-C2'-N1'	2.97	120.41	117.14
2	A	701	FAD	C4A-C5A-N7A	-2.90	106.38	109.40
2	A	701	FAD	C4-C4X-N5	2.87	122.31	118.23
2	A	701	FAD	C4X-C10-N10	2.80	120.58	116.48
6	A	705[A]	TZD	C6'-C5'-C4'	2.73	119.44	115.72
6	A	705[A]	TZD	C6'-N1'-C2'	2.69	120.55	115.96
5	A	704	G87	C17-C18-N19	-2.64	120.98	124.11
2	A	701	FAD	C4X-C4-N3	2.59	119.77	113.19
5	A	704	G87	O12-S10-N09	2.48	112.94	106.73
2	A	701	FAD	C10-C4X-N5	-2.46	119.63	124.86
2	A	701	FAD	C4-N3-C2	-2.45	121.11	125.64
6	A	705[A]	TZD	C5'-C6'-N1'	-2.42	119.78	123.82
2	A	701	FAD	O4-C4-C4X	-2.42	120.18	126.60
5	A	704	G87	C13-N14-N15	2.38	106.91	102.93
6	A	705[A]	TZD	C5'-C35-N3	-2.32	109.43	113.26
5	A	704	G87	O11-S10-C13	2.30	111.22	108.71
7	A	706[B]	TP9	N1'-C2'-N3'	-2.17	121.80	125.54
7	A	706[B]	TP9	C6'-C5'-C4'	2.16	118.67	115.72
2	A	701	FAD	C10-N1-C2	2.16	121.22	116.90
2	A	701	FAD	C9A-C5X-N5	-2.14	120.10	122.43
2	A	701	FAD	C4X-C10-N1	-2.12	119.80	124.73

There are no chirality outliers.

All (6) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
6	A	705[A]	TZD	P1-O11-P2-O21
7	A	706[B]	TP9	C4-C5-C6-C7
2	A	701	FAD	P-O3P-PA-O5B
7	A	706[B]	TP9	PB-O3A-PA-O7
5	A	704	G87	C04-N09-S10-O11
2	A	701	FAD	O4B-C4B-C5B-O5B



There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	705[A]	TZD	2	0
2	A	701	FAD	1	0
5	A	704	G87	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	600/682 (87%)	-0.08	30 (5%) 28 34	26, 38, 63, 108	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	277	VAL	5.2
1	A	276	ALA	4.5
1	A	681	GLY	3.9
1	A	80	LEU	3.8
1	A	442	GLU	3.7
1	A	280	PHE	3.5
1	A	437	ILE	3.5
1	A	441	PRO	3.4
1	A	436[A]	SER	3.2
1	A	279	GLU	3.1
1	A	677	GLU	3.0
1	A	438	GLU	2.9
1	A	99[A]	LEU	2.8
1	A	680	GLY	2.7
1	A	482	THR	2.7
1	A	79	GLN	2.6
1	A	77	LYS	2.6
1	A	278	SER	2.5
1	A	125	ASP	2.4
1	A	439	ASN	2.4
1	A	473	ILE	2.4
1	A	682	LYS	2.3
1	A	296	LYS	2.3
1	A	670	GLN	2.2
1	A	476	ILE	2.1
1	A	284	ALA	2.1
1	A	679	THR	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	75	LYS	2.0
1	A	431	ILE	2.0
1	A	83	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	G87	A	704	26/26	0.94	0.10	43,56,63,77	0
2	FAD	A	701	53/53	0.98	0.09	25,33,38,49	0
7	TP9	A	706[B]	25/25	0.98	0.11	27,31,34,37	25
3	K	A	702	1/1	0.99	0.10	39,39,39,39	0
6	TZD	A	705[A]	27/27	0.99	0.11	24,29,40,43	27
4	MG	A	703	1/1	0.99	0.08	29,29,29,29	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around G87 A 704:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD A 701:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around TP9 A 706 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around TZD A 705 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.