

# Full wwPDB X-ray Structure Validation Report (i)

#### May 20, 2021 – 02:10 PM JST

PDB ID	:	7DIE
Title	:	Crystal structure of M. penetrans Ferritin
Authors	:	Wang, w.m.; Zhang, y.; Wang, h.f.
Deposited on		
Resolution	:	1.90  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.18
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.18

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Matria	Whole archive	Similar resolution
Metric	$(\# {\rm Entries})$	$(\# \text{Entries}, \text{ resolution range}(\text{\AA}))$
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	А	179	% <b>88%</b>	•• 8%					
1	В	179	88%	•• 8%					
1	С	179	87%	• • 8%					
1	D	179	2% <b>8</b> 7%	• • 8%					
1	Е	179	% 87%	•• 8%					
1	F	179	87%	• • 8%					



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 8877 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	164	Total	С	Ν	0	S	0	5	0
	Л	104	1362	869	223	262	8	0	5	0
1	В	164	Total	С	Ν	Ο	$\mathbf{S}$	0	4	0
	D	104	1351	862	221	260	8	0	4	0
1	С	164	Total	С	Ν	Ο	S	0	3	0
	U	104	1345	859	220	258	8	0	0	0
1	D	164	Total	С	Ν	Ο	$\mathbf{S}$	0	5	0
	D	104	1362	869	223	262	8	0	5	0
1	Е	164	Total	С	Ν	Ο	$\mathbf{S}$	0	3	0
	Ľ	104	1345	859	220	258	8	0	5	0
1	F	164	Total	С	Ν	Ο	S	0	2	0
	T,	104	1337	855	218	256	8			0

• Molecule 1 is a protein called Ferritin.

• Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	8	Total Fe 8 8	0	0
2	В	6	Total Fe 6 6	0	0
2	С	4	Total Fe 4 4	0	0
2	D	3	Total Fe 3 3	0	0
2	Ε	6	Total Fe 6 6	0	0
2	F	9	Total Fe 9 9	0	0

• Molecule 3 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	123	Total O 123 123	0	0
3	В	122	Total O   122 122	0	0
3	С	123	Total O 123 123	0	0
3	D	121	Total O   121 121	0	0
3	Е	129	Total O 129 129	0	0
3	F	121	Total O   121 121	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Ferritin







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 4	Depositor
Cell constants	151.33Å 151.33Å 125.21Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	47.86 - 1.90	Depositor
Resolution (A)	47.86 - 1.90	EDS
% Data completeness	99.6 (47.86-1.90)	Depositor
(in resolution range)	99.6 (47.86-1.90)	EDS
R <sub>merge</sub>	(Not available)	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.68 (at 1.90 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
D D.	0.153 , $0.177$	Depositor
$R, R_{free}$	0.154 , $0.177$	DCC
$R_{free}$ test set	2000 reflections $(1.81%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	31.7	Xtriage
Anisotropy	0.284	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.34 , $49.7$	EDS
L-test for twinning <sup>2</sup>	$< L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	0.012 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8877	wwPDB-VP
Average B, all atoms $(Å^2)$	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.87% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
IVIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.48	0/1384	0.52	0/1866	
1	В	0.50	0/1373	0.52	0/1852	
1	С	0.48	0/1367	0.50	0/1844	
1	D	0.49	0/1384	0.51	0/1866	
1	Ε	0.53	1/1367~(0.1%)	0.52	0/1844	
1	F	0.45	0/1359	0.51	0/1833	
All	All	0.49	1/8234~(0.0%)	0.51	0/11105	

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	Е	140	GLU	CD-OE2	-5.44	1.19	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1362	0	1363	7	0
1	В	1351	0	1352	6	0
1	С	1345	0	1348	8	0
1	D	1362	0	1363	10	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Е	1345	0	1348	9	0
1	F	1337	0	1343	6	0
2	А	8	0	0	0	0
2	В	6	0	0	0	0
2	С	4	0	0	0	0
2	D	3	0	0	0	0
2	Ε	6	0	0	0	0
2	F	9	0	0	0	0
3	А	123	0	0	3	0
3	В	122	0	0	1	0
3	С	123	0	0	1	0
3	D	121	0	0	3	0
3	Е	129	0	0	4	0
3	F	121	0	0	2	0
All	All	8877	0	8117	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All $(43)$ close contacts	within the	ne same	asymmetric	$\operatorname{unit}$	are listed	below,	sorted	by their	$\operatorname{clash}$
magnitude.									

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:55[B]:ASP:OD1	3:A:301:HOH:O	1.81	0.97
1:F:55[B]:ASP:OD1	3:F:301:HOH:O	1.86	0.93
1:E:141:GLU:OE2	3:E:301:HOH:O	2.04	0.76
1:D:141:GLU:OE1	3:D:301:HOH:O	2.05	0.74
1:E:55[B]:ASP:OD1	3:E:302:HOH:O	2.08	0.72
1:F:155:ASN:ND2	1:F:158:ASN:H	1.91	0.68
1:C:111[A]:ASN:ND2	3:C:301:HOH:O	2.27	0.67
1:B:14:ASP:O	1:B:18:LYS:HG3	1.96	0.66
1:B:55[B]:ASP:OD1	3:B:301:HOH:O	2.12	0.65
1:D:155:ASN:ND2	1:D:158:ASN:H	1.94	0.65
1:E:104:LYS:NZ	3:E:304:HOH:O	2.28	0.62
1:A:155:ASN:ND2	1:A:158:ASN:H	1.98	0.61
1:E:155:ASN:ND2	1:E:158:ASN:H	2.00	0.59
1:E:155:ASN:HD22	1:E:155:ASN:C	2.04	0.59
1:A:155:ASN:HD22	1:A:155:ASN:C	2.06	0.59
1:C:155:ASN:H	1:C:158:ASN:HD22	1.50	0.59
1:E:155:ASN:H	1:E:158:ASN:HD22	1.51	0.57
1:F:155:ASN:HD22	1:F:155:ASN:C	2.07	0.57
1:B:155:ASN:ND2	1:B:158:ASN:H	2.03	0.56



		Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:D:155:ASN:H	1:D:158:ASN:HD22	1.53	0.56
1:D:155:ASN:C	1:D:155:ASN:HD22	2.09	0.56
1:B:155:ASN:H	1:B:158:ASN:HD22	1.54	0.54
1:A:155:ASN:H	1:A:158:ASN:HD22	1.55	0.54
1:C:155:ASN:ND2	1:C:158:ASN:H	2.05	0.54
1:B:155:ASN:H	1:B:158:ASN:ND2	2.07	0.51
1:D:155:ASN:H	1:D:158:ASN:ND2	2.09	0.51
1:B:155:ASN:HD22	1:B:155:ASN:C	2.15	0.50
1:E:155:ASN:H	1:E:158:ASN:ND2	2.09	0.50
1:C:155:ASN:H	1:C:158:ASN:ND2	2.09	0.50
1:C:155:ASN:C	1:C:155:ASN:HD22	2.15	0.49
1:E:111[A]:ASN:OD1	3:E:303:HOH:O	2.21	0.46
1:C:79:GLN:NE2	1:D:79:GLN:OE1	2.43	0.46
1:A:155:ASN:H	1:A:158:ASN:ND2	2.15	0.45
1:D:66[A]:LYS:NZ	3:D:307:HOH:O	2.50	0.44
1:A:141:GLU:HG3	3:A:394:HOH:O	2.17	0.43
1:D:137:GLU:OE2	1:F:70:LYS:HE3	2.18	0.43
1:A:111[A]:ASN:ND2	3:A:304:HOH:O	2.51	0.42
1:F:79:GLN:CD	1:F:79:GLN:H	2.23	0.42
1:C:101:MET:HA	1:C:104:LYS:HE2	2.01	0.41
1:F:43:ASP:HB2	3:F:397:HOH:O	2.19	0.41
1:D:153[B]:ASN:ND2	3:D:311:HOH:O	2.53	0.41
1:E:155:ASN:ND2	1:E:155:ASN:C	2.74	0.41
1:C:79:GLN:HG3	1:D:79:GLN:OE1	2.20	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles
1	А	167/179~(93%)	164 (98%)	3~(2%)	0	100 100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	В	166/179~(93%)	163~(98%)	3~(2%)	0	100	100
1	С	165/179~(92%)	163 (99%)	2(1%)	0	100	100
1	D	167/179~(93%)	165 (99%)	2(1%)	0	100	100
1	Ε	165/179~(92%)	162 (98%)	3~(2%)	0	100	100
1	F	164/179~(92%)	162 (99%)	2(1%)	0	100	100
All	All	994/1074~(93%)	979~(98%)	15~(2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	А	156/166~(94%)	153~(98%)	3~(2%)	57	53
1	В	155/166~(93%)	152~(98%)	3~(2%)	57	53
1	С	154/166~(93%)	151~(98%)	3~(2%)	57	53
1	D	156/166~(94%)	154 (99%)	2(1%)	69	68
1	Е	154/166~(93%)	152~(99%)	2(1%)	69	68
1	F	153/166~(92%)	150 (98%)	3(2%)	55	51
All	All	928/996~(93%)	912~(98%)	16 (2%)	59	57

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	17	SER
1	А	135	SER
1	А	155	ASN
1	В	135	SER
1	В	155	ASN
1	В	166	LYS
1	С	135	SER
1	С	141	GLU



Conti	Continuea from previous page						
Mol	Chain	$\mathbf{Res}$	Type				
1	С	155	ASN				
1	D	135	SER				
1	D	155	ASN				
1	Е	135	SER				
1	Е	155	ASN				
1	F	18	LYS				
1	F	135	SER				
1	F	155	ASN				

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	79	GLN
1	А	155	ASN
1	А	158	ASN
1	А	161	ASN
1	В	155	ASN
1	В	158	ASN
1	В	161	ASN
1	С	155	ASN
1	С	158	ASN
1	D	155	ASN
1	D	158	ASN
1	Е	155	ASN
1	Е	158	ASN
1	F	155	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



## 5.6 Ligand geometry (i)

Of 36 ligands modelled in this entry, 36 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	А	164/179~(91%)	-0.52	1 (0%) 89 90	25, 30, 49, 72	0
1	В	164/179~(91%)	-0.51	0 100 100	25, 30, 51, 66	0
1	С	164/179~(91%)	-0.54	0 100 100	24, 30, 48, 64	0
1	D	164/179~(91%)	-0.44	3 (1%) 68 71	24,  30,  50,  70	0
1	Ε	164/179~(91%)	-0.52	1 (0%) 89 90	26, 31, 50, 70	0
1	F	164/179~(91%)	-0.58	0 100 100	26, 31, 51, 69	0
All	All	984/1074~(91%)	-0.52	5 (0%) 91 92	24,  30,  51,  72	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	86	VAL	4.1
1	D	85	LEU	3.1
1	Е	86	VAL	2.9
1	D	84	GLU	2.3
1	А	86	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FE	Е	205	1/1	0.40	0.14	92,92,92,92	1
2	FE	F	207	1/1	0.56	0.12	101,101,101,101	1
2	FE	А	203	1/1	0.62	0.11	62,62,62,62	1
2	FE	А	202	1/1	0.73	0.23	91,91,91,91	0
2	FE	С	204	1/1	0.76	0.18	67,67,67,67	1
2	FE	С	202	1/1	0.79	0.24	93,93,93,93	0
2	FE	F	202	1/1	0.81	0.29	92,92,92,92	0
2	$\mathbf{FE}$	Ε	206	1/1	0.82	0.25	74,74,74,74	1
2	FE	F	208	1/1	0.84	0.30	71,71,71,71	1
2	$\mathbf{FE}$	Е	202	1/1	0.85	0.25	92,92,92,92	0
2	FE	F	209	1/1	0.85	0.25	$53,\!53,\!53,\!53$	1
2	$\mathbf{FE}$	В	202	1/1	0.86	0.20	89,89,89,89	0
2	$\mathbf{FE}$	В	203	1/1	0.87	0.28	$67,\!67,\!67,\!67$	1
2	$\mathbf{FE}$	В	206	1/1	0.88	0.22	87,87,87,87	1
2	FE	D	202	1/1	0.89	0.30	93,93,93,93	0
2	FE	А	207	1/1	0.89	0.41	84,84,84,84	1
2	FE	А	208	1/1	0.93	0.33	68,68,68,68	1
2	FE	F	203	1/1	0.94	0.21	64,64,64,64	1
2	FE	А	204	1/1	0.95	0.13	56, 56, 56, 56	1
2	FE	Е	203	1/1	0.96	0.22	$55,\!55,\!55,\!55$	1
2	FE	F	201	1/1	0.98	0.06	36,36,36,36	1
2	$\mathbf{FE}$	С	203	1/1	0.98	0.06	39,39,39,39	1
2	$\mathbf{FE}$	А	201	1/1	0.98	0.07	36,36,36,36	1
2	FE	F	206	1/1	0.98	0.06	40,40,40,40	1
2	$\mathbf{FE}$	Е	204	1/1	0.98	0.07	36,36,36,36	1
2	FE	D	201	1/1	0.98	0.06	36,36,36,36	1
2	FE	А	206	1/1	0.98	0.11	36,36,36,36	1
2	FE	В	205	1/1	0.99	0.07	35,35,35,35	1
2	FE	D	203	1/1	0.99	0.09	36,36,36,36	1
2	FE	Е	201	1/1	0.99	0.04	45,45,45,45	0
2	FE	В	201	1/1	0.99	0.07	34,34,34,34	1
2	FE	В	204	1/1	0.99	0.05	31,31,31,31	1
2	FE	С	201	1/1	1.00	0.05	$35,\!35,\!35,\!35$	1
2	FE	А	205	1/1	1.00	0.03	32,32,32,32	1
2	FE	F	204	1/1	1.00	0.33	$55,\!55,\!55,\!55$	1
2	$\mathbf{FE}$	F	205	1/1	1.00	0.04	35,35,35,35	1




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















































































































































































## 6.5 Other polymers (i)

There are no such residues in this entry.

