

Full wwPDB X-ray Structure Validation Report (i)

Jun 1, 2020 – 11:42 pm BST

PDB ID : 1DJP

Title : CRYSTAL STRUCTURE OF PSEUDOMONAS 7A GLUTAMINASE-ASP

ARAGINASE WITH THE INHIBITOR DON COVALENTLY BOUND IN

THE ACTIVE SITE

Authors: Ortlund, E.; Lacount, M.W.; Lewinski, K.; Lebioda, L.

Deposited on : 1999-12-03

Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at $\frac{\text{https://www.wwpdb.org/validation/2017/XrayValidationReportHelp}}{\text{with specific help available everywhere you see the (i) symbol.}$

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	A	330	84%	15%	•
1	В	330	88%	12%	



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5201 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called GLUTAMINASE-ASPARAGINASE.


	\mathbf{Mol}	Chain	Residues	${f Atoms}$			ZeroOcc	AltConf	Trace		
	1	Λ	220	Total	С	N	О	S	0	0	0
	1	1 A 330	330	2485	1550	440	487	8	0	U	
ĺ	1	D	330	Total	С	N	О	S	0	0	0
	1	Б	330	2485	1550	440	487	8	0	U	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1111	ASN	ASP	SEE REMARK 999	UNP P10182
A	1113	VAL	THR	SEE REMARK 999	UNP P10182
A	1114	GLN	LEU	SEE REMARK 999	UNP P10182
A	1115	LYS	ASN	SEE REMARK 999	UNP P10182
A	1263	VAL	LEU	SEE REMARK 999	UNP P10182
A	1264	VAL	THR	SEE REMARK 999	UNP P10182
A	1269	GLN	THR	SEE REMARK 999	UNP P10182
A	1273	ASN	THR	SEE REMARK 999	UNP P10182
A	1317	ALA	VAL	SEE REMARK 999	UNP P10182
A	1318	MET	GLU	SEE REMARK 999	UNP P10182
A	1319	VAL	LEU	SEE REMARK 999	UNP P10182
A	1322	THR	VAL	SEE REMARK 999	UNP P10182
В	3111	ASN	ASP	SEE REMARK 999	UNP P10182
В	3113	VAL	THR	SEE REMARK 999	UNP P10182
В	3114	GLN	LEU	SEE REMARK 999	UNP P10182
В	3115	LYS	ASN	SEE REMARK 999	UNP P10182
В	3263	VAL	LEU	SEE REMARK 999	UNP P10182
В	3264	VAL	THR	SEE REMARK 999	UNP P10182
В	3269	GLN	THR	SEE REMARK 999	UNP P10182
В	3273	ASN	THR	SEE REMARK 999	UNP P10182
В	3317	ALA	VAL	SEE REMARK 999	UNP P10182
В	3318	MET	GLU	SEE REMARK 999	UNP P10182
В	3319	VAL	LEU	SEE REMARK 999	UNP P10182
В	3322	THR	VAL	SEE REMARK 999	UNP P10182



• Molecule 2 is 5,5-dihydroxy-6-oxo-L-norleucine (three-letter code: DO2) (formula: $C_6H_{11}NO_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 10				0	0
2	В	1	Total 10	C 6		O 3	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	102	Total O 102 102	0	0
3	В	109	Total O 109 109	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: GLUTAMINASE-ASPARAGINASE





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	C 2 2 21	Depositor	
Cell constants	78.51Å 135.91Å 137.58Å	Depositor	
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	48.35 - 1.90	Depositor	
% Data completeness	80.7 (48.35-1.90)	Depositor	
(in resolution range)	00.7 (10.00 1.00)		
R_{merge}	0.08	Depositor	
R_{sym}	(Not available)	Depositor	
Refinement program	CNS	Depositor	
R, R_{free}	0.189 , 0.228	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	5201	wwPDB-VP	
Average B, all atoms (Å ²)	23.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: DO2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.34	0/2520	0.62	1/3413 (0.0%)	
1	В	0.35	0/2520	0.63	0/3413	
All	All	0.34	0/5040	0.63	$1/6826 \ (0.0\%)$	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
1	A	1158	LEU	CA-CB-CG	5.39	127.69	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2485	0	2503	24	0
1	В	2485	0	2503	25	0
2	A	10	0	8	0	0
2	В	10	0	7	5	0
3	A	102	0	0	1	0
3	В	109	0	0	0	0
All	All	5201	0	5021	47	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1159:VAL:HG22	1:A:1166:GLN:HB2	1.64	0.78
1:A:1039:VAL:HG13	1:A:1043:LYS:HD3	1.67	0.76
1:A:1021:ILE:H	1:A:1021:ILE:HD12	1.62	0.64
1:B:3150:LYS:HE2	1:B:3153:ARG:NH2	2.12	0.63
1:B:3207:HIS:H	1:B:3210:ASN:ND2	1.95	0.63
1:A:1150:LYS:HD2	1:A:1153:ARG:HD2	1.82	0.60
1:A:1323:LYS:NZ	1:A:1323:LYS:HB3	2.17	0.59
1:A:1058:ARG:HB2	1:A:1058:ARG:HH21	1.69	0.56
1:B:3304:VAL:HG12	1:B:3306:HIS:H	1.69	0.56
1:A:1229:SER:HB3	1:A:1253:GLY:HA2	1.86	0.56
1:B:3043:LYS:HD3	1:B:3043:LYS:O	2.06	0.55
1:A:1048:VAL:O	1:A:1051:LEU:HB2	2.06	0.55
1:B:3010:ALA:O	1:B:3055:ALA:HB1	2.07	0.55
1:B:3034:TYR:CE2	2:B:4001:DO2:HG	2.40	0.55
1:B:3034:TYR:CE2	2:B:4001:DO2:CG	2.88	0.55
1:B:3229:SER:HB3	1:B:3253:GLY:HA2	1.90	0.53
1:B:3048:VAL:O	1:B:3051:LEU:HB2	2.08	0.52
1:A:1010:ALA:O	1:A:1055:ALA:HB1	2.09	0.52
1:A:1226:ILE:HG12	1:A:1250:ILE:HB	1.92	0.51
1:B:3078:LEU:O	1:B:3082:VAL:HG23	2.12	0.50
1:B:3016:ALA:HA	1:B:3097:THR:OG1	2.13	0.49
1:B:3020:THR:CB	2:B:4001:DO2:OE1	2.57	0.48
1:B:3226:ILE:HG12	1:B:3250:ILE:HB	1.97	0.47
1:B:3291:ARG:HA	1:B:3304:VAL:HG21	1.97	0.46
1:B:3008:LYS:HG2	1:B:3009:LEU:H	1.82	0.45
1:A:1115:LYS:HE2	1:A:1211:SER:O	2.17	0.45
1:A:1155:LYS:HD3	1:A:1199:TRP:CZ3	2.52	0.45
1:A:1315:ILE:HD13	1:A:1318:MET:CE	2.46	0.45
1:B:3048:VAL:HB	1:B:3051:LEU:HD22	2.00	0.44
1:A:1175:ILE:HG21	1:A:1178:LYS:HE2	1.98	0.44
1:A:1021:ILE:H	1:A:1021:ILE:CD1	2.24	0.43
1:A:1290:LEU:HD23	1:A:1293:ALA:HB2	2.00	0.43
1:A:1262:ARG:NH1	1:B:3071:THR:HG23	2.34	0.42
1:B:3207:HIS:H	1:B:3210:ASN:HD21	1.63	0.42
1:A:1292:ASN:HA	1:A:1295:GLN:O	2.19	0.42
1:B:3020:THR:HB	2:B:4001:DO2:OE1	2.20	0.42
1:B:3034:TYR:HE2	2:B:4001:DO2:CG	2.32	0.41

Continued on next page...



~ · · · · · · · · · · · · · · · · · · ·	e		
Continued	trom	nremous	naae
-	110116	picolous	puyc

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:A:1265:PRO:HG2	3:A:307:HOH:O	2.20	0.41
1:A:1067:SER:HB2	1:A:1102:THR:OG1	2.21	0.41
1:A:1076:LEU:O	1:A:1080:LYS:HG3	2.21	0.41
1:B:3271:ARG:HA	1:B:3275:THR:O	2.21	0.41
1:A:1309:ASN:HB2	1:A:1310:PRO:CD	2.51	0.41
1:B:3270:LEU:HA	1:B:3270:LEU:HD12	1.91	0.41
1:B:3050:GLU:H	1:B:3050:GLU:CD	2.24	0.41
1:B:3137:MET:N	1:B:3137:MET:SD	2.91	0.40
1:A:1118:LYS:HA	1:A:1119:PRO:HD3	1.96	0.40
1:A:1226:ILE:HD12	1:B:3230:TYR:CD1	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percen	tiles
1	A	$328/330 \ (99\%)$	317 (97%)	10 (3%)	1 (0%)	41	31
1	В	328/330~(99%)	319 (97%)	8 (2%)	1 (0%)	41	31
All	All	656/660 (99%)	636 (97%)	18 (3%)	2 (0%)	41	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1208	THR
1	В	3208	THR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	$265/265 \; (100\%)$	250 (94%)	15 (6%)	20 11		
1	В	$265/265 \; (100\%)$	254 (96%)	11 (4%)	30 20		
All	All	$530/530 \; (100\%)$	504 (95%)	26 (5%)	25 15		

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1015	LEU
1	A	1021	ILE
1	A	1035	GLN
1	A	1042	ASP
1	A	1058	ARG
1	A	1085	LEU
1	A	1137	MET
1	A	1140	LEU
1	A	1158	LEU
1	A	1241	LEU
1	A	1267	LEU
1	A	1304	VAL
1	A	1323	LYS
1	A	1325	GLN
1	A	1330	LEU
1	В	3058	ARG
1	В	3085	LEU
1	В	3115	LYS
1	В	3140	LEU
1	В	3166	GLN
1	В	3209	VAL
1	В	3210	ASN
1	В	3241	LEU
1	В	3247	LYS
1	В	3270	LEU
1	В	3290	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

\mathbb{N}	Iol	Chain	${ m Res}$	\mathbf{Type}
	1	A	1035	GLN

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	A	1149	ASN
1	A	1166	GLN
1	A	1244	ASN
1	A	1269	GLN
1	A	1273	ASN
1	A	1285	GLN
1	A	1306	HIS
1	В	3166	GLN
1	В	3210	ASN
1	В	3217	GLN
1	В	3223	GLN
1	В	3269	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	\mathbf{B}_{0}	ond leng	gths	В	ond ang	gles
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DO2	A	4000	1	5,9,11	0.48	0	4,10,15	1.79	2 (50%)
2	DO2	В	4001	1	5,9,11	0.50	0	4,10,15	1.86	2 (50%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DO2	A	4000	1	-	3/4/9/13	-
2	DO2	В	4001	1	-	3/4/9/13	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}(^{o})$
2	A	4000	DO2	OE1-CE-CD	-2.68	109.33	126.89
2	В	4001	DO2	OE1-CE-CD	-2.67	109.36	126.89
2	В	4001	DO2	CB-CG-CD	-2.26	109.58	114.49
2	A	4000	DO2	CB-CG-CD	-2.07	109.99	114.49

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	4000	DO2	C-CA-CB-CG
2	A	4000	DO2	N-CA-CB-CG
2	A	4000	DO2	CE-CD-CG-CB
2	В	4001	DO2	C-CA-CB-CG
2	В	4001	DO2	N-CA-CB-CG
2	В	4001	DO2	CE-CD-CG-CB

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	4001	DO2	5	0

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

