



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 06:45 am BST

PDB ID : 2DKA  
Title : Crystal structure of N-acetylglucosamine-phosphate mutase, a member of the alpha-D-phosphohexomutase superfamily, in the apo-form  
Authors : Nishitani, Y.; Maruyama, D.; Nonaka, T.; Kita, A.; Fukami, T.A.; Mio, T.; Yamada-Okabe, H.; Yamada-Okabe, T.; Miki, K.  
Deposited on : 2006-04-07  
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	544	 6% 80% 15% • 5%
1	B	544	 3% 70% 14% 15%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8413 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoacetylglucosamine mutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	519	Total 4027	C 2568	N 663	O 788	S 8	0	0	0
1	B	461	Total 3584	C 2290	N 586	O 701	S 7	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	408	Total 408	O 408	0	0
2	B	394	Total 394	O 394	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.17Å 130.22Å 77.96Å 90.00° 106.68° 90.00°	Depositor
Resolution (Å)	29.56 – 1.93 29.56 – 1.93	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.56-1.93) 99.7 (29.56-1.93)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.84 (at 1.93Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.192 , 0.222 0.193 , 0.189	Depositor DCC
$R_{free}$ test set	4285 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.3	Xtrriage
Anisotropy	0.271	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 60.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8413	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/4104	0.56	0/5572
1	B	0.31	0/3659	0.57	0/4973
All	All	0.31	0/7763	0.57	0/10545

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4027	0	3949	63	0
1	B	3584	0	3519	48	0
2	A	408	0	0	8	0
2	B	394	0	0	2	0
All	All	8413	0	7468	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:338:ASN:HD21	1:A:381:ASP:H	1.22	0.87

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:ALA:HA	1:A:76:LYS:HE2	1.56	0.87
1:A:338:ASN:ND2	1:A:381:ASP:H	1.82	0.77
1:A:338:ASN:HD22	1:A:338:ASN:C	1.87	0.76
1:B:57:LYS:HE3	1:B:80:PRO:HG2	1.69	0.75
1:A:482:ASN:ND2	1:A:485:ARG:HH21	1.88	0.72
1:A:482:ASN:HD21	1:A:485:ARG:HH21	1.36	0.71
1:B:32:LYS:HD3	1:B:72:ASP:OD1	1.94	0.67
1:A:22:PHE:H	1:A:98:ASN:HD21	1.46	0.64
1:A:138:ALA:HB1	1:A:170:PHE:HB2	1.80	0.64
1:B:136:VAL:HG12	1:B:165:GLN:HB3	1.81	0.62
1:A:409:PRO:HG3	1:A:415:ALA:HA	1.82	0.61
1:B:189:LYS:HE3	2:B:582:HOH:O	2.01	0.60
1:A:464:PRO:HD2	1:A:526:ALA:O	2.03	0.59
1:B:54:LEU:HB3	1:B:57:LYS:HE2	1.86	0.57
1:B:464:PRO:HD2	1:B:526:ALA:O	2.05	0.57
1:A:11:LEU:HD11	1:A:101:PRO:HB3	1.87	0.57
1:A:3:ILE:HD12	1:A:160:PRO:HD3	1.88	0.55
1:A:76:LYS:HD3	2:A:685:HOH:O	2.06	0.55
1:A:65:ALA:HB3	1:A:142:ARG:CZ	2.36	0.55
1:B:169:LEU:HG	1:B:227:VAL:HG23	1.87	0.55
1:A:36:LEU:HD21	1:A:74:GLY:HA2	1.89	0.54
1:B:22:PHE:CD2	1:B:39:VAL:HG21	2.42	0.54
1:A:527:ASP:OD2	1:A:528:THR:HG23	2.08	0.54
1:A:84:MET:HE1	2:A:951:HOH:O	2.08	0.53
1:A:180:ARG:NE	2:A:874:HOH:O	2.42	0.53
1:B:409:PRO:HB3	1:B:414:GLU:HB3	1.90	0.53
1:A:342:VAL:HG12	1:A:373:LEU:HD23	1.91	0.53
1:B:138:ALA:HB1	1:B:170:PHE:HB2	1.91	0.52
1:A:367:PRO:HG3	1:A:483:ALA:HA	1.89	0.52
1:A:306:LYS:HD2	1:A:307:PHE:N	2.24	0.52
1:A:259:LEU:CD1	1:A:260:ASN:ND2	2.73	0.52
1:A:180:ARG:HH11	1:A:180:ARG:HG3	1.76	0.51
1:B:24:TYR:CD2	1:B:90:GLU:HG2	2.45	0.51
1:B:418:ILE:O	1:B:422:GLN:HG3	2.11	0.51
1:B:115:VAL:HG22	1:B:119:LYS:HE3	1.92	0.50
1:B:218:ASP:OD2	1:B:284:LYS:NZ	2.44	0.50
1:A:11:LEU:HD11	1:A:101:PRO:CB	2.42	0.50
1:A:415:ALA:O	1:A:418:ILE:HG22	2.10	0.50
1:A:306:LYS:HD2	1:A:307:PHE:H	1.77	0.49
1:A:337:LEU:HA	1:A:381:ASP:OD1	2.12	0.49
1:A:338:ASN:HD21	1:A:381:ASP:N	2.03	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:529:GLN:O	1:A:532:VAL:HG12	2.11	0.49
1:A:306:LYS:HE3	1:A:307:PHE:O	2.13	0.48
1:B:31:MET:HB2	2:B:823:HOH:O	2.13	0.48
1:A:480:THR:HG22	1:A:487:LEU:HD13	1.95	0.48
1:A:256:PRO:HA	2:A:803:HOH:O	2.13	0.48
1:B:410:ASN:HD22	1:B:411:ASN:N	2.11	0.48
1:B:331:ASP:CG	1:B:334:LYS:HD3	2.33	0.47
1:A:11:LEU:HB3	1:A:12:PRO:HD3	1.97	0.47
1:B:274:LEU:CD1	1:B:280:PRO:HB3	2.44	0.47
1:A:86:GLU:OE1	1:B:375:HIS:HE1	1.97	0.47
1:A:275:PRO:HG2	1:A:278:VAL:CG2	2.45	0.47
1:A:4:GLU:HG3	1:A:115:VAL:HG21	1.96	0.47
1:A:32:LYS:HB3	1:A:34:ASP:OD1	2.14	0.47
1:A:115:VAL:O	1:A:119:LYS:HG3	2.15	0.47
1:B:63:ILE:O	1:B:64:THR:HB	2.14	0.47
1:A:76:LYS:NZ	2:A:683:HOH:O	2.47	0.46
1:A:338:ASN:ND2	1:A:338:ASN:C	2.61	0.46
1:B:22:PHE:CE2	1:B:39:VAL:HG21	2.50	0.46
1:A:38:TYR:O	1:A:41:PHE:HB3	2.16	0.45
1:A:505:PRO:O	1:A:506:ASN:HB2	2.16	0.45
1:A:416:LYS:O	1:A:420:VAL:HG23	2.16	0.45
1:A:43:VAL:O	1:A:46:ILE:HG22	2.16	0.45
1:B:410:ASN:C	1:B:410:ASN:HD22	2.19	0.45
1:B:276:LYS:O	1:B:277:ASN:HB2	2.17	0.45
1:A:287:ALA:HB1	1:A:296:LEU:HD11	1.98	0.45
1:A:136:VAL:HG12	1:A:165:GLN:HB3	1.98	0.45
1:B:350:SER:OG	1:B:463:LEU:HG	2.18	0.44
1:B:171:THR:HG23	1:B:231:LYS:HG3	1.99	0.44
1:B:43:VAL:O	1:B:46:ILE:HG22	2.17	0.44
1:B:366:THR:HB	1:B:367:PRO:HD2	1.99	0.44
1:A:3:ILE:HG23	1:A:4:GLU:N	2.33	0.44
1:B:364:ARG:HG3	1:B:364:ARG:HH11	1.82	0.43
1:A:224:ALA:HB2	1:A:264:GLY:HA2	1.99	0.43
1:B:255:GLN:HA	1:B:256:PRO:HD2	1.75	0.43
1:B:324:GLN:NE2	1:B:359:LEU:HA	2.33	0.43
1:B:348:ASN:ND2	1:B:350:SER:H	2.16	0.43
2:A:627:HOH:O	1:B:55:GLN:HG2	2.19	0.43
1:A:301:GLN:NE2	2:A:832:HOH:O	2.47	0.42
1:A:432:VAL:HG22	1:A:433:GLY:N	2.34	0.42
1:B:224:ALA:HB2	1:B:264:GLY:HA2	2.00	0.42
1:B:64:THR:OG1	1:B:292:ASP:OD1	2.32	0.42

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:172:THR:HB	1:B:173:PRO:HD3	2.02	0.42
1:A:275:PRO:HG2	1:A:278:VAL:HG23	2.02	0.42
1:A:76:LYS:HB3	1:A:76:LYS:NZ	2.35	0.42
1:B:509:SER:HA	1:B:524:ALA:HB1	2.02	0.42
1:B:508:ARG:O	1:B:525:GLU:O	2.38	0.42
1:A:346:TYR:OH	1:A:512:ARG:HD3	2.20	0.42
2:A:626:HOH:O	1:B:57:LYS:HE2	2.19	0.42
1:B:241:HIS:CE1	1:B:242:LYS:HG3	2.54	0.41
1:B:38:TYR:O	1:B:41:PHE:HB3	2.20	0.41
1:B:295:ARG:HG2	1:B:391:HIS:HB2	2.02	0.41
1:A:31:MET:CG	1:A:36:LEU:HD13	2.50	0.41
1:A:368:THR:HG22	1:A:369:GLY:N	2.35	0.41
1:A:465:ASN:HA	1:A:532:VAL:HG21	2.02	0.41
1:A:408:LYS:HA	1:A:409:PRO:HD2	1.96	0.41
1:B:30:ARG:O	1:B:31:MET:HB3	2.21	0.41
1:B:52:LYS:HG3	1:B:133:ALA:HB2	2.03	0.41
1:B:52:LYS:HD3	1:B:129:LEU:HA	2.03	0.41
1:A:61:VAL:HG11	1:A:152:THR:HG21	2.02	0.41
1:A:136:VAL:HG11	1:A:182:LEU:CD1	2.51	0.41
1:A:259:LEU:HD13	1:A:260:ASN:ND2	2.36	0.41
1:B:279:LYS:HA	1:B:280:PRO:HD2	1.87	0.41
1:B:331:ASP:HA	1:B:332:PRO:HD2	1.97	0.41
1:B:364:ARG:HG3	1:B:376:GLU:OE1	2.20	0.40
1:A:463:LEU:O	1:A:464:PRO:C	2.58	0.40
1:B:3:ILE:HG23	1:B:4:GLU:N	2.36	0.40
1:A:356:GLU:O	1:A:360:LYS:HA	2.21	0.40
1:A:521:ARG:HH11	1:A:521:ARG:HG2	1.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	503/544 (92%)	487 (97%)	13 (3%)	3 (1%)	25	13
1	B	451/544 (83%)	432 (96%)	18 (4%)	1 (0%)	47	39
All	All	954/1088 (88%)	919 (96%)	31 (3%)	4 (0%)	34	24

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	20	VAL
1	A	19	GLY
1	A	31	MET
1	B	19	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	442/480 (92%)	438 (99%)	4 (1%)	78	75
1	B	396/480 (82%)	394 (100%)	2 (0%)	88	88
All	All	838/960 (87%)	832 (99%)	6 (1%)	84	81

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	143	GLU
1	A	170	PHE
1	A	338	ASN
1	A	495	ASP
1	B	170	PHE
1	B	410	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (26) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	98	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	120	ASN
1	A	277	ASN
1	A	282	ASN
1	A	301	GLN
1	A	324	GLN
1	A	325	GLN
1	A	338	ASN
1	A	343	GLN
1	A	482	ASN
1	A	494	GLN
1	A	506	ASN
1	A	529	GLN
1	A	531	ASN
1	B	5	GLN
1	B	204	GLN
1	B	255	GLN
1	B	271	ASN
1	B	277	ASN
1	B	301	GLN
1	B	325	GLN
1	B	343	GLN
1	B	348	ASN
1	B	375	HIS
1	B	410	ASN
1	B	413	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	519/544 (95%)	0.17	31 (5%) 21 28	12, 25, 50, 60	0
1	B	461/544 (84%)	0.11	19 (4%) 37 44	12, 23, 47, 60	0
All	All	980/1088 (90%)	0.14	50 (5%) 28 35	12, 24, 50, 60	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	463	LEU	6.6
1	B	346	TYR	6.2
1	B	509	SER	6.0
1	A	488	VAL	5.5
1	A	478	PHE	5.0
1	A	480	THR	4.8
1	B	526	ALA	4.6
1	A	490	PRO	4.4
1	B	368	THR	4.4
1	A	487	LEU	4.3
1	B	527	ASP	3.9
1	A	479	LYS	3.7
1	A	103	PRO	3.6
1	B	367	PRO	3.6
1	A	1	MET	3.5
1	B	214	ASN	3.4
1	A	407	TYR	3.4
1	B	103	PRO	3.2
1	A	110	GLU	3.2
1	B	506	ASN	3.1
1	B	102	SER	3.1
1	A	481	THR	3.0
1	A	330	ILE	3.0
1	A	5	GLN	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	2.9
1	A	492	GLY	2.9
1	B	507	GLY	2.8
1	B	372	HIS	2.7
1	B	464	PRO	2.7
1	A	424	PHE	2.6
1	A	514	SER	2.5
1	B	5	GLN	2.5
1	B	524	ALA	2.5
1	A	17	PRO	2.5
1	A	257	ASN	2.5
1	A	19	GLY	2.4
1	A	102	SER	2.4
1	A	519	ALA	2.3
1	A	410	ASN	2.3
1	A	333	THR	2.3
1	A	69	PRO	2.2
1	B	334	LYS	2.2
1	A	528	THR	2.2
1	A	483	ALA	2.2
1	B	333	THR	2.1
1	A	428	ILE	2.1
1	A	26	THR	2.1
1	A	386	PHE	2.1
1	A	3	ILE	2.0
1	A	20	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.