

# Full wwPDB X-ray Structure Validation Report (i)

Aug 20, 2020 – 05:17 PM BST

PDB ID 6DN8

> Title SPRY domain-containing SOCS box protein complexed 2

> > (GZJ)VDINNN(CY3) Cyclic peptide inhibitor

: Law, R.H.P.; Caradoc-Davies, T.T.; Norton, R.S. Authors

Deposited on 2018-06-06

1.75 Å(reported) Resolution

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity 4.02b-467

> 1.8.5 (274361), CSD as541be (2020) Mogul

Xtriage (Phenix) 1.13 EDS 2.13.1

Percentile statistics 20191225.v01 (using entries in the PDB archive December 25th 2019)

> Refmac 5.8.0158

7.0.044 (Gargrove) CCP4

Ideal geometry (proteins) Engh & Huber (2001) Ideal geometry (DNA, RNA) Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) 2.13.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	229	81%	15%
1	С	229	82%	14%
1	Е	229	79% 7%	14%
2	В	8	13% 75% 13%	13%
2	D	8	88%	13%
2	F	8	75%	25%



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5289 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called SPRY domain-containing SOCS box protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	195	Total	С	N	О	S	0	1	0
1	Λ	190	1520	963	283	270	4	U	1	0
1	С	198	Total	С	N	О	S	0	0	0
1			1528	972	278	274	4	0	0	
1	T.	196	Total	С	N	О	S	0	0	0
1		196	1514	964	279	267	4	0	0	U

There are 69 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	MET	-	initiating methionine	UNP Q96A44
A	6	HIS	-	expression tag	UNP Q96A44
A	7	HIS	_	expression tag	UNP Q96A44
A	8	HIS	-	expression tag	UNP Q96A44
A	9	HIS	-	expression tag	UNP Q96A44
A	10	HIS	_	expression tag	UNP Q96A44
A	11	HIS	-	expression tag	UNP Q96A44
A	12	SER	-	expression tag	UNP Q96A44
A	13	SER	_	expression tag	UNP Q96A44
A	14	GLY	-	expression tag	UNP Q96A44
A	15	VAL	_	expression tag	UNP Q96A44
A	16	ASP	-	expression tag	UNP Q96A44
A	17	LEU	-	expression tag	UNP Q96A44
A	18	GLY	_	expression tag	UNP Q96A44
A	19	THR	-	expression tag	UNP Q96A44
A	20	GLU	_	expression tag	UNP Q96A44
A	21	ASN	-	expression tag	UNP Q96A44
A	22	LEU	_	expression tag	UNP Q96A44
A	23	TYR	=	expression tag	UNP Q96A44
A	24	PHE	-	expression tag	UNP Q96A44
A	25	GLN	-	expression tag	UNP Q96A44
A	26	SER	-	expression tag	UNP Q96A44
A	27	MET	_	expression tag	UNP Q96A44

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	5	MET	-	initiating methionine	UNP Q96A44
С	6	HIS	-	expression tag	UNP Q96A44
С	7	HIS	-	expression tag	UNP Q96A44
С	8	HIS	-	expression tag	UNP Q96A44
С	9	HIS	-	expression tag	UNP Q96A44
С	10	HIS	-	expression tag	UNP Q96A44
С	11	HIS	-	expression tag	UNP Q96A44
С	12	SER	_	expression tag	UNP Q96A44
С	13	SER	-	expression tag	UNP Q96A44
С	14	GLY	-	expression tag	UNP Q96A44
С	15	VAL	-	expression tag	UNP Q96A44
С	16	ASP	_	expression tag	UNP Q96A44
С	17	LEU	-	expression tag	UNP Q96A44
С	18	GLY	-	expression tag	UNP Q96A44
С	19	THR	-	expression tag	UNP Q96A44
С	20	GLU	-	expression tag	UNP Q96A44
С	21	ASN	_	expression tag	UNP Q96A44
С	22	LEU	_	expression tag	UNP Q96A44
С	23	TYR	-	expression tag	UNP Q96A44
С	24	PHE	-	expression tag	UNP Q96A44
С	25	GLN	-	expression tag	UNP Q96A44
С	26	SER	-	expression tag	UNP Q96A44
С	27	MET	-	expression tag	UNP Q96A44
Е	5	MET	-	initiating methionine	UNP Q96A44
Е	6	HIS	_	expression tag	UNP Q96A44
Е	7	HIS	-	expression tag	UNP Q96A44
Е	8	HIS	_	expression tag	UNP Q96A44
Е	9	HIS	-	expression tag	UNP Q96A44
Е	10	HIS	_	expression tag	UNP Q96A44
Е	11	HIS	_	expression tag	UNP Q96A44
Е	12	SER	=	expression tag	UNP Q96A44
Е	13	SER	_	expression tag	UNP Q96A44
Е	14	GLY	-	expression tag	UNP Q96A44
Е	15	VAL		expression tag	UNP Q96A44
Е	16	ASP		expression tag	UNP Q96A44
Е	17	LEU	=	expression tag	UNP Q96A44
Е	18	GLY	-	expression tag	UNP Q96A44
Е	19	THR	-	expression tag	UNP Q96A44
Е	20	GLU		expression tag	UNP Q96A44
Е	21	ASN	-	expression tag	UNP Q96A44
Е	22	LEU	-	expression tag	UNP Q96A44
Е	23	TYR	-	expression tag	UNP Q96A44

Continued on next page...




~ 1	e		
Continued	trom	nremous	naae
	110110	production	paycon

Chain	Residue	Modelled	Actual	Comment	Reference
Е	24	PHE	-	expression tag	UNP Q96A44
Е	25	GLN	-	expression tag	UNP Q96A44
Е	26	SER	-	expression tag	UNP Q96A44
Е	27	MET	ı	expression tag	UNP Q96A44

 $\bullet$  Molecule 2 is a protein called (GZJ) VDINNN(CY3) Cyclic peptide inhibitor.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	В	7	Total C N O 56 33 10 13	0	0	0
2	D	7	Total C N O 56 33 10 13	0	0	0
2	F	6	Total C N O 47 27 9 11	0	0	0


 $\bullet$  Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $\mathrm{C_2H_3O_2}).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0
3	Е	1	Total C O 4 2 2	0	0


 $\bullet$  Molecule 4 is SULFATE ION (three-letter code: SO4) (formula:  $\mathrm{O_4S}).$ 





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	С	1	Total O S 5 4 1	0	0
4	С	1	Total O S 5 4 1	0	0

• Molecule 5 is 2-AMINO-3-MERCAPTO-PROPIONAMIDE (three-letter code: CY3) (formula:  $C_3H_8N_2OS$ ).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
5	B	1	Total	С	N	О	S	0	0
'	В	1	7	3	2	1	1	0	0
5	D	1	Total	С	N	О	S	0	0
)	D	1	7	3	2	1	1	0	U
5	E.	1	Total	С	N	О	S	0	0
3	I,	1	7	3	2	1	1	0	U

#### • Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	167	Total O 168 168	0	1
6	В	6	Total O 6 6	0	0
6	С	185	Total O 185 185	0	0
6	D	9	Total O 9 9	0	0
6	E	142	Total O 142 142	0	0
6	F	10	Total O 10 10	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SPRY domain-containing SOCS box protein 4



• Molecule 2: (GZJ)VDINNN(CY3) Cyclic peptide inhibitor



$\alpha$ : $\mathbf{D}$		de la companya de la
Chain D:	88%	13%
CIICIII D .	5570	1370



• Molecule 2: (GZJ)VDINNN(CY3) Cyclic peptide inhibitor

Chain F: 75% 25%





# 4 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 2 21 21	Depositor	
Cell constants	49.02Å 109.37Å 118.10Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	45.27 - 1.75	Depositor	
Resolution (A)	40.12 - 1.75	EDS	
% Data completeness	99.8 (45.27-1.75)	Depositor	
(in resolution range)	99.9 (40.12-1.75)	EDS	
$R_{merge}$	0.08	Depositor	
$R_{sym}$	(Not available)	Depositor	
$< I/\sigma(I) > 1$	2.36 (at 1.75Å)	Xtriage	
Refinement program	BUSTER 2.10.3	Depositor	
P. P.	0.193 , 0.210	Depositor	
$R, R_{free}$	0.201 , $0.221$	DCC	
$R_{free}$ test set	3178 reflections $(4.91%)$	wwPDB-VP	
Wilson B-factor (Å <sup>2</sup> )	21.6	Xtriage	
Anisotropy	0.422	Xtriage	
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.32, 47.3	EDS	
L-test for twinning <sup>2</sup>	$ < L > = 0.45, < L^2> = 0.28$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
$F_o, F_c$ correlation	0.95	EDS	
Total number of atoms	5289	wwPDB-VP	
Average B, all atoms $(Å^2)$	28.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



 $<sup>^{1}</sup>$ Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CY3, GZJ, SO4, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z  > 5
1	A	0.52	0/1561	0.67	0/2125
1	С	0.51	0/1571	0.66	0/2142
1	E	0.49	0/1556	0.66	0/2119
2	В	0.39	0/46	0.57	0/62
2	D	0.58	0/46	0.52	0/62
2	F	0.59	0/46	0.63	0/62
All	All	0.51	0/4826	0.66	0/6572

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	В	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	В	1	GZJ	Mainchain

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1520	0	1479	5	0
1	С	1528	0	1480	6	0
1	Ε	1514	0	1473	10	0
2	В	56	0	42	0	0
2	D	56	0	42	0	0
2	F	47	0	41	0	0
3	A	4	0	3	0	0
3	С	4	0	3	0	0
3	Ε	4	0	3	0	0
4	A	5	0	0	0	0
4	С	10	0	0	0	0
5	В	7	0	5	0	0
5	D	7	0	5	0	0
5	F	7	0	6	0	0
6	A	168	0	0	0	0
6	В	6	0	0	0	0
6	С	185	0	0	0	0
6	D	9	0	0	0	0
6	Ε	142	0	0	0	0
6	F	10	0	0	0	0
All	All	5289	0	4582	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{ll}  ext{Interatomic} \  ext{distance } ( ext{Å}) \end{array}$	Clash overlap (Å)
1:E:37:LEU:HD21	1:E:191:ILE:HD11	1.66	0.75
1:C:191:ILE:HD13	1:E:63:LEU:HD11	1.69	0.73
1:E:84:THR:HB	1:E:126:SER:HB3	1.74	0.70
1:C:41:LEU:HD13	1:E:63:LEU:HD13	1.82	0.61
1:C:178:LEU:HB2	1:C:191:ILE:HB	1.85	0.57
1:A:126:SER:HB3	1:A:132:LEU:HD13	1.87	0.55
1:E:126:SER:HB2	1:E:132:LEU:HD13	1.90	0.54
1:C:164:PHE:CD1	1:E:121:ARG:HG2	2.43	0.53
1:C:40:LEU:HD22	1:C:97:LEU:HG	1.97	0.47
1:C:126:SER:HB3	1:C:132:LEU:HD13	1.98	0.46
1:E:178:LEU:HB2	1:E:191:ILE:HB	1.97	0.45
1:E:116:GLY:HA3	1:E:140:TRP:O	2.17	0.44

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${f distance}({f \AA})$	overlap (Å)
1:A:75:PHE:O	1:A:221:GLU:HA	2.19	0.43
1:A:37:LEU:HD21	1:A:191:ILE:HD11	2.01	0.42
1:E:46:ALA:HB1	1:E:50:VAL:HB	2.03	0.41
1:A:75:PHE:CD2	1:A:87:ILE:HG21	2.56	0.41
1:A:97:LEU:HD23	1:A:230:LEU:HD12	2.03	0.40
1:E:75:PHE:CE2	1:E:222:VAL:HB	2.56	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$_{ m ntiles}$
1	A	$192/229 \ (84\%)$	188 (98%)	4 (2%)	0	100	100
1	С	$196/229 \; (86\%)$	195 (100%)	1 (0%)	0	100	100
1	E	$192/229 \ (84\%)$	188 (98%)	4 (2%)	0	100	100
2	В	5/8 (62%)	5 (100%)	0	0	100	100
2	D	5/8 (62%)	5 (100%)	0	0	100	100
2	F	4/8 (50%)	3 (75%)	1 (25%)	0	100	100
All	All	594/711 (84%)	584 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	154/182~(85%)	153 (99%)	1 (1%)	86 79
1	С	154/182~(85%)	152 (99%)	2 (1%)	69 54
1	E	152/182 (84%)	151 (99%)	1 (1%)	84 75
2	В	6/6 (100%)	6 (100%)	0	100 100
2	D	6/6 (100%)	6 (100%)	0	100 100
2	F	6/6 (100%)	6 (100%)	0	100 100
All	All	478/564 (85%)	474 (99%)	4 (1%)	81 72

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	146	ARG
1	С	121	ARG
1	С	127	VAL
1	E	177	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	78	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Pos	Link	$ \mathbf{B} $	ond leng	${ m gths}$	Bond angles		
			res		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GZJ	В	1	2,5	7,8,9	0.42	0	8,9,11	0.66	0



Mol	Type Cl	Chain	Res	Link	$\mathbf{B}_{0}$	ond leng	${ m gths}$	В	ond ang	gles
MIOI		Chain		LILK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GZJ	D	1	2,5	7,8,9	0.43	0	8,9,11	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GZJ	В	1	2,5	-	0/7/8/10	-
2	GZJ	D	1	2,5	-	0/7/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

# 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuna	Chain	Res	Link	В	Bond lengths			Bond angles		
MIOI	Type	Chain			Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2	
4	SO4	С	302	_	4,4,4	0.76	0	6,6,6	0.30	0	
5	CY3	В	101	2	6,6,6	0.25	0	6,7,7	0.65	0	
5	CY3	F	101	2	6,6,6	0.39	0	6,7,7	0.89	0	
5	CY3	D	101	2	6,6,6	0.37	0	6,7,7	0.27	0	



Mol	Т	Chain	Res	Link	В	ond len	${ m gths}$	Bond angles		
10101	Type	Chain		LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ACT	E	301	_	1,3,3	0.18	0	0,3,3	0.00	-
3	ACT	С	301	_	1,3,3	2.99	1 (100%)	0,3,3	0.00	-
3	ACT	A	301	-	1,3,3	2.21	1 (100%)	0,3,3	0.00	-
4	SO4	С	303	-	4,4,4	0.54	0	6,6,6	0.93	0
4	SO4	A	302	_	4,4,4	0.72	0	6,6,6	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CY3	В	101	2	-	0/6/6/6	-
5	CY3	F	101	2	-	1/6/6/6	_
5	CY3	D	101	2	-	0/6/6/6	_

#### All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$\operatorname{Ideal}( ext{\AA})$
3	С	301	ACT	СН3-С	2.99	1.52	1.48
3	A	301	ACT	СН3-С	2.21	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	F	101	CY3	N-CA-CB-SG

There are no ring outliers.

No monomer is involved in short contacts.

# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	195/229~(85%)	-0.01	4 (2%) 63 71	13, 21, 57, 118	0
1	С	$198/229 \ (86\%)$	-0.05	2 (1%) 82 87	14, 22, 46, 74	0
1	E	$196/229 \ (85\%)$	0.00	6 (3%) 49 55	15, 25, 56, 72	0
2	В	6/8 (75%)	0.58	1 (16%) 1 2	24, 28, 35, 51	0
2	D	6/8 (75%)	-0.23	0 100 100	19, 23, 28, 30	0
2	F	6/8 (75%)	0.02	0 100 100	18, 21, 26, 35	0
All	All	607/711 (85%)	-0.02	13 (2%) 63 71	13, 23, 56, 118	0

All (13) RSRZ outliers are listed below:

Mol	Chain	${f Res}$	$\mathbf{Type}$	RSRZ
1	A	171	PHE	3.7
1	A	165	LEU	3.6
2	В	2	VAL	3.5
1	A	164	PHE	3.5
1	Ε	35	ALA	2.7
1	Ε	230	LEU	2.6
1	Ε	163	ALA	2.6
1	С	167	PRO	2.6
1	A	163	ALA	2.5
1	Ε	40	LEU	2.2
1	С	232	PRO	2.1
1	Ε	232	PRO	2.0
1	Ε	37	LEU	2.0

# 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B-factors}({f \AA}^2)$	Q < 0.9
2	GZJ	В	1	9/10	0.86	0.22	52,54,58,60	0
2	GZJ	D	1	9/10	0.92	0.10	26,28,36,39	0

# 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

# 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B-factors}({f \AA}^2)$	Q < 0.9
3	ACT	E	301	4/4	0.83	0.14	52,52,53,54	0
3	ACT	A	301	4/4	0.85	0.20	29,38,41,43	0
3	ACT	С	301	4/4	0.87	0.13	39,40,41,43	0
5	CY3	В	101	7/7	0.87	0.16	52,54,59,61	0
5	CY3	F	101	7/7	0.88	0.14	30,31,35,37	0
4	SO4	A	302	5/5	0.94	0.14	49,49,50,50	0
4	SO4	С	302	5/5	0.95	0.18	44,47,48,50	0
4	SO4	С	303	5/5	0.96	0.10	36,40,41,42	5
5	CY3	D	101	7/7	0.97	0.10	22,25,28,28	0

# 6.5 Other polymers (i)

There are no such residues in this entry.

