Full wwPDB X-ray Structure Validation Report (i)

Aug 23, 2023 – 01:19 AM EDT

PDB ID	:	3DOB
Title	:	Peptide-binding domain of Heat shock 70 kDa protein F44E5.5 from C.elegans.
Authors	:	Osipiuk, J.; Hatzos, C.; Gu, M.; Zhang, R.; Voisine, C.; Morimoto, R.I.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on	:	2008-07-03
Resolution	:	2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	152	14% 88% 7% • 5%
1	B	152	9% 78% 18% • •

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2393 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heat shock 70 kDa protein F44E5.5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	145	1127	685	207	231	4	0	0	0
1	B	148	1150	700	213	233	4	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	393	SER	-	expression tag	UNP Q9XTL8
A	394	ASN	-	expression tag	UNP Q9XTL8
A	395	ALA	-	expression tag	UNP Q9XTL8
A	480	ASN	ASP	engineered mutation	UNP Q9XTL8
B	393	SER	-	expression tag	UNP Q9XTL8
B	394	ASN	-	expression tag	UNP Q9XTL8
B	395	ALA	-	expression tag	UNP Q9XTL8
B	480	ASN	ASP	engineered mutation	UNP Q9XTL8

- Molecule 2 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
2	A	1	4	2	1	1	0	0
			Total	C	O	S		
2	B	1	4	2	1	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	56	56	56	0	0
			Total	O		
3	B	52	52	52	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Heat shock 70 kDa protein F44E5.5

Chain A:



• Molecule 1: Heat shock 70 kDa protein F44E5.5

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α, β, γ	102.01Å 102.01Å 80.68Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.80 – 2.39 38.74 – 2.39	Depositor EDS
% Data completeness (in resolution range)	99.8 (38.80-2.39) 99.7 (38.74-2.39)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.61 (at 2.39Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.184 , 0.224 0.183 , 0.224	Depositor DCC
R_{free} test set	1992 reflections (10.25%)	wwPDB-VP
Wilson B-factor (Å^2)	48.5	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 48.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	0.023 for -h,-k,l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	2393	wwPDB-VP
Average B, all atoms (Å^2)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.87% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.77	1/1140 (0.1%)	0.81	1/1538 (0.1%)
1	B	0.69	0/1163	0.76	0/1567
All	All	0.73	1/2303 (0.0%)	0.78	1/3105 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	513	GLN	CD-OE1	6.04	1.37	1.24

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	482	ASP	CB-CG-OD1	6.19	123.87	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1127	0	1101	4	0
1	B	1150	0	1133	17	0
2	A	4	0	5	1	0
2	B	4	0	5	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	56	0	0	0	0
3	B	52	0	0	3	0
All	All	2393	0	2244	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:410:VAL:HG21	1:B:537:VAL:HG21	1.43	1.00
1:B:511:LEU:HD21	1:B:519:MET:HE1	1.52	0.90
1:B:511:LEU:HD21	1:B:519:MET:CE	2.07	0.85
1:B:540:ARG:HB2	3:B:38:HOH:O	1.82	0.78
1:B:540:ARG:HA	3:B:84:HOH:O	1.88	0.73
1:B:494:LYS:HD2	3:B:59:HOH:O	1.93	0.68
1:B:452:ARG:HH12	1:B:533:GLN:HE22	1.56	0.53
1:B:511:LEU:CD2	1:B:516:ILE:HG13	2.43	0.49
1:A:512:THR:HB	1:A:515:ASP:OD1	2.12	0.49
1:B:511:LEU:HD23	1:B:516:ILE:HG13	1.96	0.48
1:B:510:ARG:O	1:B:510:ARG:HD2	2.15	0.47
1:B:426:CYS:HB3	2:B:602:BME:H12	1.96	0.47
1:B:460:PHE:CD2	1:B:502:ILE:HG12	2.50	0.47
1:B:417:ARG:O	1:B:418:ASN:HB2	2.16	0.44
1:B:399:PRO:HB2	1:B:516:ILE:HD13	2.00	0.44
1:B:496:THR:HB	1:B:498:LYS:HE3	1.99	0.44
1:A:426:CYS:HB3	2:A:601:BME:H12	2.00	0.44
1:A:515:ASP:OD1	1:A:515:ASP:N	2.51	0.43
1:B:477:VAL:HA	1:B:490:SER:O	2.20	0.42
1:B:532:GLU:O	1:B:536:ARG:HG3	2.20	0.41
1:A:516:ILE:O	1:A:520:VAL:HG13	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	141/152 (93%)	139 (99%)	2 (1%)	0	100	100
1	B	144/152 (95%)	137 (95%)	7 (5%)	0	100	100
All	All	285/304 (94%)	276 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/129 (96%)	117 (94%)	7 (6%)	21	34
1	B	126/129 (98%)	117 (93%)	9 (7%)	14	23
All	All	250/258 (97%)	234 (94%)	16 (6%)	17	28

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	470	ARG
1	A	475	ILE
1	A	502	ILE
1	A	513	GLN
1	A	515	ASP
1	A	528	LYS
1	A	537	VAL
1	B	420	ARG
1	B	423	THR
1	B	440	SER
1	B	507	GLU
1	B	511	LEU
1	B	514	SER
1	B	518	ARG
1	B	534	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	540	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	BME	A	601	-	3,3,3	0.19	0	1,2,2	1.17	0
2	BME	B	602	-	3,3,3	0.35	0	1,2,2	1.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BME	A	601	-	-	0/1/1/1	-
2	BME	B	602	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	BME	1	0
2	B	602	BME	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	145/152 (95%)	0.71	22 (15%)	2	1	29, 39, 78, 85	0
1	B	148/152 (97%)	0.63	13 (8%)	10	9	29, 45, 63, 82	0
All	All	293/304 (96%)	0.67	35 (11%)	4	4	29, 43, 76, 85	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	507	GLU	5.8
1	B	509	GLY	5.7
1	A	514	SER	5.3
1	B	508	LYS	5.2
1	A	425	ALA	4.5
1	A	518	ARG	4.0
1	B	506	ASN	4.0
1	A	478	THR	3.7
1	A	426	CYS	3.6
1	B	425	ALA	3.6
1	B	534	ARG	3.5
1	A	414	LEU	3.3
1	B	510	ARG	3.3
1	B	535	GLU	3.3
1	B	426	CYS	3.2
1	A	512	THR	3.1
1	B	478	THR	3.0
1	A	424	LYS	3.0
1	A	526	PHE	3.0
1	A	477	VAL	2.9
1	A	489	VAL	2.8
1	B	433	ALA	2.8
1	A	521	HIS	2.7
1	B	505	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	517	ASP	2.6
1	A	507	GLU	2.4
1	A	427	LYS	2.3
1	A	428	THR	2.3
1	A	536	ARG	2.2
1	A	528	LYS	2.2
1	A	519	MET	2.2
1	A	490	SER	2.2
1	A	491	ALA	2.2
1	B	477	VAL	2.1
1	A	524	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	BME	B	602	4/4	0.96	0.12	45,46,46,47	0
2	BME	A	601	4/4	0.97	0.16	45,47,49,50	0

6.5 Other polymers (i)

There are no such residues in this entry.