



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 20, 2023 – 09:28 PM JST

PDB ID : 7DP3  
Title : Human MCM8 N-terminal domain  
Authors : Li, J.; Liu, L.; Liu, Y.  
Deposited on : 2020-12-17  
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	 2% 78% 6% 16%
1	B	331	 7% 74% 10% 16%

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4677 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA helicase MCM8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	278	2207	1402	374	416	15	0	2	0
1	B	279	2210	1400	376	419	15	0	1	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	46	MET	-	initiating methionine	UNP Q9UJA3
A	47	GLY	-	expression tag	UNP Q9UJA3
A	48	SER	-	expression tag	UNP Q9UJA3
A	49	SER	-	expression tag	UNP Q9UJA3
A	50	HIS	-	expression tag	UNP Q9UJA3
A	51	HIS	-	expression tag	UNP Q9UJA3
A	52	HIS	-	expression tag	UNP Q9UJA3
A	53	HIS	-	expression tag	UNP Q9UJA3
A	54	HIS	-	expression tag	UNP Q9UJA3
A	55	HIS	-	expression tag	UNP Q9UJA3
A	56	SER	-	expression tag	UNP Q9UJA3
A	57	GLN	-	expression tag	UNP Q9UJA3
A	58	GLY	-	expression tag	UNP Q9UJA3
A	59	SER	-	expression tag	UNP Q9UJA3
A	60	MET	-	expression tag	UNP Q9UJA3
B	46	MET	-	initiating methionine	UNP Q9UJA3
B	47	GLY	-	expression tag	UNP Q9UJA3
B	48	SER	-	expression tag	UNP Q9UJA3
B	49	SER	-	expression tag	UNP Q9UJA3
B	50	HIS	-	expression tag	UNP Q9UJA3
B	51	HIS	-	expression tag	UNP Q9UJA3
B	52	HIS	-	expression tag	UNP Q9UJA3
B	53	HIS	-	expression tag	UNP Q9UJA3
B	54	HIS	-	expression tag	UNP Q9UJA3
B	55	HIS	-	expression tag	UNP Q9UJA3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	56	SER	-	expression tag	UNP Q9UJA3
B	57	GLN	-	expression tag	UNP Q9UJA3
B	58	GLY	-	expression tag	UNP Q9UJA3
B	59	SER	-	expression tag	UNP Q9UJA3
B	60	MET	-	expression tag	UNP Q9UJA3

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	145	Total	O	0	0
			145	145		
3	B	113	Total	O	0	0
			113	113		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA helicase MCM8



- Molecule 1: DNA helicase MCM8



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.02Å 52.03Å 137.22Å 90.00° 91.86° 90.00°	Depositor
Resolution (Å)	48.65 – 2.55 48.65 – 2.55	Depositor EDS
% Data completeness (in resolution range)	87.7 (48.65-2.55) 87.3 (48.65-2.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.69 (at 2.54Å)	Xtrriage
Refinement program	PHENIX 1.17_3644	Depositor
R, $R_{free}$	0.214 , 0.272 0.214 , 0.272	Depositor DCC
$R_{free}$ test set	1883 reflections (9.74%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.5	Xtrriage
Anisotropy	0.128	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 38.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.094 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	4677	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	0/2252	0.41	0/3049
1	B	0.24	0/2253	0.45	0/3050
All	All	0.24	0/4505	0.43	0/6099

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2207	0	2223	12	0
1	B	2210	0	2218	19	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	145	0	0	1	0
3	B	113	0	0	1	0
All	All	4677	0	4441	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:269:CYS:HB3	1:B:270:ARG:HA	1.83	0.61
1:B:340:ALA:HB1	1:B:346:ASN:HA	1.82	0.61
1:B:303:GLU:HB3	1:B:306:ARG:HD3	1.86	0.57
1:B:150:GLU:OE2	1:B:336:LYS:NZ	2.37	0.52
1:A:297:MET:HG3	1:A:308:PRO:HB3	1.90	0.52
1:B:96:ILE:HD13	1:B:158:LEU:HD23	1.91	0.51
1:B:284:VAL:HG23	3:B:585:HOH:O	2.10	0.51
1:B:313[B]:CYS:SG	1:B:359:ALA:HB2	2.50	0.50
1:A:96:ILE:HD13	1:A:158:LEU:HD23	1.94	0.50
1:A:332:THR:OG1	1:A:361:SER:OG	2.29	0.49
1:B:261:PRO:N	1:B:262:THR:HA	2.27	0.49
1:A:149:PRO:HB3	1:A:218:TYR:HB2	1.94	0.48
1:B:85:VAL:HG11	1:B:266:VAL:HA	1.94	0.48
1:A:227:ARG:NH2	1:A:312:GLU:OE2	2.44	0.47
1:B:269:CYS:H	1:B:270:ARG:HB3	1.80	0.46
1:B:319:LEU:HD22	1:B:362:ILE:HG22	1.97	0.46
1:A:319:LEU:HD22	1:A:362:ILE:HG22	1.96	0.46
1:B:207:LEU:HD21	1:B:221:LEU:HD13	1.96	0.46
1:B:364:ASN:OD1	1:B:365:SER:N	2.49	0.45
1:B:263:LYS:HG2	1:B:265:PRO:HA	1.99	0.44
1:B:149:PRO:HB3	1:B:218:TYR:HB2	2.00	0.44
1:A:112:LYS:NZ	3:A:505:HOH:O	2.44	0.44
1:A:162:GLN:HA	1:A:165:THR:HG22	2.00	0.44
1:A:208:LYS:HE2	1:A:208:LYS:HB3	1.87	0.44
1:B:299:ASP:HA	1:B:300:ASP:C	2.37	0.44
1:A:240:PHE:HB3	1:A:274:PHE:HB3	2.01	0.42
1:B:84:GLU:OE1	1:B:84:GLU:N	2.44	0.42
1:A:227:ARG:NH1	1:A:292:LYS:HD2	2.35	0.41
1:A:122:LEU:HD23	1:A:197:ARG:HB2	2.03	0.41
1:B:247:GLU:OE1	1:B:266:VAL:HG21	2.21	0.41
1:B:116:GLU:OE2	1:B:168:LEU:HD11	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries



of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/331 (83%)	265 (97%)	9 (3%)	0	100	100
1	B	276/331 (83%)	257 (93%)	19 (7%)	0	100	100
All	All	550/662 (83%)	522 (95%)	28 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/293 (85%)	249 (100%)	0	100	100
1	B	249/293 (85%)	249 (100%)	0	100	100
All	All	498/586 (85%)	498 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	278/331 (83%)	-0.06	8 (2%) 51 59	8, 36, 101, 180	0
1	B	279/331 (84%)	0.26	23 (8%) 11 14	11, 44, 130, 190	0
All	All	557/662 (84%)	0.10	31 (5%) 24 29	8, 39, 122, 190	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	268	VAL	8.7
1	B	246	GLY	6.8
1	A	349	ASP	5.0
1	B	342	GLU	4.8
1	B	348	ASN	4.8
1	B	350	LYS	4.4
1	B	267	PRO	4.4
1	B	265	PRO	4.2
1	B	345	ARG	4.1
1	A	256	GLY	3.9
1	B	242	CYS	3.6
1	B	305	GLY	3.6
1	B	343	GLY	3.5
1	B	261	PRO	3.5
1	B	243	ALA	3.2
1	B	269	CYS	3.2
1	B	303	GLU	3.1
1	B	275	THR	3.1
1	A	175	LEU	3.1
1	B	245	CYS	3.0
1	A	299	ASP	3.0
1	B	260	LEU	3.0
1	B	241	LEU	2.9
1	B	299	ASP	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	348	ASN	2.6
1	A	177	ALA	2.5
1	B	247	GLU	2.4
1	A	305	GLY	2.1
1	B	298	SER	2.1
1	A	178	GLN	2.1
1	B	278	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	B	400	1/1	0.95	0.12	103,103,103,103	0
2	ZN	A	400	1/1	0.99	0.11	57,57,57,57	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.