

Full wwPDB X-ray Structure Validation Report (i)

Jul 17, 2023 – 04:13 PM EDT

PDB ID : 8DQ3

Title: X-ray crystal structure of Aggregatibacter actinomycetemcomitans diman-

ganese(II) class Id ribonucleotide reductase beta subunit

Authors: Rose, H.R.; Jung, J.J.; Boal, A.K.

Deposited on : 2022-07-18

Resolution : 1.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.34

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.34

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	348	83%		15%
1	В	348	82%		15%
1	С	348	81%	_	15%
1	D	348	81%	_	15%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 10571 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Ribonucleoside-diphosphate reductase.

Mol	Chain	Residues	${f Atoms}$			ZeroOcc	AltConf	Trace		
1	Λ	296	Total	С	N	О	S	0	3	0
1	A	290	2494	1624	402	460	8	0	3	
1	В	296	Total	С	N	О	S	0	2	0
1	Ъ	290	2486	1617	401	459	9	0	2	0
1	С	296	Total	С	N	О	S	0	2	0
1		290	2490	1621	402	459	8	0	<u> </u>	
1	D	297	Total	С	N	О	S	0	0	0
1	ע	291	2483	1614	401	460	8		U	

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	GLY	-	expression tag	UNP A0A142G3G1
A	-17	SER	-	expression tag	UNP A0A142G3G1
A	-16	SER	-	expression tag	UNP A0A142G3G1
A	-15	HIS	-	expression tag	UNP A0A142G3G1
A	-14	HIS	-	expression tag	UNP A0A142G3G1
A	-13	HIS	-	expression tag	UNP A0A142G3G1
A	-12	HIS	-	expression tag	UNP A0A142G3G1
A	-11	HIS	-	expression tag	UNP A0A142G3G1
A	-10	HIS	-	expression tag	UNP A0A142G3G1
A	-9	SER	-	expression tag	UNP A0A142G3G1
A	-8	SER	-	expression tag	UNP A0A142G3G1
A	-7	GLY	-	expression tag	UNP A0A142G3G1
A	-6	LEU	-	expression tag	UNP A0A142G3G1
A	-5	VAL	-	expression tag	UNP A0A142G3G1
A	-4	PRO	-	expression tag	UNP A0A142G3G1
A	-3	ARG	-	expression tag	UNP A0A142G3G1
A	-2	GLY	-	expression tag	UNP A0A142G3G1
A	-1	SER	-	expression tag	UNP A0A142G3G1
A	0	HIS	-	expression tag	UNP A0A142G3G1
В	-18	GLY	-	expression tag	UNP A0A142G3G1
В	-17	SER	-	expression tag	UNP A0A142G3G1



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
В	-16	SER	-	expression tag	UNP A0A142G3G1
В	-15	HIS	-	expression tag	UNP A0A142G3G1
В	-14	HIS	-	expression tag	UNP A0A142G3G1
В	-13	HIS	-	expression tag	UNP A0A142G3G1
В	-12	HIS	-	expression tag	UNP A0A142G3G1
В	-11	HIS	-	expression tag	UNP A0A142G3G1
В	-10	HIS	-	expression tag	UNP A0A142G3G1
В	-9	SER	-	expression tag	UNP A0A142G3G1
В	-8	SER	-	expression tag	UNP A0A142G3G1
В	-7	GLY	-	expression tag	UNP A0A142G3G1
В	-6	LEU	-	expression tag	UNP A0A142G3G1
В	-5	VAL	-	expression tag	UNP A0A142G3G1
В	-4	PRO	-	expression tag	UNP A0A142G3G1
В	-3	ARG	-	expression tag	UNP A0A142G3G1
В	-2	GLY	-	expression tag	UNP A0A142G3G1
В	-1	SER	-	expression tag	UNP A0A142G3G1
В	0	HIS	-	expression tag	UNP A0A142G3G1
С	-18	GLY	-	expression tag	UNP A0A142G3G1
С	-17	SER	-	expression tag	UNP A0A142G3G1
С	-16	SER	-	expression tag	UNP A0A142G3G1
С	-15	HIS	-	expression tag	UNP A0A142G3G1
С	-14	HIS	-	expression tag	UNP A0A142G3G1
С	-13	HIS	-	expression tag	UNP A0A142G3G1
С	-12	HIS	-	expression tag	UNP A0A142G3G1
С	-11	HIS	-	expression tag	UNP A0A142G3G1
С	-10	HIS	-	expression tag	UNP A0A142G3G1
С	-9	SER	-	expression tag	UNP A0A142G3G1
С	-8	SER	-	expression tag	UNP A0A142G3G1
С	-7	GLY	_	expression tag	UNP A0A142G3G1
С	-6	LEU	-	expression tag	UNP A0A142G3G1
С	-5	VAL	-	expression tag	UNP A0A142G3G1
С	-4	PRO	-	expression tag	UNP A0A142G3G1
С	-3	ARG	-	expression tag	UNP A0A142G3G1
С	-2	GLY	-	expression tag	UNP A0A142G3G1
С	-1	SER	-	expression tag	UNP A0A142G3G1
С	0	HIS	-	expression tag	UNP A0A142G3G1
D	-18	GLY	-	expression tag	UNP A0A142G3G1
D	-17	SER	-	expression tag	UNP A0A142G3G1
D	-16	SER	-	expression tag	UNP A0A142G3G1
D	-15	HIS	-	expression tag	UNP A0A142G3G1
D	-14	HIS	-	expression tag	UNP A0A142G3G1
D	-13	HIS	-	expression tag	tinued on next rage



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-12	HIS	-	expression tag	UNP A0A142G3G1
D	-11	HIS	-	expression tag	UNP A0A142G3G1
D	-10	HIS	-	expression tag	UNP A0A142G3G1
D	-9	SER	-	expression tag	UNP A0A142G3G1
D	-8	SER	-	expression tag	UNP A0A142G3G1
D	-7	GLY	-	expression tag	UNP A0A142G3G1
D	-6	LEU	-	expression tag	UNP A0A142G3G1
D	-5	VAL	-	expression tag	UNP A0A142G3G1
D	-4	PRO	-	expression tag	UNP A0A142G3G1
D	-3	ARG	-	expression tag	UNP A0A142G3G1
D	-2	GLY	-	expression tag	UNP A0A142G3G1
D	-1	SER	-	expression tag	UNP A0A142G3G1
D	0	HIS	-	expression tag	UNP A0A142G3G1

• Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Mn 2 2	0	0
2	В	2	Total Mn 2 2	0	0
2	С	2	Total Mn 2 2	0	0
2	D	2	Total Mn 2 2	0	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0
3	С	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0


 \bullet Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\mathrm{C_2H_6O_2}).$





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	С	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0

 \bullet Molecule 5 is FORMIC ACID (three-letter code: FMT) (formula: $\mathrm{CH_2O_2}).$



M	ol	Chain	Residues	Atoms		ZeroOcc	AltConf	
Ę	ó	A	1	Total 3	C 1	O 2	0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total C O 3 1 2	0	0
5	В	1	Total C O 3 1 2	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	152	Total O 152 152	0	0
6	В	171	Total O 171 171	0	0
6	С	134	Total O 134 134	0	0
6	D	128	Total O 128 128	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ribonucleoside-diphosphate reductase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	52.21Å 156.54Å 80.72Å	Depositor
a, b, c, α , β , γ	90.00° 91.09° 90.00°	Depositor
Resolution (Å)	44.22 - 1.67	Depositor
resolution (A)	44.22 - 1.67	EDS
% Data completeness	99.5 (44.22-1.67)	Depositor
(in resolution range)	99.5 (44.22-1.67)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.35 (at 1.67Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.170 , 0.195	Depositor
it, it free	0.170 , 0.196	DCC
R_{free} test set	7461 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	12.9	Xtriage
Anisotropy	0.004	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.39 , 46.5	EDS
L-test for twinning ²	$< L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	0.047 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10571	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, EDO, MG, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.36	0/2567	0.52	0/3463	
1	В	0.38	0/2555	0.53	0/3446	
1	С	0.34	0/2559	0.52	0/3451	
1	D	0.35	0/2545	0.51	0/3432	
All	All	0.36	0/10226	0.52	0/13792	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2494	0	2437	3	0
1	В	2486	0	2431	6	0
1	С	2490	0	2437	9	0
1	D	2483	0	2428	9	0
2	A	2	0	0	0	0
2	В	2	0	0	0	0
2	С	2	0	0	0	0
2	D	2	0	0	0	0
3	A	1	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
4	A	4	0	6	0	0
4	С	4	0	6	0	0
4	D	4	0	6	1	0
5	A	3	0	1	0	0
5	В	6	0	2	0	0
6	A	152	0	0	0	0
6	В	171	0	0	1	0
6	С	134	0	0	2	0
6	D	128	0	0	0	0
All	All	10571	0	9754	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:15:GLU:OE1	6:B:501:HOH:O	2.13	0.67
1:D:145:LYS:H	4:D:404:EDO:H11	1.61	0.66
1:B:64:LYS:NZ	1:B:250:GLU:HG2	2.14	0.61
1:C:183:LYS:NZ	6:C:502:HOH:O	2.29	0.58
1:B:230:LEU:HD11	1:B:277:LEU:HD13	1.92	0.51
1:C:107:LYS:HD2	1:D:16:ILE:HD13	1.93	0.51
1:B:64:LYS:HZ2	1:B:250:GLU:HG2	1.76	0.49
1:C:291:LYS:O	1:C:294:GLU:HG2	2.12	0.48
1:D:125:ASN:O	1:D:128:GLN:HG2	2.13	0.48
1:B:201:GLU:O	1:B:205:ARG:HG3	2.15	0.47
1:C:16:ILE:HD12	1:D:74:VAL:HG12	1.98	0.46
1:B:116:LYS:HD3	1:B:116:LYS:HA	1.76	0.46
1:C:22:PRO:HA	1:C:25:LEU:HD23	1.99	0.45
1:D:159:MET:SD	1:D:230:LEU:HG	2.56	0.45
1:A:21:TYR:OH	1:A:217:GLU:OE2	2.30	0.45
1:A:116:LYS:HD3	1:A:116:LYS:HA	1.73	0.44
1:C:166:HIS:CD2	1:C:204:GLY:HA3	2.52	0.44
1:A:230:LEU:HD11	1:A:277:LEU:HD13	2.01	0.43
1:C:16:ILE:HD13	1:D:107:LYS:HD2	2.00	0.43
1:D:171:SER:HB3	1:D:266:ILE:HG21	2.00	0.43
1:C:184:ASN:OD1	1:C:187:LYS:NZ	2.52	0.42



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:C:292:GLU:OE2	6:C:501:HOH:O	2.22	0.41
1:D:201:GLU:O	1:D:205:ARG:HG3	2.21	0.41
1:D:116:LYS:HA	1:D:116:LYS:HD3	1.79	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	297/348~(85%)	293 (99%)	4 (1%)	0	100	100
1	В	296/348~(85%)	291 (98%)	5 (2%)	0	100	100
1	С	296/348~(85%)	292 (99%)	4 (1%)	0	100	100
1	D	295/348~(85%)	290 (98%)	5 (2%)	0	100	100
All	All	1184/1392~(85%)	1166 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	$272/315\ (86\%)$	269 (99%)	3 (1%)	73 61	
1	В	$271/315\ (86\%)$	269 (99%)	2 (1%)	84 76	



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	С	271/315 (86%)	270 (100%)	1 (0%)	91	86	
1	D	270/315 (86%)	269 (100%)	1 (0%)	91	86	
All	All	1084/1260 (86%)	1077 (99%)	7 (1%)	86	79	

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	107	LYS
1	A	144	ARG
1	A	172	GLN
1	В	107	LYS
1	В	172	GLN
1	С	172	GLN
1	D	172	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 18 ligands modelled in this entry, 12 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain Res L		Link	В	ond leng	gths	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	FMT	В	405	-	2,2,2	0.68	0	1,1,1	0.23	0
4	EDO	D	404	-	3,3,3	0.48	0	2,2,2	0.44	0
4	EDO	A	404	-	3,3,3	0.60	0	2,2,2	0.12	0
4	EDO	С	404	-	3,3,3	0.49	0	2,2,2	0.44	0
5	FMT	A	405	-	2,2,2	0.68	0	1,1,1	0.24	0
5	FMT	В	404	-	2,2,2	0.69	0	1,1,1	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	D	404	-	-	1/1/1/1	-
4	EDO	С	404	-	-	0/1/1/1	-
4	EDO	A	404	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	404	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

\mathbf{Mol}	Chain	Res	Type	Clashes	Symm-Clashes
4	D	404	EDO	1	0

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	296/348~(85%)	-0.16	3 (1%) 82 85	6, 14, 25, 39	0
1	В	296/348~(85%)	-0.19	1 (0%) 94 94	6, 12, 24, 37	0
1	С	296/348~(85%)	-0.04	6 (2%) 65 69	7, 15, 27, 40	0
1	D	297/348 (85%)	-0.10	3 (1%) 82 85	7, 15, 28, 46	0
All	All	1185/1392 (85%)	-0.12	13 (1%) 80 83	6, 14, 26, 46	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	301	LEU	4.4
1	D	302	SER	4.2
1	A	302	SER	3.6
1	С	301	LEU	3.2
1	D	6	LYS	3.2
1	С	302	SER	3.2
1	С	163[A]	PHE	2.6
1	D	128	GLN	2.6
1	В	301	LEU	2.2
1	С	287	PRO	2.2
1	С	291	LYS	2.2
1	С	14	PHE	2.1
1	A	163[A]	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	MG	С	403	1/1	0.77	0.26	38,38,38,38	0
4	EDO	D	404	4/4	0.78	0.19	27,27,27,38	0
5	FMT	A	405	3/3	0.84	0.16	22,22,27,28	0
3	MG	A	403	1/1	0.87	0.14	27,27,27,27	0
5	FMT	В	405	3/3	0.88	0.13	27,27,32,33	0
3	MG	D	403	1/1	0.90	0.29	31,31,31,31	0
5	FMT	В	404	3/3	0.94	0.08	22,22,24,30	0
4	EDO	A	404	4/4	0.94	0.09	17,19,19,19	0
3	MG	В	403	1/1	0.95	0.15	29,29,29,29	0
4	EDO	С	404	4/4	0.97	0.13	18,20,22,22	0
2	MN	С	401	1/1	0.99	0.08	9,9,9,9	0
2	MN	A	401	1/1	1.00	0.08	7,7,7,7	0
2	MN	С	402	1/1	1.00	0.06	9,9,9,9	0
2	MN	D	401	1/1	1.00	0.06	7,7,7,7	0
2	MN	D	402	1/1	1.00	0.08	7,7,7,7	0
2	MN	A	402	1/1	1.00	0.07	8,8,8,8	0
2	MN	В	401	1/1	1.00	0.08	6,6,6,6	0
2	MN	В	402	1/1	1.00	0.06	6,6,6,6	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.










Electron density around MN A 401:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around MN C 402: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around MN D 401: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around MN D 402: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c \ (\mathrm{at}\ 0.7\ \mathrm{rmsd}) \ \mathrm{in}\ \mathrm{gray}$


 ${
m mF}_o{
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)
















6.5 Other polymers (i)

There are no such residues in this entry.

