

Full wwPDB X-ray Structure Validation Report (i)

May 26, 2020 – 04:37 pm BST

PDB ID : 1DTH

Title : METALLOPROTEASE

Authors : Botos, I.; Scapozza, L.; Zhang, D.; Liotta, L.A.; Meyer, E.F.

Deposited on : 1996-02-12

Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

Mol Probity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# { m Entries})$	$(\# ext{Entries}, ext{resolution range}(ext{Å}))$
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	A	203	78%	20%	•
1	В	203	79%	17%	•

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BAT	В	973	-	-	X	-



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3250 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ATROLYSIN C.

\mathbf{Mol}	Chain	Residues	${f Atoms}$			ZeroOcc	AltConf	Trace		
1	Δ	202	Total	С	N	О	S	0	0	0
1	Λ	202	1591	1014	263	303	11	U	0	
1	D	202	Total	С	N	О	S	0	0	0
1	Б	202	1591	1014	263	303	11	U	0	U

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0

• Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total Ca 1 1	0	0
3	A	1	Total Ca 1 1	0	0

• Molecule 4 is 4-(N-HYDROXYAMINO)-2R-ISOBUTYL-2S-(2-THIENYLTHIOMETHYL) SUCCINYL-L-PHENYLALANINE-N-METHYLAMIDE (three-letter code: BAT) (formula: $C_{23}H_{31}N_3O_4S_2$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf			
1	Λ.	1	Total	С	N	О	S	0	0	
4	A	1	32	23	3	4	2	0		
1	D	1	Total	С	N	О	S	0	0	
4	Б	1	32	23	3	4	2		U	




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: ATROLYSIN C





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 65	Depositor
Cell constants	97.21Å 97.21Å 87.91Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 - 2.00	Depositor
% Data completeness	92.6 (8.00-2.00)	Depositor
(in resolution range)	,	Берозног
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.168 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3250	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA, BAT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.44	0/1621	0.74	0/2184	
1	В	0.48	0/1621	0.73	0/2184	
All	All	0.46	0/3242	0.73	0/4368	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	$\mathbf{H}(\mathbf{model})$	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	1591	0	1530	37	0
1	В	1591	0	1530	38	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
4	A	32	0	29	18	0
4	В	32	0	31	22	0
All	All	3250	0	3120	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.



All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap(A)
1:B:59:TRP:CZ3	1:B:59:TRP:CZ2	2.42	1.03
1:A:59:TRP:CZ2	1:A:59:TRP:CZ3	2.43	1.01
1:A:81:TRP:CZ2	1:A:81:TRP:CZ3	2.42	1.01
1:B:81:TRP:CZ2	1:B:81:TRP:CZ3	2.42	1.00
1:A:176:TYR:CE2	4:A:972:BAT:H6	1.99	0.97
1:A:108:LEU:HD22	4:A:972:BAT:H121	1.63	0.80
1:A:176:TYR:HE2	4:A:972:BAT:H6	1.45	0.80
1:A:176:TYR:CD2	4:A:972:BAT:H6	2.18	0.79
1:A:167:ARG:HB2	4:A:972:BAT:S2	2.29	0.72
1:A:169:GLY:O	4:A:972:BAT:S2	2.49	0.71
1:A:110:LEU:HA	4:A:972:BAT:H161	1.72	0.70
1:B:108:LEU:HD22	4:B:973:BAT:H111	1.76	0.67
4:A:972:BAT:H112	4:A:972:BAT:C2	2.25	0.67
1:B:108:LEU:HB2	4:B:973:BAT:H111	1.76	0.66
1:B:110:LEU:HA	4:B:973:BAT:H161	1.80	0.63
1:B:170:LEU:HA	4:B:973:BAT:S1	2.38	0.63
1:A:75:LEU:HD23	1:A:125:ILE:HG13	1.81	0.62
1:A:108:LEU:HB2	4:A:972:BAT:H10	1.82	0.61
1:B:170:LEU:HA	4:B:973:BAT:S2	2.41	0.60
1:A:176:TYR:HE2	4:A:972:BAT:C6	2.16	0.57
4:A:972:BAT:O1	4:A:972:BAT:H112	2.04	0.57
1:A:17:ARG:HB3	1:A:100:ALA:HB3	1.88	0.56
1:A:75:LEU:HD22	1:A:123:ILE:HD12	1.87	0.56
1:B:108:LEU:CB	4:B:973:BAT:H111	2.36	0.55
1:B:170:LEU:CA	4:B:973:BAT:S1	2.95	0.55
1:A:169:GLY:HA2	4:A:972:BAT:H91	1.90	0.54
1:A:170:LEU:HA	4:A:972:BAT:S2	2.48	0.54
1:B:35:GLU:O	1:B:38:ASN:HB2	2.08	0.53
1:A:108:LEU:N	1:A:108:LEU:HD12	2.24	0.52
1:A:108:LEU:HD22	4:A:972:BAT:H111	1.91	0.52
4:A:972:BAT:H23	4:A:972:BAT:H92	1.92	0.52
1:B:108:LEU:HD13	4:B:973:BAT:H121	1.91	0.52
1:B:169:GLY:O	4:B:973:BAT:S2	2.67	0.51
1:A:105:GLU:O	1:A:106:GLU:HB2	2.11	0.51
1:A:172:LYS:HD3	1:A:172:LYS:H	1.75	0.51
1:A:194:LYS:N	1:A:195:PRO:HD3	2.27	0.50
1:B:170:LEU:CB	4:B:973:BAT:H113	2.42	0.50
1:B:196:GLN:NE2	1:B:196:GLN:H	2.10	0.49
1:B:169:GLY:HA2	4:B:973:BAT:H92	1.94	0.49
1:B:12:VAL:HG22	1:B:53:LEU:HD21	1.93	0.49

Continued on next page...



Continued from previous page...

Continuea from prev		Interatomic	Clash	
Atom-1	Atom-2	${\rm distance} \; (\mathring{\rm A})$	overlap (Å)	
1:B:108:LEU:HB2	4:B:973:BAT:C11	2.41	0.49	
1:A:116:MET:O	1:A:117:CYS:HB2	2.13	0.48	
1:B:108:LEU:CD2	4:B:973:BAT:H111	2.42	0.48	
1:B:151:GLU:HB2	1:B:185:TYR:CZ	2.49	0.48	
1:B:165:ILE:HA	4:B:973:BAT:H6	1.96	0.47	
1:B:170:LEU:HB2	4:B:973:BAT:H113	1.96	0.47	
1:B:170:LEU:N	4:B:973:BAT:H91	2.30	0.47	
1:A:113:LEU:O	1:A:115:THR:HG23	2.15	0.47	
1:B:75:LEU:HG	1:B:110:LEU:HD23	1.97	0.47	
1:A:108:LEU:HD23	1:A:129:HIS:CD2	2.50	0.47	
1:A:163:LEU:H	1:A:163:LEU:HD12	1.80	0.47	
1:B:186:TYR:CE1	1:B:190:LEU:HD13	2.50	0.46	
1:A:165:ILE:O	4:A:972:BAT:S1	2.74	0.45	
1:B:26:LEU:HG	1:B:58:ILE:HD13	1.97	0.45	
1:A:108:LEU:HD23	1:A:129:HIS:CG	2.52	0.45	
1:A:101:ILE:HG23	1:A:103:LEU:HD13	1.99	0.45	
1:A:134:LEU:O	1:A:138:VAL:HG13	2.16	0.45	
1:B:139:THR:HG23	4:B:973:BAT:O2	2.17	0.44	
1:A:45:ARG:HD3	1:B:48:ASN:ND2	2.32	0.44	
1:A:103:LEU:HB2	1:A:107:THR:O	2.18	0.44	
1:A:49:ILE:HD11	1:A:186:TYR:CZ	2.53	0.44	
4:A:972:BAT:C23	4:A:972:BAT:H92	2.47	0.44	
1:A:103:LEU:HB3	1:A:107:THR:HB	2.00	0.43	
1:B:142:HIS:HB3	4:B:973:BAT:HO2	1.84	0.43	
1:A:176:TYR:CE2	4:A:972:BAT:C6	2.86	0.43	
1:B:169:GLY:C	4:B:973:BAT:S1	2.96	0.43	
1:B:108:LEU:HD23	1:B:129:HIS:CG	2.54	0.43	
1:A:111:ALA:HA	1:A:112:PRO:HD3	1.86	0.42	
1:A:145:GLY:O	1:A:150:MET:HB2	2.19	0.42	
1:A:189:PHE:CE2	1:A:195:PRO:HG2	2.55	0.42	
4:B:973:BAT:O1	4:B:973:BAT:H112	2.20	0.42	
1:B:36:ILE:O	1:B:40:ILE:HG13	2.20	0.42	
1:B:17:ARG:HB3	1:B:100:ALA:HB3	2.03	0.41	
1:B:183:MET:HG2	1:B:187:GLU:OE1	2.21	0.41	
1:B:194:LYS:O	1:B:194:LYS:HD3	2.21	0.40	
1:B:169:GLY:O	4:B:973:BAT:S1	2.80	0.40	
1:B:112:PRO:HG3	1:B:121:LEU:HD12	2.02	0.40	
1:B:170:LEU:HB2	4:B:973:BAT:S1	2.61	0.40	
1:B:144:LEU:O	1:B:147:ASN:HB2	2.22	0.40	

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	200/203~(98%)	189 (94%)	10 (5%)	1 (0%)	29 23
1	В	200/203~(98%)	189 (94%)	10 (5%)	1 (0%)	29 23
All	All	400/406 (98%)	378 (94%)	20 (5%)	2 (0%)	29 23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	104	ASP
1	В	104	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	182/183 (100%)	173 (95%)	9 (5%)	25 21
1	В	182/183 (100%)	168 (92%)	14 (8%)	13 8
All	All	364/366 (100%)	341 (94%)	23 (6%)	18 13

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	LEU
1	A	60	SER
1	A	103	LEU
1	A	105	GLU

Continued on next page...



Continued from previous page...

			Das paye
Mol	Chain	Res	Type
1	A	121	LEU
1	A	138	VAL
1	A	172	LYS
1	A	180	ASP
1	A	197	CYS
1	В	26	LEU
1	В	27	ASN
1	В	48	ASN
1	В	53	LEU
1	В	62	GLU
1	В	71	SER
1	В	75	LEU
1	В	104	ASP
1	В	105	GLU
1	В	113	LEU
1	В	127	GLN
1	В	174	ARG
1	В	187	GLU
1	В	196	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	${ m Res}$	\mathbf{Type}
1	A	27	ASN
1	A	127	GLN
1	В	48	ASN
1	В	76	ASN
1	В	192	GLN
1	В	196	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Т	Chain	Dec	Link	Bo	nd leng	ths	В	ond ang	cles
MIOI	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	$\mid \# Z > 2$
4	BAT	A	972	2	31,33,33	2.61	5 (16%)	31,43,43	2.92	9 (29%)
4	BAT	В	973	2	31,33,33	1.43	4 (12%)	31,43,43	3.50	7 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BAT	A	972	2	-	10/35/37/37	0/2/2/2
4	BAT	В	973	2	-	9/35/37/37	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
4	A	972	BAT	O2-N1	-13.04	1.07	1.40
4	В	973	BAT	O2-N1	-4.24	1.29	1.40
4	В	973	BAT	C2-N1	4.10	1.39	1.33
4	A	972	BAT	C1-C2	3.48	1.57	1.51
4	В	973	BAT	C6-C5	2.24	1.47	1.39
4	A	972	BAT	C6-C5	2.23	1.47	1.39
4	A	972	BAT	O1-C2	-2.10	1.19	1.23
4	В	973	BAT	C6-C7	2.08	1.40	1.34
4	A	972	BAT	C6-C7	2.01	1.40	1.34

All (16) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
4	В	973	BAT	C1-C2-N1	-12.08	101.22	115.97
4	A	972	BAT	C9-C8-C1	11.95	129.00	112.29
4	В	973	BAT	C9-C8-C1	10.47	126.93	112.29
4	В	973	BAT	O1-C2-C1	7.19	130.39	121.73
4	A	972	BAT	C1-C2-N1	-4.29	110.73	115.97
4	В	973	BAT	C17-C14-C15	4.22	121.20	110.25
4	A	972	BAT	C8-C13-N2	-4.16	111.03	116.00
4	A	972	BAT	C17-C14-C15	4.13	120.96	110.25
4	В	973	BAT	C3-S1-C4	3.89	110.01	102.20
4	A	972	BAT	C12-C10-C9	-3.71	97.46	111.11
4	A	972	BAT	C11-C10-C9	3.62	124.42	111.11
4	В	973	BAT	O1-C2-N1	3.45	127.74	123.27
4	A	972	BAT	C16-N3-C15	-2.97	117.07	122.22
4	A	972	BAT	O1-C2-C1	2.91	125.23	121.73
4	В	973	BAT	C16-N3-C15	-2.30	118.23	122.22
4	A	972	BAT	O3-C13-C8	2.28	124.48	121.73

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	972	BAT	N2-C13-C8-C9
4	В	973	BAT	N2-C13-C8-C9
4	A	972	BAT	C17-C14-C15-O4
4	A	972	BAT	C17-C14-C15-N3
4	В	973	BAT	C17-C14-C15-O4
4	В	973	BAT	C17-C14-C15-N3
4	В	973	BAT	C3-C1-C8-C13
4	В	973	BAT	C11-C10-C9-C8
4	A	972	BAT	C3-C1-C2-O1
4	В	973	BAT	O3-C13-C8-C9
4	В	973	BAT	C3-C1-C8-C9
4	A	972	BAT	C8-C1-C2-O1
4	A	972	BAT	C3-C1-C2-N1
4	В	973	BAT	C2-C1-C8-C13
4	A	972	BAT	C2-C1-C3-S1
4	A	972	BAT	C8-C1-C2-N1
4	A	972	BAT	C3-C1-C8-C13
4	A	972	BAT	C2-C1-C8-C13
4	В	973	BAT	C17-C14-N2-C13


There are no ring outliers.

2 monomers are involved in 40 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	972	BAT	18	0
4	В	973	BAT	22	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

