



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2021 – 09:08 PM EDT

PDB ID : 3E2Q
Title : Crystal Structure Reduced PutA86-630 Mutant Y540S Complexed with trans-4-hydroxy-L-proline
Authors : Tanner, J.J.
Deposited on : 2008-08-06
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	551	 10% 78% 7% 15%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FAD	A	2001	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3930 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proline dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	468	3584	2268	632	667	17	0	4	0

There are 7 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	540	SER	TYR	engineered mutation	UNP P09546
A	631	HIS	-	expression tag	UNP P09546
A	632	HIS	-	expression tag	UNP P09546
A	633	HIS	-	expression tag	UNP P09546
A	634	HIS	-	expression tag	UNP P09546
A	635	HIS	-	expression tag	UNP P09546
A	636	HIS	-	expression tag	UNP P09546

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is 4-HYDROXYPROLINE (three-letter code: HYP) (formula: $C_5H_9NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	9	5	1	3	0	0

- Molecule 4 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			16	10	6		
4	A	1	Total	C	O	0	0
			16	10	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	252	Total	O	0	0
			252	252		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proline dehydrogenase



4 Data and refinement statistics i

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	73.08Å 141.65Å 145.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.43 – 1.75 40.43 – 1.74	Depositor EDS
% Data completeness (in resolution range)	98.8 (40.43-1.75) 98.9 (40.43-1.74)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 1.74Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.201 , 0.228 0.200 , 0.225	Depositor DCC
R_{free} test set	3814 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	25.5	Xtrriage
Anisotropy	0.709	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.003 for -h,-l,-k	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3930	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HYP, 1PE, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/3655	0.48	0/4962

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3584	0	3528	21	0
2	A	53	0	28	2	0
3	A	9	0	8	0	0
4	A	32	0	44	1	0
5	A	252	0	0	1	0
All	All	3930	0	3608	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:540:SER:HB2	2:A:2001:FAD:HM72	1.68	0.74
1:A:361:ARG:HG2	1:A:361:ARG:HH21	1.63	0.62
4:A:2003:1PE:H122	5:A:2250:HOH:O	2.05	0.56
1:A:116:LEU:HD23	1:A:120:VAL:HG12	1.88	0.54
1:A:565:SER:O	1:A:569:ARG:HG3	2.07	0.54
1:A:358:LEU:HD23	1:A:393:LEU:HD21	1.90	0.52
1:A:241:GLU:N	1:A:242:PRO:CD	2.73	0.51
1:A:402:VAL:HG22	1:A:431:ARG:HB2	1.94	0.49
1:A:106:ALA:O	1:A:110:LEU:HG	2.12	0.49
1:A:400:GLY:HA2	1:A:429:MET:O	2.14	0.48
1:A:503:TYR:OH	1:A:505:PRO:HA	2.14	0.47
1:A:256:GLY:HA3	1:A:549:LEU:HD21	1.96	0.46
1:A:294:ALA:HA	1:A:335:PRO:HG3	1.98	0.46
1:A:377:LEU:HD23	1:A:409:ARG:CZ	2.45	0.45
1:A:298:GLN:HE21	1:A:302:VAL:HG23	1.80	0.44
1:A:361:ARG:HG2	1:A:361:ARG:NH2	2.33	0.43
1:A:101:ARG:O	1:A:408:LYS:HD3	2.18	0.43
1:A:361:ARG:HG3	1:A:396:TRP:CG	2.52	0.43
2:A:2001:FAD:H1'2	2:A:2001:FAD:H9	1.15	0.43
1:A:586:THR:O	1:A:590:LYS:HG3	2.20	0.42
1:A:349:LEU:HD23	1:A:349:LEU:O	2.20	0.41
1:A:175:ILE:HA	1:A:176:PRO:HD3	1.87	0.41
1:A:368:ASN:HD21	1:A:402:VAL:HG23	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	468/551 (85%)	462 (99%)	6 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	354/450 (79%)	348 (98%)	6 (2%)	60 42

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99	TYR
1	A	101	ARG
1	A	168	LEU
1	A	241	GLU
1	A	254	LEU
1	A	406	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	113	GLN
1	A	271	ASN
1	A	298	GLN
1	A	305	GLN
1	A	316	ASN
1	A	368	ASN
1	A	497	GLN
1	A	534	ASN
1	A	546	HIS
1	A	560	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	1PE	A	2002	-	15,15,15	0.54	0	14,14,14	1.39	0
3	HYP	A	1	-	6,9,9	3.62	4 (66%)	5,12,12	2.22	1 (20%)
2	FAD	A	2001	-	51,58,58	3.08	20 (39%)	60,89,89	2.98	24 (40%)
4	1PE	A	2003	-	15,15,15	0.54	0	14,14,14	1.60	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	1PE	A	2002	-	-	2/13/13/13	-
3	HYP	A	1	-	-	0/0/13/13	0/1/1/1
2	FAD	A	2001	-	1/1/9/9	10/30/50/50	0/6/6/6
4	1PE	A	2003	-	-	4/13/13/13	-

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	FAD	O2'-C2'	-11.15	1.19	1.43
2	A	2001	FAD	C2B-C1B	-7.67	1.42	1.53
2	A	2001	FAD	C10-N1	6.23	1.41	1.33
3	A	1	HYP	CB-CG	-5.75	1.41	1.52
2	A	2001	FAD	C1'-N10	-5.54	1.42	1.48
2	A	2001	FAD	O4-C4	5.50	1.38	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	FAD	C9A-N10	-5.14	1.31	1.38
2	A	2001	FAD	C3B-C4B	-4.62	1.41	1.53
3	A	1	HYP	CD-CG	-4.58	1.42	1.53
3	A	1	HYP	CB-CA	-4.28	1.45	1.54
2	A	2001	FAD	C2B-C3B	-4.20	1.41	1.53
2	A	2001	FAD	PA-O1A	3.84	1.64	1.50
2	A	2001	FAD	C6A-N6A	3.65	1.47	1.34
2	A	2001	FAD	P-O1P	3.56	1.63	1.50
2	A	2001	FAD	C9A-C5X	-2.97	1.36	1.42
2	A	2001	FAD	C4-N3	2.85	1.38	1.33
2	A	2001	FAD	C8A-N7A	2.76	1.39	1.34
2	A	2001	FAD	C4X-C10	-2.64	1.36	1.38
2	A	2001	FAD	C4-C4X	-2.32	1.37	1.41
2	A	2001	FAD	C2-N1	-2.30	1.33	1.38
2	A	2001	FAD	C4X-N5	2.26	1.36	1.33
3	A	1	HYP	CA-N	-2.14	1.40	1.50
2	A	2001	FAD	C2A-N3A	2.12	1.35	1.32
2	A	2001	FAD	C4A-N3A	-2.03	1.32	1.35

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	FAD	C1'-N10-C9A	-11.14	109.53	118.29
2	A	2001	FAD	C1'-N10-C10	9.13	126.58	118.41
2	A	2001	FAD	O2'-C2'-C3'	6.80	125.63	109.10
2	A	2001	FAD	N3A-C2A-N1A	-6.55	118.45	128.68
2	A	2001	FAD	O4B-C4B-C5B	4.93	125.59	109.37
2	A	2001	FAD	O2'-C2'-C1'	4.83	121.23	109.59
3	A	1	HYP	CB-CG-CD	4.79	109.14	103.27
2	A	2001	FAD	C4-N3-C2	3.79	118.34	115.14
2	A	2001	FAD	O4'-C4'-C5'	3.68	118.18	109.92
2	A	2001	FAD	O3B-C3B-C4B	3.64	121.58	111.05
2	A	2001	FAD	C6-C5X-N5	3.49	122.90	119.05
2	A	2001	FAD	C4X-N5-C5X	3.47	120.23	116.77
2	A	2001	FAD	C5X-C9A-N10	3.43	120.20	117.72
2	A	2001	FAD	C4'-C3'-C2'	-3.37	106.36	113.36
2	A	2001	FAD	O3B-C3B-C2B	3.35	122.65	111.82
2	A	2001	FAD	C5B-C4B-C3B	3.12	126.89	115.18
2	A	2001	FAD	O2B-C2B-C1B	2.34	119.48	110.85
2	A	2001	FAD	C4-C4X-N5	2.26	121.18	118.60
2	A	2001	FAD	C5'-C4'-C3'	2.24	116.53	112.20
2	A	2001	FAD	C9A-C5X-N5	-2.13	119.03	122.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	FAD	P-O3P-PA	-2.11	125.58	132.83
2	A	2001	FAD	O4B-C1B-C2B	-2.09	103.87	106.93
2	A	2001	FAD	C10-C4X-N5	-2.06	119.83	121.26
2	A	2001	FAD	C9-C9A-C5X	2.03	123.35	119.88
2	A	2001	FAD	O2B-C2B-C3B	2.03	118.39	111.82
4	A	2003	1PE	OH4-C13-C23	2.01	119.46	110.39

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	2001	FAD	C4B

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2001	FAD	O4B-C4B-C5B-O5B
2	A	2001	FAD	C2'-C1'-N10-C9A
2	A	2001	FAD	C2'-C1'-N10-C10
2	A	2001	FAD	N10-C1'-C2'-O2'
2	A	2001	FAD	C1'-C2'-C3'-O3'
2	A	2001	FAD	C1'-C2'-C3'-C4'
2	A	2001	FAD	O3'-C3'-C4'-C5'
2	A	2001	FAD	C3B-C4B-C5B-O5B
4	A	2002	1PE	OH4-C13-C23-OH3
4	A	2002	1PE	OH5-C14-C24-OH4
2	A	2001	FAD	C2'-C3'-C4'-C5'
4	A	2003	1PE	C24-C14-OH5-C25
4	A	2003	1PE	C12-C22-OH3-C23
4	A	2003	1PE	OH2-C12-C22-OH3
4	A	2003	1PE	C13-C23-OH3-C22
2	A	2001	FAD	C3'-C4'-C5'-O5'


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2001	FAD	2	0
4	A	2003	1PE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	468/551 (84%)	0.59	57 (12%) 4 6	20, 35, 75, 173	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	152	LEU	12.4
1	A	159	SER	7.5
1	A	155	PHE	7.3
1	A	153	GLN	7.3
1	A	156	SER	6.8
1	A	151	LEU	6.5
1	A	243	LEU	5.8
1	A	145	ALA	5.6
1	A	184	LEU	5.1
1	A	164	ALA	4.8
1	A	142	SER	4.4
1	A	148	VAL	4.3
1	A	240	GLY	4.3
1	A	158	SER	4.3
1	A	149	GLN	4.1
1	A	242	PRO	3.9
1	A	160	GLN	3.9
1	A	244	ILE	3.9
1	A	450	LEU	3.8
1	A	163	VAL	3.8
1	A	342	TYR	3.7
1	A	143	GLY	3.6
1	A	138	GLN	3.6
1	A	403	ILE	3.5
1	A	141	ALA	3.5
1	A	430	ILE	3.3
1	A	609	ARG	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	157	LEU	3.3
1	A	165	LEU	3.3
1	A	168	LEU	3.1
1	A	432	LEU	3.1
1	A	183	ALA	3.1
1	A	90	VAL	3.1
1	A	150	GLY	2.9
1	A	147	MET	2.8
1	A	167	CYS	2.7
1	A	162	GLY	2.7
1	A	161	GLU	2.6
1	A	433	VAL	2.6
1	A	140	ASN	2.6
1	A	401	PHE	2.6
1	A	480[A]	ILE	2.5
1	A	371	ALA	2.5
1	A	577	LEU	2.5
1	A	169	ALA	2.4
1	A	271	ASN	2.4
1	A	402	VAL	2.4
1	A	369	ILE	2.4
1	A	88	GLN	2.3
1	A	166	MET	2.3
1	A	481	TYR	2.3
1	A	575	LEU	2.3
1	A	154	GLU	2.3
1	A	328	ILE	2.2
1	A	93	ALA	2.1
1	A	247	GLY	2.0
1	A	241	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	1PE	A	2002	16/16	0.77	0.14	48,57,60,60	0
4	1PE	A	2003	16/16	0.79	0.17	54,60,72,76	0
3	HYP	A	1	9/9	0.92	0.16	32,38,39,41	0
2	FAD	A	2001	53/53	0.94	0.12	17,23,28,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.