Full wwPDB X-ray Structure Validation Report (i)

Jun 12, 2024 – 01:57 PM EDT

PDB ID	:	1E54
Title	:	Anion-selective porin from Comamonas acidovorans
Authors	:	Zeth, K.; Diederichs, K.; Welte, W.; Engelhardt, H.
Deposited on	:	2000-07-17
Resolution	:	2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	332	79% 20% .
2	B	8	38% 50% 12%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 2630 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called OUTER MEMBRANE PORIN PROTEIN 32.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	332	2461	1546	430	483	2	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	PCA	GLN	modified residue	UNP P24305

- Molecule 2 is a protein called OMP32.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	8	65	37	12	16	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	1	1	1	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	97	Total O 97 97	0	0
5	B	1	Total O 1 1	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: OUTER MEMBRANE PORIN PROTEIN 32

Chain A: 79% 20%



• Molecule 2: OMP32

Chain B: 38% 50% 12%

4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α, β, γ	107.25Å 107.25Å 140.59Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.10	Depositor
% Data completeness (in resolution range)	89.4 (20.00-2.10)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
Refinement program	CNS 0.9	Depositor
R, R_{free}	0.200 , 0.235	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2630	wwPDB-VP
Average B, all atoms (Å^2)	34.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PCA, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.36	0/2498	0.65	0/3370
2	B	0.47	0/66	0.51	0/88
All	All	0.36	0/2564	0.65	0/3458

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2461	0	2396	56	0
2	B	65	0	48	4	0
3	A	1	0	0	0	0
4	A	5	0	0	0	0
5	A	97	0	0	3	0
5	B	1	0	0	0	0
All	All	2630	0	2444	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:305:LEU:H	1:A:321:GLN:HE22	1.07	0.97
1:A:101:LEU:HD13	1:A:207:THR:HB	1.54	0.89
1:A:305:LEU:H	1:A:321:GLN:NE2	1.78	0.80
1:A:106:LEU:HD11	1:A:220:SER:HB3	1.65	0.77
1:A:106:LEU:H	1:A:258:GLN:HE22	1.38	0.70
1:A:161:THR:HG22	1:A:166:ASP:HA	1.75	0.68
1:A:238:THR:HG22	5:A:2071:HOH:O	1.93	0.67
1:A:159:LYS:O	1:A:161:THR:HG23	1.96	0.66
1:A:121:TRP:CG	1:A:238:THR:HG21	2.30	0.66
1:A:305:LEU:N	1:A:321:GLN:HE22	1.88	0.66
1:A:80:SER:HB3	1:A:89:ARG:HG3	1.78	0.64
1:A:32:SER:HB3	1:A:36:THR:HA	1.80	0.63
1:A:18:ASN:CG	1:A:322:THR:HG22	2.21	0.60
1:A:315:VAL:HG22	1:A:316:GLN:H	1.70	0.56
1:A:19:LYS:HA	1:A:24:ASP:O	2.05	0.55
1:A:106:LEU:H	1:A:258:GLN:NE2	2.03	0.55
1:A:221:GLY:HA2	1:A:242:MET:HE2	1.88	0.55
1:A:2:SER:HA	1:A:44:THR:O	2.07	0.54
1:A:15:ALA:HB3	1:A:310:VAL:HG21	1.90	0.53
1:A:252:VAL:CG1	1:A:281:ASN:HB2	2.39	0.52
1:A:249:VAL:O	1:A:249:VAL:HG12	2.09	0.52
1:A:155:ALA:HB3	1:A:171:TYR:HB3	1.93	0.51
1:A:236:ILE:HD12	1:A:236:ILE:N	2.26	0.51
1:A:157:ASP:HB3	1:A:170:ARG:HB2	1.92	0.50
1:A:92:ARG:HH11	1:A:135:ASN:ND2	2.10	0.50
1:A:249:VAL:CG2	1:A:255:VAL:HG23	2.41	0.49
1:A:106:LEU:N	1:A:258:GLN:HE22	2.06	0.49
1:A:286:THR:OG1	1:A:330:HIS:HD2	1.95	0.49
1:A:251:GLY:C	1:A:252:VAL:HG23	2.33	0.49
1:A:270:LYS:O	1:A:298:LYS:HB2	2.12	0.49
1:A:315:VAL:HG22	1:A:316:GLN:N	2.28	0.49
1:A:94:LEU:HD23	1:A:99:GLN:HA	1.95	0.49
1:A:106:LEU:HD11	1:A:220:SER:CB	2.40	0.48
1:A:161:THR:HG22	1:A:166:ASP:CA	2.43	0.48
1:A:272:HIS:O	1:A:295:LEU:HD12	2.14	0.48
2:B:407:THR:OG1	2:B:408:SER:N	2.44	0.48
1:A:18:ASN:N	1:A:18:ASN:HD22	2.12	0.48
1:A:143:PRO:HG2	1:A:145:PHE:CZ	2.49	0.47
1:A:59:GLY:HA3	1:A:72:ASN:O	2.15	0.46
1:A:204:ASP:OD2	1:A:229:LYS:HE3	2.16	0.45
1:A:121:TRP:CD1	1:A:238:THR:HG21	2.52	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:LEU:HD11	1:A:133:ARG:NH2	2.32	0.45
1:A:189:LEU:HD23	1:A:206:ILE:HG13	1.99	0.45
1:A:252:VAL:HG12	1:A:281:ASN:HB2	1.98	0.44
2:B:401:ASP:N	2:B:401:ASP:OD1	2.51	0.44
1:A:101:LEU:CD1	1:A:207:THR:HB	2.39	0.44
1:A:284:LYS:HB2	1:A:284:LYS:NZ	2.33	0.43
1:A:159:LYS:HB3	1:A:159:LYS:HE2	1.80	0.42
1:A:332:PHE:OXT	2:B:405:ASN:HB2	2.18	0.42
1:A:111:GLY:O	1:A:114:PRO:HD3	2.20	0.42
1:A:189:LEU:CD2	1:A:206:ILE:HG13	2.50	0.42
1:A:18:ASN:ND2	1:A:322:THR:H	2.17	0.42
1:A:12:THR:HG22	5:A:2008:HOH:O	2.20	0.41
1:A:89:ARG:C	1:A:90:LEU:HD12	2.40	0.41
1:A:4:VAL:HB	2:B:403:TRP:CZ3	2.55	0.41
1:A:172:ILE:HG23	1:A:172:ILE:O	2.21	0.41
1:A:130:ASN:C	1:A:130:ASN:HD22	2.24	0.41
1:A:118:PHE:HB2	5:A:2036:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/332 (99%)	315 (96%)	15 (4%)	0	100	100
2	B	6/8 (75%)	5 (83%)	1 (17%)	0	100	100
All	All	336/340 (99%)	320 (95%)	16 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	244/244 (100%)	235 (96%)	9 (4%)	34 35
2	B	7/7 (100%)	6 (86%)	1 (14%)	3 1
All	All	251/251 (100%)	241 (96%)	10 (4%)	31 32

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	LEU
1	A	18	ASN
1	A	84	ASN
1	A	130	ASN
1	A	242	MET
1	A	252	VAL
1	A	257	LEU
1	A	284	LYS
1	A	305	LEU
2	B	401	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	ASN
1	A	130	ASN
1	A	135	ASN
1	A	160	GLN
1	A	180	ASN
1	A	224	GLN
1	A	258	GLN
1	A	264	GLN
1	A	281	ASN
1	A	291	ASN
1	A	297	ASN
1	A	316	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	321	GLN
1	A	330	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	PCA	A	1	1	7,8,9	4.31	1 (14%)	9,10,12	2.04	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	PCA	CA-N	-11.21	1.33	1.46

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	1	PCA	CB-CA-N	3.69	113.89	103.30
1	A	1	PCA	CB-CA-C	2.49	116.12	112.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	1	PCA	O-C-CA	-2.22	118.95	124.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	SO4	A	600	-	4,4,4	1.89	2 (50%)	6,6,6	0.89	0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	600	SO4	O1-S	3.13	1.62	1.46
4	A	600	SO4	O3-S	-2.09	1.30	1.47

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.