Full wwPDB X-ray Structure Validation Report (i)

Jan 2, 2024 – 06:45 pm GMT

PDB ID	:	5EBI
Title	:	Crystal structure of a DNA-RNA chimera in complex with Ba2+ ions: a case of unusual multi-domain twinning
Authors	:	Gilski, M.; Drozdzal, P.; Kierzek, R.; Jaskolski, M.
Deposited on	:	2015-10-19
Resolution	:	1.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1619 (1.14-1.06)
Clashscore	141614	1671 (1.14-1.06)
RSRZ outliers	127900	1588 (1.14-1.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	6	100%
1	B	6	100%
1	C	6	67% 33%
1	D	6	100%
1	E	6	100%
1	F	6	83% 17%
1	G	6	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	H	6	83% 17%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BA	A	105	-	-	X	-

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 1278 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is DNA/RNA hybrid called DNA/RNA (5'-D(*C)-R(P*G)-D(P*C)-R(P*G)-D(P*C)-R(P*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	6	130	59	24	41	6	0	2	0
1	B	6	123	57	24	37	5	0	0	0
1	C	6	130	59	24	41	6	0	2	0
1	D	6	123	57	24	37	5	0	0	0
1	E	6	123	57	24	37	5	0	0	0
1	F	6	144	63	24	49	8	0	5	0
1	G	6	123	57	24	37	5	0	0	0
1	H	6	137	61	24	45	7	0	4	0

- Molecule 2 is BARIUM ION (three-letter code: BA) (formula: Ba).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ba		
2	A	6	6	6	0	0
2	B	1	1	1	0	0
2	C	2	2	2	0	0
2	D	2	2	2	0	0
2	F	3	3	3	0	0
2	G	1	1	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ba		
2	H	1	1	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	31	31	31	0	0
			Total	O		
3	B	20	21	21	0	1
			Total	O		
3	C	30	30	30	0	0
			Total	O		
3	D	28	28	28	0	0
			Total	O		
3	E	36	38	38	0	2
			Total	O		
3	F	21	21	21	0	0
			Total	O		
3	G	35	36	36	0	1
			Total	O		
3	H	23	24	24	0	1

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA/RNA (5'-D(*C)-R(P*G)-D(P*C)-R(P*G)-D(P*C)-R(P*G)-3')

Chain A: 

There are no outlier residues recorded for this chain.

• Molecule 1: DNA/RNA (5'-D(*C)-R(P*G)-D(P*C)-R(P*G)-D(P*C)-R(P*G)-3')

Chain B: 

There are no outlier residues recorded for this chain.

• Molecule 1: DNA/RNA (5'-D(*C)-R(P*G)-D(P*C)-R(P*G)-D(P*C)-R(P*G)-3')

Chain C:

• Molecule 1: DNA/RNA (5'-D(*C)-R(P*G)-D(P*C)-R(P*G)-D(P*C)-R(P*G)-3')

Chain D: 

There are no outlier residues recorded for this chain.

• Molecule 1: DNA/RNA (5'-D(*C)-R(P*G)-D(P*C)-R(P*G)-D(P*C)-R(P*G)-3')

Chain E: 

There are no outlier residues recorded for this chain.

• Molecule 1: DNA/RNA (5'-D(*C)-R(P*G)-D(P*C)-R(P*G)-D(P*C)-R(P*G)-3')

Chain F:

• Molecule 1: DNA/RNA (5'-D(*C)-R(P*G)-D(P*C)-R(P*G)-D(P*C)-R(P*G)-3')

Chain G:

There are no outlier residues recorded for this chain.

• Molecule 1: DNA/RNA (5'-D(*C)-R(P*G)-D(P*C)-R(P*G)-D(P*C)-R(P*G)-3')

Chain H:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	35.72Å 44.10Å 35.72Å 90.00° 119.98° 90.00°	Depositor
Resolution (Å)	30.94 – 1.09 30.94 – 1.09	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.94-1.09) 99.7 (30.94-1.09)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.63 (at 1.09Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.115 , 0.169 0.118 , 0.149	Depositor DCC
R_{free} test set	996 reflections (2.49%)	wwPDB-VP
Wilson B-factor (Å^2)	8.2	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 84.4	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.239 for -h-l,k,h 0.239 for l,k,-h-l 0.254 for h,-k,-h-l 0.480 for -h-l,-k,l 0.251 for l,-k,h	Xtriage
Reported twinning fraction	0.279 for H, K, L 0.103 for L, -K, H 0.275 for H+L, -K, -L 0.113 for L, K, -H-L 0.110 for -H-L, K, H 0.121 for -H, -K, H+L	Depositor
Outliers	2 of 40052 reflections (0.005%)	Xtriage
F_o,F_c correlation	0.98	EDS
Total number of atoms	1278	wwPDB-VP
Average B, all atoms (Å^2)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 61.33 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3321e-05. The detected translational NCS is most*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

likely also responsible for the elevated intensity ratio.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.53	0/184	1.34	0/284
1	B	1.25	0/137	1.15	0/211
1	C	1.55	0/184	1.53	2/284 (0.7%)
1	D	1.41	0/137	1.12	0/211
1	E	1.39	0/137	1.11	0/211
1	F	1.57	0/249	1.16	0/385
1	G	1.39	0/137	1.11	0/211
1	H	1.38	0/231	1.21	0/357
All	All	1.45	0/1396	1.23	2/2154 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	2[A]	G	P-O3'-C3'	7.52	128.73	119.70
1	C	2[B]	G	P-O3'-C3'	7.52	128.73	119.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	130	0	60	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	123	0	68	0	0
1	C	130	0	60	1	0
1	D	123	0	68	0	0
1	E	123	0	68	0	0
1	F	144	0	52	1	0
1	G	123	0	68	0	0
1	H	137	0	52	1	0
2	A	6	0	0	3	0
2	B	1	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	1	0
2	F	3	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	A	31	0	0	1	0
3	B	21	0	0	0	0
3	C	30	0	0	0	0
3	D	28	0	0	1	0
3	E	38	0	0	0	0
3	F	21	0	0	1	0
3	G	36	0	0	0	0
3	H	24	0	0	1	0
All	All	1278	0	496	7	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:102:BA:BA	3:D:205:HOH:O	1.56	1.10
1:F:11[A]:DC:OP1	3:F:202:HOH:O	1.68	1.09
1:H:11[A]:DC:OP1	3:H:202:HOH:O	1.74	1.05
2:A:105:BA:BA	2:A:106:BA:BA	0.98	0.97
2:A:104:BA:BA	2:A:105:BA:BA	1.70	0.87
2:A:105:BA:BA	3:A:216:HOH:O	1.94	0.75
1:C:6:G:N3	1:C:6:G:H2'	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

There are no protein molecules in this entry.

5.3.2 Protein sidechains (i)

There are no protein molecules in this entry.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 16 ligands modelled in this entry, 16 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	6/6 (100%)	-0.59	0	100	100	8, 10, 12, 15	0
1	B	6/6 (100%)	-0.58	0	100	100	10, 12, 13, 14	0
1	C	6/6 (100%)	-0.45	0	100	100	8, 9, 13, 14	0
1	D	6/6 (100%)	-0.60	0	100	100	10, 11, 13, 14	0
1	E	6/6 (100%)	-0.50	0	100	100	10, 12, 13, 15	0
1	F	6/6 (100%)	-0.59	0	100	100	8, 10, 13, 14	0
1	G	6/6 (100%)	-0.50	0	100	100	10, 12, 13, 15	0
1	H	6/6 (100%)	-0.54	0	100	100	9, 10, 13, 14	0
All	All	48/48 (100%)	-0.54	0	100	100	8, 11, 14, 15	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	BA	D	102	1/1	0.98	0.05	18,18,18,18	1
2	BA	A	101	1/1	0.99	0.05	18,18,18,18	1
2	BA	A	103	1/1	1.00	0.02	11,11,11,11	1
2	BA	A	104	1/1	1.00	0.02	8,8,8,8	1
2	BA	A	105	1/1	1.00	0.06	20,20,20,20	1
2	BA	A	106	1/1	1.00	0.03	13,13,13,13	1
2	BA	B	101	1/1	1.00	0.01	9,9,9,9	1
2	BA	C	101	1/1	1.00	0.03	7,7,7,7	1
2	BA	C	102	1/1	1.00	0.03	11,11,11,11	1
2	BA	D	101	1/1	1.00	0.02	9,9,9,9	1
2	BA	A	102	1/1	1.00	0.02	8,8,8,8	1
2	BA	F	101	1/1	1.00	0.03	10,10,10,10	1
2	BA	F	102	1/1	1.00	0.03	10,10,10,10	1
2	BA	F	103	1/1	1.00	0.02	14,14,14,14	1
2	BA	G	101	1/1	1.00	0.01	8,8,8,8	1
2	BA	H	101	1/1	1.00	0.02	9,9,9,9	1

6.5 Other polymers (i)

There are no such residues in this entry.