

# Full wwPDB X-ray Structure Validation Report (i)

#### Aug 31, 2020 - 09:43 AM BST

PDB ID	:	1EDB
$\operatorname{Title}$	:	CRYSTALLOGRAPHIC AND FLUORESCENCE STUDIES OF THE IN-
		TERACTION OF HALOALKANE DEHALOGENASE WITH HALIDE
		IONS: STUDIES WITH HALIDE COMPOUNDS REVEAL A HALIDE
		BINDING SITE IN THE ACTIVE SITE
Authors	:	Verschueren, K.H.G.; Dijkstra, B.W.
Deposited on	:	1993-05-13
Resolution	:	2.01  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	NOT EXECUTED
$\mathrm{EDS}$	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$		
Clashscore	141614	9178 (2.00-2.00)		
Ramachandran outliers	138981	9054 (2.00-2.00)		
Sidechain outliers	138945	9053 (2.00-2.00)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain					
1	A	310	76%	20%	••			



## 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2671 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called HALOALKANE DEHALOGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	310	Total 2479	C 1596	N 406	O 462	S 15	0	0	0

• Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Cl 1 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	191	Total O 191 191	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: HALOALKANE DEHALOGENASE





## 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 21 21 2	Depositor	
Cell constants	95.40Å $72.90$ Å $41.40$ Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor	
Resolution (Å)	15.00 - 2.01	Depositor	
% Data completeness	(Not available) $(15.00-2.01)$	Depositor	
(in resolution range)	(1000 available) (10.00 2.01)		
$R_{merge}$	(Not available)	Depositor	
$R_{sym}$	(Not available)	Depositor	
Refinement program	TNT	Depositor	
$R, R_{free}$	0.175 , (Not available)	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	2671	wwPDB-VP	
Average B, all atoms $(Å^2)$	12.0	wwPDB-VP	



## 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.70	1/2552~(0.0%)	1.74	61/3470~(1.8%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	Ideal(Å)
1	А	308	GLU	CG-CD	5.91	1.60	1.51

All (61) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	А	307	ALA	CB-CA-C	-16.17	85.85	110.10
1	А	212	ASP	CB-CG-OD2	-11.08	108.33	118.30
1	А	188	ASP	CB-CG-OD1	10.54	127.79	118.30
1	А	229	ARG	NE-CZ-NH1	10.48	125.54	120.30
1	А	170	ASP	CB-CG-OD1	9.94	127.24	118.30
1	А	184	ASP	CB-CG-OD2	-9.93	109.36	118.30
1	А	137	ASP	CB-CG-OD2	-9.62	109.64	118.30
1	А	193	ARG	NE-CZ-NH1	9.26	124.93	120.30
1	А	40	ASP	CB-CG-OD1	9.06	126.45	118.30
1	А	93	ASP	CB-CG-OD2	-8.89	110.30	118.30
1	А	308	GLU	CB-CA-C	8.82	128.04	110.40
1	А	103	ARG	NE-CZ-NH1	8.69	124.64	120.30



Continued from previous page							
Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	116	ARG	NE-CZ-NH1	8.55	124.57	120.30
1	А	45	ASP	CB-CG-OD2	-8.51	110.65	118.30
1	А	124	ASP	CB-CG-OD1	8.48	125.93	118.30
1	А	212	ASP	CB-CG-OD1	8.15	125.64	118.30
1	А	143	ARG	NE-CZ-NH2	-8.10	116.25	120.30
1	А	188	ASP	CB-CG-OD2	-8.10	111.01	118.30
1	А	124	ASP	CB-CG-OD2	-7.79	111.29	118.30
1	А	45	ASP	CB-CG-OD1	7.61	125.14	118.30
1	А	89	ASP	CB-CG-OD2	-7.48	111.56	118.30
1	А	16	ASP	CB-CG-OD2	-7.46	111.58	118.30
1	А	40	ASP	CB-CG-OD2	-7.22	111.80	118.30
1	А	16	ASP	CB-CG-OD1	7.15	124.73	118.30
1	А	269	TYR	CB-CG-CD1	-7.13	116.72	121.00
1	А	27	ASP	CB-CG-OD1	7.11	124.70	118.30
1	А	286	ASP	CB-CG-OD1	6.76	124.39	118.30
1	А	116	ARG	NE-CZ-NH2	-6.67	116.96	120.30
1	А	184	ASP	CB-CG-OD1	6.67	124.30	118.30
1	А	286	ASP	CB-CG-OD2	-6.48	112.47	118.30
1	А	266	ASP	CB-CG-OD2	-6.47	112.48	118.30
1	А	48	ASP	CB-CG-OD1	6.41	124.07	118.30
1	А	114	ASP	CB-CG-OD1	6.38	124.04	118.30
1	А	204	SER	CB-CA-C	5.99	121.48	110.10
1	А	199	THR	CA-CB-CG2	-5.99	104.01	112.40
1	А	307	ALA	CA-C-N	-5.93	104.15	117.20
1	А	243	PHE	CB-CG-CD2	-5.93	116.65	120.80
1	А	206	TYR	CB-CG-CD2	-5.88	117.47	121.00
1	А	134	PRO	N-CA-CB	5.84	110.31	103.30
1	А	13	SER	CB-CA-C	-5.67	99.33	110.10
1	А	57	PRO	N-CA-C	-5.66	97.38	112.10
1	А	293	GLU	CB-CA-C	5.62	121.65	110.40
1	А	175	TRP	CA-CB-CG	5.54	124.23	113.70
1	А	89	ASP	CB-CG-OD1	5.54	123.29	118.30
1	А	154	ASP	CB-CG-OD1	5.52	123.27	118.30
1	А	274	LEU	CB-CG-CD2	-5.44	101.75	111.00
1	А	308	GLU	C-N-CA	5.41	135.22	121.70
1	А	48	ASP	CB-CG-OD2	-5.40	113.44	118.30
1	А	269	TYR	CB-CG-CD2	5.28	124.17	121.00
1	А	119	THR	N-CA-C	-5.25	96.82	111.00
1	А	230	ASP	CB-CG-OD1	5.24	123.02	118.30
1	А	154	ASP	CB-CG-OD2	-5.18	113.64	118.30
1	А	308	GLU	CB-CG-CD	5.16	128.13	114.20
1	А	272	LYS	O-C-N	-5.07	114.59	122.70



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	А	103	ARG	NE-CZ-NH2	-5.06	117.77	120.30
1	А	26	ASP	CB-CG-OD2	-5.06	113.75	118.30
1	А	257	GLY	N-CA-C	-5.06	100.45	113.10
1	А	216	GLN	N-CA-C	5.05	124.65	111.00
1	А	114	ASP	CB-CG-OD2	-5.04	113.76	118.30
1	А	137	ASP	CB-CG-OD1	5.03	122.83	118.30
1	А	229	ARG	NH1-CZ-NH2	-5.01	113.89	119.40

Continued from previous page...

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	307	ALA	Mainchain

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2479	0	2379	30	0
2	А	1	0	0	0	0
3	А	191	0	0	5	0
All	All	2671	0	2379	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:292:GLN:HE21	1:A:292:GLN:H	1.09	0.93
1:A:225:MET:HE1	3:A:592:HOH:O	1.86	0.75
1:A:153:THR:HG21	1:A:158:GLN:HB2	1.68	0.75
1:A:142:LYS:HG2	1:A:143:ARG:HG3	1.69	0.74
1:A:112:ARG:HH11	1:A:112:ARG:HG3	1.54	0.71
1:A:124:ASP:OD1	1:A:289:HIS:NE2	2.26	0.66



A 4 1	A 4 a ma D	Interatomic	Clash
Atom-1	Atom-2	$distance ( m \AA)$	overlap (Å)
1:A:220:ARG:HD3	3:A:470:HOH:O	1.97	0.64
1:A:99:PHE:HA	3:A:592:HOH:O	1.98	0.63
1:A:308:GLU:HB2	3:A:587:HOH:O	2.02	0.59
1:A:309:THR:HG22	1:A:309:THR:O	2.04	0.57
1:A:262:LEU:HD23	3:A:483:HOH:O	2.04	0.56
1:A:154:ASP:HB2	1:A:155:PRO:HD2	1.88	0.54
1:A:112:ARG:HH11	1:A:112:ARG:CG	2.21	0.54
1:A:292:GLN:NE2	1:A:292:GLN:H	1.92	0.53
1:A:309:THR:O	1:A:310:GLU:HG2	2.10	0.52
1:A:154:ASP:CG	1:A:157:THR:HG22	2.31	0.51
1:A:307:ALA:HB3	1:A:308:GLU:HG3	1.93	0.51
1:A:154:ASP:HB2	1:A:155:PRO:CD	2.45	0.47
1:A:251:GLN:HE22	1:A:309:THR:HB	1.82	0.45
1:A:151:LEU:HD13	1:A:241:ILE:HG13	1.99	0.45
1:A:292:GLN:HE21	1:A:292:GLN:N	1.93	0.43
1:A:112:ARG:NH1	1:A:112:ARG:CG	2.78	0.43
1:A:161:PHE:N	1:A:161:PHE:CD1	2.86	0.43
1:A:272:LYS:HG2	1:A:272:LYS:O	2.12	0.43
1:A:154:ASP:OD1	1:A:157:THR:HG22	2.20	0.42
1:A:125:TRP:CH2	1:A:225:MET:CE	3.01	0.42
1:A:31:TYR:N	1:A:32:PRO:CD	2.83	0.42
1:A:251:GLN:NE2	1:A:309:THR:HB	2.36	0.41
1:A:154:ASP:OD2	1:A:157:THR:HG22	2.20	0.40
1:A:5:ILE:HG21	1:A:5:ILE:HD13	1.74	0.40

Continued from previous page...

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	308/310~(99%)	291~(94%)	16~(5%)	1 (0%)	41 37



All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	148	ASN

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	262/262~(100%)	246~(94%)	16 (6%)	18 14

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	А	1	MET
1	А	9	ASP
1	А	14	ASN
1	А	15	LEU
1	А	34	LEU
1	А	73	SER
1	А	113	LEU
1	А	116	ARG
1	А	123	GLN
1	А	157	THR
1	А	162	SER
1	А	204	SER
1	А	222	PHE
1	А	266	ASP
1	А	292	GLN
1	А	308	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	14	ASN
1	А	104	ASN
1	А	123	GLN



Continued from previous page...

Mol	Chain	Res	Type
1	А	251	GLN
1	А	292	GLN
1	А	297	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

#### 5.7 Other polymers (i)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

## 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

