Full wwPDB X-ray Structure Validation Report (i)

Mar 13, 2024 – 04:24 PM JST

PDB ID	:	5EE9
Title	:	Complex structure of OSYCHF1 with GMP-PNP
Authors	:	Li, X.; Chen, Z.
Deposited on	:	2015-10-22
Resolution	:	2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	84% 7% 10%
1	B	395	2% 83% 7% • 9%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5729 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Obg-like ATPase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	357	2717	1761	439	506	11	0	0	0
1	B	359	2716	1756	442	507	11	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q6Z1J6
B	0	SER	-	expression tag	UNP Q6Z1J6

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter code: GNP) (formula: $C_{10}H_{17}N_6O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	32	10	6	13	3	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total K 1 1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 6 3 3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	127	Total O 127 127	0	0
6	B	129	Total O 129 129	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Obg-like ATPase 1

Chain A:



- Molecule 1: Obg-like ATPase 1

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	70.45Å 112.06Å 115.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.75 41.53 – 2.75	Depositor EDS
% Data completeness (in resolution range)	97.4 (50.00-2.75) 97.5 (41.53-2.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.69 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.229 , 0.262 0.229 , 0.259	Depositor DCC
R_{free} test set	1133 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å^2)	45.6	Xtriage
Anisotropy	0.132	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.31 , 38.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	5729	wwPDB-VP
Average B, all atoms (Å^2)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 54.96 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.3882e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, GNP, MG, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.42	2/2777 (0.1%)	0.46	0/3767
1	B	0.43	2/2778 (0.1%)	0.48	0/3766
All	All	0.42	4/5555 (0.1%)	0.47	0/7533

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	250	TRP	CD2-CE2	5.10	1.47	1.41
1	A	206	TRP	CD2-CE2	5.06	1.47	1.41
1	B	75	TRP	CD2-CE2	5.06	1.47	1.41
1	B	206	TRP	CD2-CE2	5.02	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2717	0	2551	11	0
1	B	2716	0	2501	17	0
2	A	32	0	13	0	0
3	A	1	0	0	0	0
4	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	6	0	8	1	0
6	A	127	0	0	0	0
6	B	129	0	0	0	0
All	All	5729	0	5073	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:328:GLN:HE21	1:B:328:GLN:H	1.25	0.83
1:B:328:GLN:HE21	1:B:328:GLN:N	1.93	0.65
1:B:334:HIS:HD2	1:B:336:ASP:H	1.47	0.62
1:B:75:TRP:CH2	1:B:353:LYS:HA	2.38	0.57
1:B:330:ALA:HB1	1:B:337:PHE:HB3	1.87	0.56
1:A:330:ALA:HB1	1:A:337:PHE:HB3	1.90	0.54
1:B:336:ASP:HA	1:B:339:ARG:HG2	1.90	0.53
1:A:229:VAL:HG13	1:A:261:PRO:HA	1.91	0.53
1:A:322:ARG:O	1:A:323:GLN:HB2	2.12	0.49
1:B:187:LEU:HD21	1:B:214:LEU:HG	1.93	0.49
1:A:352:LEU:HD12	1:A:381:ILE:HD13	1.94	0.49
1:B:37:LYS:HB2	5:B:402:GOL:H12	1.95	0.48
1:B:144:VAL:HA	1:B:147:LEU:HD12	1.95	0.48
1:B:169:LEU:O	1:B:173:MET:HB2	2.13	0.48
1:A:337:PHE:N	1:A:337:PHE:CD1	2.84	0.46
1:A:337:PHE:N	1:A:337:PHE:HD1	2.14	0.46
1:A:75:TRP:CZ3	1:A:79:LEU:HD22	2.52	0.45
1:A:191:VAL:HG21	1:A:214:LEU:HD21	1.98	0.44
1:B:35:VAL:HG12	1:B:127:ARG:HB2	2.00	0.43
1:B:26:LYS:HD3	1:B:93:ASN:ND2	2.34	0.42
1:B:187:LEU:HD13	1:B:210:ASP:HB3	2.01	0.42
1:B:142:ASP:HA	1:B:143:PRO:HD2	1.92	0.42
1:B:223:LYS:HA	1:B:224:PRO:HD3	1.96	0.42
1:A:270:LEU:HA	1:A:273:MET:HE3	2.01	0.41
1:A:203:PHE:CZ	1:A:255:GLY:HA3	2.56	0.41
1:B:26:LYS:HD3	1:B:93:ASN:HD21	1.85	0.41
1:B:75:TRP:HZ3	1:B:352:LEU:HD13	1.85	0.40
1:A:140:SER:HB2	1:A:145:ARG:NH2	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	351/395 (89%)	341 (97%)	9 (3%)	1 (0%)	41	60
1	B	353/395 (89%)	346 (98%)	7 (2%)	0	100	100
All	All	704/790 (89%)	687 (98%)	16 (2%)	1 (0%)	51	75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	367	TYR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/336 (80%)	261 (97%)	7 (3%)	46	66
1	B	257/336 (76%)	247 (96%)	10 (4%)	32	52
All	All	525/672 (78%)	508 (97%)	17 (3%)	39	59

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	LEU
1	A	202	ARG
1	A	225	VAL
1	A	234	LYS
1	A	252	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	264	CYS
1	A	352	LEU
1	B	22	SER
1	B	168	ASP
1	B	173	MET
1	B	220	LEU
1	B	244	LEU
1	B	252	GLN
1	B	264	CYS
1	B	328	GLN
1	B	352	LEU
1	B	387	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	ASN
1	A	194	HIS
1	A	285	ASN
1	A	323	GLN
1	A	387	ASN
1	B	164	ASN
1	B	194	HIS
1	B	328	GLN
1	B	334	HIS
1	B	369	GLN
1	B	387	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	GOL	B	402	-	5,5,5	0.33	0	5,5,5	0.39	0
2	GNP	A	401	3	29,34,34	1.78	5 (17%)	33,54,54	2.00	9 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	402	-	-	4/4/4/4	-
2	GNP	A	401	3	-	3/14/38/38	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	GNP	PB-N3B	4.45	1.75	1.63
2	A	401	GNP	C5-C6	4.38	1.48	1.41
2	A	401	GNP	PG-N3B	4.33	1.74	1.63
2	A	401	GNP	C5-C4	2.63	1.47	1.40
2	A	401	GNP	O4'-C1'	2.01	1.43	1.41

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	401	GNP	C2-N3-C4	5.04	121.11	115.36
2	A	401	GNP	C2-N1-C6	4.09	122.43	115.93
2	A	401	GNP	C5-C6-N1	-4.06	117.88	123.43
2	A	401	GNP	C4-C5-C6	-3.54	117.42	120.80
2	A	401	GNP	N3-C2-N1	-3.40	122.69	127.22
2	A	401	GNP	PB-O3A-PA	-3.23	121.24	132.62

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	401	GNP	C3'-C2'-C1'	2.97	105.45	100.98
2	A	401	GNP	O1G-PG-N3B	-2.67	107.83	111.77
2	A	401	GNP	C4-C5-N7	-2.66	106.63	109.40

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	GNP	PB-N3B-PG-O1G
2	A	401	GNP	PG-N3B-PB-O1B
2	A	401	GNP	PG-N3B-PB-O3A
5	B	402	GOL	O1-C1-C2-C3
5	B	402	GOL	C1-C2-C3-O3
5	B	402	GOL	O1-C1-C2-O2
5	B	402	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	402	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand GNP A 401



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	357/395 (90%)	0.05	1 (0%)	94	96	21, 42, 60, 78	0
1	B	359/395 (90%)	0.14	8 (2%)	62	70	18, 44, 69, 79	0
All	All	716/790 (90%)	0.10	9 (1%)	77	84	18, 43, 65, 79	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	173	MET	3.9
1	B	187	LEU	2.9
1	B	172	SER	2.8
1	B	34	ASN	2.7
1	B	169	LEU	2.6
1	B	159	ILE	2.4
1	B	177	ASN	2.2
1	A	111	ALA	2.2
1	B	342	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	GOL	B	402	6/6	0.87	0.24	49,52,53,53	0
4	K	B	401	1/1	0.88	0.10	54,54,54,54	0
3	MG	A	402	1/1	0.93	0.29	46,46,46,46	0
2	GNP	A	401	32/32	0.94	0.16	50,54,57,58	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GNP A 401:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.