

Full wwPDB X-ray Structure Validation Report (i)

May 15, 2020 - 01:49 pm BST

PDB ID	:	1EGQ
Title	:	ENHANCEMENT OF ENZYME ACTIVITY THROUGH THREE-PHASE
		PARTITIONING: CRYSTAL STRUCTURE OF A MODIFIED SERINE
		PROTEINASE AT 1.5 A RESOLUTION
Authors	:	Singh, R.K.; Gourinath, S.; Sharma, S.; Ray, I.; Gupta, M.N.; Singh, T.P.
Deposited on		
Resolution	:	1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


$\operatorname{MolProbity}$:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{llllllllllllllllllllllllllllllllllll$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495(1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	А	279	82%	15%	•



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2282 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called PROTEINASE K.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	279	Total 2029	C 1247	N 357	0 415	S 10	0	0	0

• Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	2	Total Ca 2 2	0	0

• Molecule 3 is ACETIC ACID (three-letter code: ACY) (formula: $C_2H_4O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0



• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	243	Total O 243 243	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PROTEINASE K





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants	68.29Å 68.29 Å 106.36 Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 - 1.55	Depositor
% Data completeness	93.0(10.00-1.55)	Depositor
(in resolution range)	35.0 (10.00-1.55)	Depositor
R_{merge}	0.07	Depositor
R _{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.193 , 0.206	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2282	wwPDB-VP
Average B, all atoms $(Å^2)$	19.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.31	0/2068	0.61	1/2810~(0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms Z		$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	А	133	LEU	CA-CB-CG	5.08	126.99	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2029	0	1935	45	0
2	А	2	0	0	0	0
3	А	8	0	8	2	0
4	А	243	0	0	4	1
All	All	2282	0	1943	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance (\AA)	overlap (Å)	
1:A:58:THR:HG22	1:A:60:TYR:H	1.37	0.90	
1:A:103:GLN:HG2	1:A:106:THR:HG23	1.63	0.80	
1:A:80:ARG:HH11	1:A:80:ARG:HB3	1.46	0.79	
1:A:58:THR:HG21	1:A:62:SER:O	1.83	0.79	
1:A:46:HIS:HD2	1:A:48:GLU:H	1.32	0.74	
1:A:207:SER:HB3	4:A:556:HOH:O	1.88	0.72	
1:A:261:LEU:H	1:A:270:ASN:HD21	1.37	0.72	
1:A:97:ASP:HA	3:A:301:ACY:H2	1.79	0.64	
1:A:80:ARG:NH1	1:A:80:ARG:HB3	2.15	0.62	
1:A:235:ALA:O	1:A:239:THR:HG23	2.00	0.61	
1:A:255:THR:HB	1:A:276:ASN:ND2	2.15	0.61	
1:A:255:THR:HB	1:A:276:ASN:HD21	1.65	0.60	
1:A:125:LYS:CG	1:A:239:THR:HA	2.31	0.60	
1:A:180:VAL:HG21	1:A:230:VAL:HG21	1.85	0.59	
1:A:27:GLU:CD	1:A:27:GLU:H	2.06	0.59	
1:A:225:MET:O	1:A:229:HIS:HD2	1.86	0.58	
1:A:64:ARG:HB2	1:A:64:ARG:HH11	1.69	0.57	
1:A:207:SER:HB2	4:A:719:HOH:O	2.05	0.55	
1:A:258:LYS:HD2	1:A:271:LEU:CD2	2.36	0.54	
1:A:24:TYR:HB3	1:A:277:TYR:CD2	2.44	0.53	
1:A:103:GLN:HG2	1:A:106:THR:CG2	2.35	0.52	
1:A:221:SER:O	3:A:303:ACY:H1	2.09	0.52	
1:A:103:GLN:HG3	1:A:105:SER:OG	2.08	0.52	
1:A:201:ILE:HG13	1:A:202:PHE:N	2.25	0.52	
1:A:164:ALA:H	1:A:194:ASN:ND2	2.09	0.51	
1:A:46:HIS:HE1	1:A:216:SER:O	1.94	0.51	
1:A:30:GLY:HA2	1:A:239:THR:HG21	1.93	0.50	
1:A:30:GLY:C	1:A:239:THR:HG21	2.33	0.49	
1:A:185:ARG:HD3	4:A:582:HOH:O	2.11	0.49	
1:A:80:ARG:HH11	1:A:80:ARG:CB	2.21	0.48	
1:A:125:LYS:HG2	1:A:239:THR:HA	1.96	0.47	
1:A:3:GLN:HE22	1:A:86:LYS:HZ1	1.61	0.47	
1:A:162:ASN:H	1:A:194:ASN:ND2	2.14	0.46	
1:A:33:SER:HB3	1:A:239:THR:HG22	1.98	0.46	
1:A:46:HIS:CD2	1:A:48:GLU:H	2.23	0.44	
1:A:180:VAL:HG21	1:A:230:VAL:CG2	2.48	0.43	
1:A:164:ALA:H	1:A:194:ASN:HD22	1.67	0.43	
1:A:24:TYR:HB3	1:A:277:TYR:HD2	1.84	0.43	
1:A:75:GLY:HA2	1:A:79:SER:HB3	2.00	0.43	
1:A:106:THR:HG21	4:A:637:HOH:O	2.19	0.42	
1:A:256:ALA:O	1:A:258:LYS:HD3	2.19	0.42	

Continued on next page...



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
1:A:180:VAL:CG2	1:A:230:VAL:HG21	2.48	0.42	
1:A:103:GLN:HB3	1:A:103:GLN:HE21	1.64	0.42	
1:A:118:LYS:NZ	1:A:119:ASN:HD21	2.17	0.42	
1:A:46:HIS:CD2	1:A:48:GLU:HB2	2.57	0.40	

Continued from previous page...

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
4:A:615:HOH:O	4:A:615:HOH:O[8_665]	2.17	0.03	

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percen	tiles
1	А	277/279~(99%)	267~(96%)	10 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Rotameric Outlie		Outliers	Percentiles	
1	А	213/213~(100%)	204~(96%)	9 (4%)	30 5	



Mol	Chain	Res	Type
1	А	103	GLN
1	А	106	THR
1	А	120	ASN
1	А	125	LYS
1	А	133	LEU
1	А	167	ARG
1	А	225	MET
1	А	240	LEU
1	А	270	ASN

All (9) residues with a non-rotameric sidechain are listed below:

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	3	GLN
1	А	5	ASN
1	А	31	GLN
1	А	46	HIS
1	А	103	GLN
1	А	119	ASN
1	А	162	ASN
1	А	168	ASN
1	А	194	ASN
1	А	229	HIS
1	А	257	ASN
1	А	270	ASN
1	А	276	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.



5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	True	Chain	Res	Link	B	ond len	gths	B	ond ang	gles
	Type	Cham	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACY	А	303	-	$1,\!3,\!3$	2.32	1 (100%)	$0,\!3,\!3$	0.00	-
3	ACY	А	301	-	1,3,3	1.06	0	$_{0,3,3}$	0.00	-

All (1) bond length outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
3	А	303	ACY	CH3-C	2.32	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	303	ACY	1	0
3	А	301	ACY	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

