



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 16, 2023 – 07:47 AM EDT

PDB ID : 2EG5  
Title : The structure of xanthosine methyltransferase  
Authors : McCarthy, A.A.; McCarthy, J.G.  
Deposited on : 2007-02-28  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	
1	C	372	
1	E	372	
1	G	372	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	XTS	A	502	-	-	X	-
3	XTS	E	2502	-	-	X	-
3	XTS	G	3502	-	-	X	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 11430 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xanthosine methyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	344	Total 2701	C 1744	N 449	O 495	S 13	0	3	0
1	C	344	Total 2681	C 1734	N 444	O 490	S 13	0	3	0
1	E	346	Total 2698	C 1745	N 444	O 494	S 15	0	4	0
1	G	343	Total 2667	C 1725	N 440	O 489	S 13	0	3	0

- Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula:  $C_{14}H_{20}N_6O_5S$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	Total 26	C 14	N 6	O 5	S 1	0	0
2	C	1	Total 26	C 14	N 6	O 5	S 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			S
2	E	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	G	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 3 is 9-[(2R,3R,4S,5R)-3,4-DIHYDROXY-5-(HYDROXYMETHYL)OXOLAN-2-YL]-3H-PURINE-2,6-DIONE (three-letter code: XTS) (formula: C<sub>10</sub>H<sub>12</sub>N<sub>4</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	Total	C	N	O	0	0
			20	10	4	6		
3	C	1	Total	C	N	O	0	0
			20	10	4	6		
3	E	1	Total	C	N	O	0	0
			20	10	4	6		
3	G	1	Total	C	N	O	0	0
			20	10	4	6		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	156	Total	O	0	0
			156	156		
4	C	132	Total	O	0	0
			132	132		
4	E	114	Total	O	0	0
			114	114		

Continued on next page...


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	G	97	Total	O	0	0
			97	97		





● Molecule 1: Xanthosine methyltransferase





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.59Å 119.80Å 116.40Å 90.00° 102.16° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 29.95 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.7 (30.00-2.20) 86.6 (29.95-2.20)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.84 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.230 , 0.284 0.231 , 0.281	Depositor DCC
$R_{free}$ test set	3830 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.7	Xtrriage
Anisotropy	0.155	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 44.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	0.089 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	11430	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.23 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7398e-03.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAH, XTS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.80	1/2765 (0.0%)	0.80	1/3739 (0.0%)
1	C	0.79	0/2745	0.79	1/3714 (0.0%)
1	E	0.67	0/2765	0.74	3/3740 (0.1%)
1	G	0.73	0/2731	0.76	0/3698
All	All	0.75	1/11006 (0.0%)	0.77	5/14891 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	132	CYS	CB-SG	-5.60	1.72	1.81

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	101	ASP	CB-CG-OD2	5.55	123.29	118.30
1	E	173	GLU	OE1-CD-OE2	-5.53	116.67	123.30
1	C	360	ILE	CB-CA-C	-5.24	101.13	111.60
1	A	173	GLU	OE1-CD-OE2	-5.11	117.17	123.30
1	E	360	ILE	CB-CA-C	-5.03	101.55	111.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	302	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2701	0	2702	69	0
1	C	2681	0	2667	71	0
1	E	2698	0	2686	73	0
1	G	2667	0	2634	66	0
2	A	26	0	19	2	0
2	C	26	0	19	3	0
2	E	26	0	19	2	0
2	G	26	0	19	2	0
3	A	20	0	12	9	0
3	C	20	0	12	6	0
3	E	20	0	12	8	0
3	G	20	0	12	7	0
4	A	156	0	0	9	0
4	C	132	0	0	11	0
4	E	114	0	0	11	0
4	G	97	0	0	6	0
All	All	11430	0	10813	278	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 278 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:24:TYR:HB3	3:E:2502:XTS:H5'2	1.21	1.17
1:A:24:TYR:HB3	3:A:502:XTS:H5'2	1.26	1.16
1:C:290:LEU:CD2	1:C:359:LEU:HD11	1.74	1.16
1:E:236:ARG:HD3	1:E:244:MET:HE1	1.28	1.14
1:G:236:ARG:HD3	1:G:244:MET:HE1	1.28	1.09

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/372 (91%)	323 (95%)	13 (4%)	3 (1%)	17	16
1	C	339/372 (91%)	324 (96%)	12 (4%)	3 (1%)	17	16
1	E	342/372 (92%)	328 (96%)	11 (3%)	3 (1%)	17	16
1	G	338/372 (91%)	324 (96%)	11 (3%)	3 (1%)	17	16
All	All	1358/1488 (91%)	1299 (96%)	47 (4%)	12 (1%)	17	16

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	177	GLY
1	C	177	GLY
1	E	177	GLY
1	G	177	GLY
1	A	23	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/327 (90%)	263 (90%)	30 (10%)	7	6
1	C	287/327 (88%)	255 (90%)	32 (11%)	6	5
1	E	290/327 (89%)	261 (90%)	29 (10%)	7	7
1	G	283/327 (86%)	256 (90%)	27 (10%)	8	8
All	All	1153/1308 (88%)	1035 (90%)	118 (10%)	7	6

5 of 118 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	331	GLU
1	G	290	LEU
1	E	161	GLN
1	G	277	CYS
1	G	172	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	25	ASN
1	E	77	GLN
1	G	125	ASN
1	G	25	ASN
1	G	96	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SAH	G	3501	-	24,28,28	1.51	4 (16%)	25,40,40	1.94	5 (20%)
2	SAH	C	1501	-	24,28,28	1.32	2 (8%)	25,40,40	1.95	6 (24%)
2	SAH	A	501	-	24,28,28	1.58	3 (12%)	25,40,40	1.95	4 (16%)
3	XTS	G	3502	-	16,22,22	1.89	4 (25%)	20,33,33	3.58	9 (45%)
2	SAH	E	2501	-	24,28,28	1.29	2 (8%)	25,40,40	1.93	5 (20%)
3	XTS	C	1502	-	16,22,22	1.82	5 (31%)	20,33,33	3.34	8 (40%)
3	XTS	A	502	-	16,22,22	2.09	4 (25%)	20,33,33	3.05	8 (40%)
3	XTS	E	2502	-	16,22,22	1.88	3 (18%)	20,33,33	2.90	6 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAH	G	3501	-	-	1/11/31/31	0/3/3/3
2	SAH	C	1501	-	-	0/11/31/31	0/3/3/3
2	SAH	A	501	-	-	1/11/31/31	0/3/3/3
3	XTS	G	3502	-	-	2/2/22/22	0/3/3/3
2	SAH	E	2501	-	-	0/11/31/31	0/3/3/3
3	XTS	C	1502	-	-	2/2/22/22	0/3/3/3
3	XTS	A	502	-	-	2/2/22/22	0/3/3/3
3	XTS	E	2502	-	-	2/2/22/22	0/3/3/3

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	XTS	O4'-C1'	5.52	1.48	1.41
2	A	501	SAH	C2-N3	4.80	1.39	1.32
2	G	3501	SAH	C2-N3	4.70	1.39	1.32
3	E	2502	XTS	O4'-C1'	4.68	1.47	1.41
2	A	501	SAH	C2-N1	4.37	1.42	1.33

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	3502	XTS	C6-N1-C2	-10.32	118.74	126.88
3	C	1502	XTS	C6-N1-C2	-9.69	119.24	126.88
3	G	3502	XTS	N3-C2-N1	8.31	120.72	115.90

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	XTS	C6-N1-C2	-7.64	120.85	126.88
3	E	2502	XTS	C6-N1-C2	-7.36	121.08	126.88

There are no chirality outliers.

5 of 10 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	502	XTS	O4'-C4'-C5'-O5'
3	E	2502	XTS	O4'-C4'-C5'-O5'
3	G	3502	XTS	O4'-C4'-C5'-O5'
3	C	1502	XTS	O4'-C4'-C5'-O5'
3	A	502	XTS	C3'-C4'-C5'-O5'

There are no ring outliers.



8 monomers are involved in 39 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	3501	SAH	2	0
2	C	1501	SAH	3	0
2	A	501	SAH	2	0
3	G	3502	XTS	7	0
2	E	2501	SAH	2	0
3	C	1502	XTS	6	0
3	A	502	XTS	9	0
3	E	2502	XTS	8	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	344/372 (92%)	-0.06	12 (3%) 44 42	30, 43, 65, 79	12 (3%)
1	C	344/372 (92%)	0.09	17 (4%) 29 28	31, 43, 65, 79	15 (4%)
1	E	346/372 (93%)	0.19	23 (6%) 18 17	31, 43, 65, 79	5 (1%)
1	G	343/372 (92%)	0.27	23 (6%) 17 16	31, 43, 65, 79	11 (3%)
All	All	1377/1488 (92%)	0.12	75 (5%) 25 24	30, 43, 66, 79	43 (3%)

The worst 5 of 75 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	177	GLY	10.9
1	C	174	LEU	10.8
1	E	176	ILE	9.9
1	G	176	ILE	8.9
1	C	172	THR	7.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	XTS	G	3502	20/20	0.69	0.26	64,65,67,67	2
3	XTS	E	2502	20/20	0.76	0.21	64,65,67,67	2
2	SAH	A	501	26/26	0.80	0.20	53,56,67,68	0
2	SAH	E	2501	26/26	0.83	0.20	53,57,67,68	0
3	XTS	A	502	20/20	0.84	0.17	64,65,67,67	2
3	XTS	C	1502	20/20	0.85	0.21	64,65,67,67	2
2	SAH	C	1501	26/26	0.87	0.17	53,56,67,68	0
2	SAH	G	3501	26/26	0.87	0.15	53,56,67,68	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around XTS E 2502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around SAH A 501:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around SAH E 2501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around XTS A 502:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)









**Electron density around XTS C 1502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around SAH C 1501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.