



Full wwPDB NMR Structure Validation Report i

Apr 20, 2024 – 06:32 PM EDT

PDB ID : 1EKH
Title : NMR STRUCTURE OF D(TTGGCCAA)2 BOUND TO CHROMOMYCIN-A3 AND COBALT
Authors : Gochin, M.
Deposited on : 2000-03-08

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA and RNA chains that are outliers for geometric criteria:

Mol	Chain	Compound	Res	Total models with violations	
				Chirality	Geometry
2	D	2GL	1	2	-
3	E	DDA	1	2	-

Continued on next page...

Continued from previous page...

Mol	Chain	Compound	Res	Total models with violations	
				Chirality	Geometry
3	F	DDA	1	2	-
5	B	CPH	23	3	-
5	A	CPH	33	3	-

2 Ensemble composition and analysis

This entry contains 6 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition [\(i\)](#)


There are 5 unique types of molecules in this entry. The entry contains 835 atoms, of which 346 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called DNA (5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3').

Mol	Chain	Residues	Atoms						Trace
1	A	8	Total	C	H	N	O	P	0
			253	78	92	30	46	7	

Mol	Chain	Residues	Atoms						Trace
1	B	8	Total	C	H	N	O	P	0
			253	78	92	30	46	7	


- Molecule 2 is an oligosaccharide called 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms						Trace
2	C	2	Total	C	H	O			0
			48	15	25	8			

Mol	Chain	Residues	Atoms						Trace
2	D	2	Total	C	H	O			0
			48	15	25	8			

- Molecule 3 is an oligosaccharide called 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose.




Mol	Chain	Residues	Atoms						Trace
3	E	3	Total	C	H	O			0
			67	21	35	11			

Mol	Chain	Residues	Atoms						Trace
3	F	3	Total	C	H	O			0
			67	21	35	11			

- Molecule 4 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms						
4	A	1	Total	Co					
			1	1					

- Molecule 5 is (1S)-5-deoxy-1-O-methyl-1-C-[(2R,3S)-3,5,7,10-tetrahydroxy-6-methyl-4-oxo-1,2,3,4-tetrahydroanthracen-2-yl]-D-xylulose (three-letter code: CPH) (formula: C₂₁H₂₄O₉).




Mol	Chain	Residues	Atoms			
			Total	C	H	O
5	A	1	49	21	21	7
5	B	1	49	21	21	7

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: DNA (5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3')




- Molecule 1: DNA (5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3')




- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose




- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose



- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose



- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose

Chain F:  100%

DDA1
DDA2
ERI3

4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: DNA (5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3')

Chain A:  25% 62% 12%


T1
T2
G3
G4
C5
C6
A7
A8

- Molecule 1: DNA (5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3')

Chain B:  12% 75% 12%


T11
T12
G13
G14
C15
C16
A17
A18

- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose

Chain C:  100%


2GL1
1GL2

- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose

Chain D:  100%

2GL1
1GL2

- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose

Chain E:  100%



- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose


Chain F: 100%



4.2.2 Score per residue for model 2

- Molecule 1: DNA (5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3')

Chain A: 88% 12%



- Molecule 1: DNA (5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3')

Chain B: 12% 62% 25%



- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose

Chain C: 50% 50%



- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose

Chain D: 50% 50%




- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose

Chain E: 100%



- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose

Chain F:  100%

DDA1
DDA2
ERI3

4.2.3 Score per residue for model 3

- Molecule 1: DNA (5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3')

Chain A:  38% 62%

T1
T2
G3
G4
C5
C6
A7
A8

- Molecule 1: DNA (5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3')

Chain B:  12% 50% 38%


T11
T12
G13
G14
C15
C16
A17
A18

- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose

Chain C:  50% 50%

2GL1
1GL2

- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose

Chain D:  100%

2GL1
1GL2

- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose

Chain E:  67% 33%

DDA1
DDA2
ERI3

- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose

Chain F:  67% 33%



4.2.4 Score per residue for model 4

- Molecule 1: DNA ($5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3'$)

Chain A:  75% 25%




- Molecule 1: DNA ($5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3'$)

Chain B:  50% 50%




- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose

Chain C:  100%




- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose

Chain D:  100%




- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose

Chain E:  100%



- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose

Chain F:  100%

DDA1
DDA2
ERI3


4.2.5 Score per residue for model 5

- Molecule 1: DNA (5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3')




T₁
T₂
G₃
G₄
G₅
A₇
A₈

- Molecule 1: DNA (5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3')



T₁
T₂
G₃
G₄
C₁₅
C₁₆
A₁₇
A₁₈

- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose



2GL1
1GL2

- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose



2GL1
1GL2

- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose



DDA1
DDA2
ERI3

- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose




DDA1
DDA2
ERI3

4.2.6 Score per residue for model 6


- Molecule 1: DNA (5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3')




- Molecule 1: DNA (5'-D(*TP*TP*GP*GP*CP*CP*AP*A)-3')




- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose




- Molecule 2: 2,6-dideoxy-4-O-methyl-alpha-D-galactopyranose-(1-3)-4-O-acetyl-2,6-dideoxy-beta-D-galactopyranose



- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose



- Molecule 3: 3-C-methyl-4-O-acetyl-alpha-L-Olivopyranose-(1-3)-beta-D-Olivopyranose-(1-3)-beta-D-Olivopyranose



5 Refinement protocol and experimental data overview i

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 6 were deposited, based on the following criterion: *MINIMIZED AVERAGED COORDINATES FROM 6 RUNS*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.0, 3.851

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CO, 1GL, 2GL, CPH, DDA, ERI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.41±0.04	1±1/180 (0.6± 0.5%)	2.69±0.15	18±3/276 (6.5± 1.0%)
1	B	1.40±0.06	1±0/180 (0.7± 0.3%)	2.62±0.22	16±3/276 (5.9± 0.9%)
All	All	1.41	14/2160 (0.6%)	2.66	204/3312 (6.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	1.2±0.4	1.8±1.3
1	B	0.8±0.7	2.0±1.3
All	All	12	23

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	B	11	DT	N1-C2	6.52	1.43	1.38	5	2
1	B	12	DT	N1-C2	5.53	1.42	1.38	5	6
1	A	2	DT	N1-C2	5.35	1.42	1.38	1	2
1	A	1	DT	N1-C2	5.12	1.42	1.38	3	3
1	A	7	DA	N9-C4	5.10	1.41	1.37	1	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	6	DC	P-O3'-C3'	18.35	141.72	119.70	5	4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	17	DA	P-O3'-C3'	16.82	139.89	119.70	6	6
1	B	13	DG	P-O3'-C3'	15.02	137.72	119.70	4	6
1	A	5	DC	P-O3'-C3'	14.90	137.57	119.70	1	3
1	A	3	DG	P-O3'-C3'	14.74	137.39	119.70	5	6
1	A	7	DA	P-O3'-C3'	14.49	137.08	119.70	1	5
1	B	15	DC	P-O3'-C3'	13.82	136.28	119.70	6	4
1	B	12	DT	P-O3'-C3'	13.69	136.13	119.70	2	4
1	A	4	DG	P-O5'-C5'	12.11	140.27	120.90	4	6
1	A	6	DC	C4'-C3'-C2'	-11.54	92.71	103.10	3	1
1	B	14	DG	P-O5'-C5'	11.03	138.55	120.90	6	6
1	A	6	DC	O4'-C1'-N1	10.70	115.49	108.00	1	3
1	A	3	DG	C4'-C3'-C2'	-10.28	93.84	103.10	5	3
1	B	14	DG	N3-C2-N2	10.24	127.07	119.90	6	6
1	B	16	DC	P-O3'-C3'	10.17	131.90	119.70	5	3
1	B	13	DG	C4'-C3'-C2'	-10.12	94.00	103.10	6	3
1	A	4	DG	N3-C2-N2	9.81	126.77	119.90	5	6
1	A	5	DC	O4'-C1'-N1	9.31	114.52	108.00	1	2
1	B	18	DA	P-O5'-C5'	9.17	135.57	120.90	5	2
1	A	8	DA	P-O5'-C5'	8.63	134.71	120.90	4	3
1	B	14	DG	P-O3'-C3'	8.53	129.94	119.70	3	6
1	A	6	DC	C6-N1-C2	-8.17	117.03	120.30	1	4
1	B	17	DA	P-O5'-C5'	8.16	133.96	120.90	6	1
1	B	12	DT	P-O5'-C5'	8.01	133.71	120.90	3	5
1	A	7	DA	P-O5'-C5'	7.87	133.50	120.90	4	3
1	A	2	DT	P-O3'-C3'	7.86	129.13	119.70	1	6
1	A	4	DG	P-O3'-C3'	7.76	129.01	119.70	2	6
1	A	3	DG	P-O5'-C5'	7.70	133.22	120.90	6	6
1	A	6	DC	P-O5'-C5'	7.58	133.03	120.90	5	4
1	A	5	DC	O4'-C4'-C3'	-7.42	101.53	104.50	1	1
1	B	11	DT	O4'-C1'-N1	7.29	113.10	108.00	5	1
1	B	16	DC	O4'-C1'-N1	7.17	113.02	108.00	6	1
1	A	4	DG	O4'-C4'-C3'	-6.98	101.71	104.50	6	6
1	B	15	DC	C4'-C3'-C2'	-6.89	96.90	103.10	6	5
1	B	15	DC	O4'-C1'-N1	6.78	112.75	108.00	4	3
1	B	13	DG	P-O5'-C5'	6.75	131.70	120.90	1	6
1	A	5	DC	P-O5'-C5'	6.18	130.80	120.90	1	1
1	A	3	DG	N3-C2-N2	6.11	124.18	119.90	3	1
1	B	15	DC	N1-C2-O2	6.06	122.53	118.90	2	4
1	B	15	DC	C1'-O4'-C4'	-6.05	104.05	110.10	4	1
1	B	11	DT	N3-C2-O2	-6.03	118.68	122.30	5	1
1	A	4	DG	C4'-C3'-C2'	-6.02	97.68	103.10	4	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	14	DG	C2-N3-C4	6.00	114.90	111.90	6	4
1	A	5	DC	N1-C2-O2	5.95	122.47	118.90	6	5
1	B	14	DG	O4'-C4'-C3'	-5.84	102.16	104.50	3	3
1	A	2	DT	P-O5'-C5'	5.81	130.19	120.90	1	4
1	B	15	DC	O4'-C4'-C3'	-5.81	102.18	104.50	4	1
1	B	16	DC	P-O5'-C5'	5.80	130.18	120.90	4	2
1	B	16	DC	C6-N1-C2	-5.77	117.99	120.30	6	3
1	A	5	DC	C4'-C3'-C2'	-5.76	97.91	103.10	4	3
1	B	13	DG	N3-C2-N2	5.74	123.92	119.90	3	2
1	A	6	DC	C6-N1-C1'	5.73	127.68	120.80	6	1
1	A	4	DG	C2-N3-C4	5.67	114.73	111.90	5	2
1	B	14	DG	C4'-C3'-C2'	-5.67	98.00	103.10	3	3
1	A	4	DG	N1-C2-N2	-5.62	111.15	116.20	3	5
1	B	14	DG	N1-C2-N2	-5.59	111.17	116.20	3	3
1	A	2	DT	O4'-C1'-N1	5.43	111.80	108.00	4	1
1	A	7	DA	C5'-C4'-C3'	-5.37	104.43	114.10	5	1
1	B	17	DA	C5'-C4'-C3'	-5.29	104.57	114.10	5	1
1	A	7	DA	C4'-C3'-O3'	5.24	122.81	109.70	6	1
1	A	7	DA	C3'-C2'-C1'	-5.16	96.30	102.50	6	1
1	B	15	DC	C5'-C4'-C3'	-5.15	104.82	114.10	4	1

All unique chiral outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Models (Total)
1	A	8	DA	C3',C4'	5
1	B	18	DA	C4',C3'	4
1	A	6	DC	C4'	1
1	B	11	DT	C1'	1
1	A	7	DA	C3'	1

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	6	DC	Sidechain	4
1	B	11	DT	Sidechain	3
1	B	14	DG	Sidechain	3
1	A	2	DT	Sidechain	3
1	A	8	DA	Sidechain	2
1	B	12	DT	Sidechain	2
1	A	1	DT	Sidechain	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	B	13	DG	Sidechain	1
1	B	18	DA	Sidechain	1
1	A	4	DG	Sidechain	1
1	B	16	DC	Sidechain	1
1	B	17	DA	Sidechain	1

6.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	161	92	92	1±1
1	A	161	92	92	1±1
2	C	23	25	24	0±0
2	D	23	25	23	0±0
3	E	32	35	33	0±0
3	F	32	35	33	0±0
5	B	28	21	19	1±1
5	A	28	21	19	0±0
All	All	2934	2076	2021	12

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:5:DC:H5'	2:C:1:2GL:H1	0.52	1.82	2	2
1:B:16:DC:H5''	5:B:23:CPH:HMEB	0.50	1.82	1	1
1:B:15:DC:H4'	5:B:23:CPH:C5	0.50	2.37	5	1
5:A:33:CPH:HC7	1:B:16:DC:O2	0.49	2.07	4	1
1:B:15:DC:H4'	5:B:23:CPH:C5A	0.48	2.38	4	2
1:B:15:DC:H5'	2:D:1:2GL:H1	0.47	1.86	2	1
1:B:15:DC:H2''	1:B:16:DC:H5''	0.47	1.86	3	1
1:A:5:DC:H2''	1:A:6:DC:H5''	0.43	1.89	6	1
5:A:33:CPH:HMEB	3:F:1:DDA:H62	0.41	1.91	3	1
1:A:3:DG:H21	3:E:3:ERI:H63	0.40	1.76	3	1

6.3 Torsion angles [\(i\)](#)

6.3.1 Protein backbone [\(i\)](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [\(i\)](#)

There are no protein molecules in this entry.

6.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [\(i\)](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	2GL	C	1	2	13,13,13	1.17±0.04	1±0 (6±2%)
2	1GL	C	2	2	10,10,11	1.05±0.06	1±0 (10±0%)
2	2GL	D	1	2	13,13,13	1.29±0.27	1±1 (7±7%)
2	1GL	D	2	2	10,10,11	1.05±0.06	1±0 (10±0%)
3	DDA	E	1	3	10,10,10	1.51±0.20	2±1 (15±7%)
3	DDA	E	2	3	9,9,10	0.82±0.23	0±0 (1±4%)
3	ERI	E	3	3	13,13,14	0.75±0.05	0±0 (0±0%)
3	DDA	F	1	3	10,10,10	1.41±0.19	1±1 (13±7%)
3	DDA	F	2	3	9,9,10	0.70±0.10	0±0 (0±0%)
3	ERI	F	3	3	13,13,14	0.72±0.05	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	2GL	C	1	2	15,18,18	1.79±0.08	6±1 (38±5%)
2	1GL	C	2	2	11,13,15	1.14±0.10	1±0 (12±4%)
2	2GL	D	1	2	15,18,18	1.91±0.33	5±1 (34±5%)
2	1GL	D	2	2	11,13,15	1.17±0.08	2±0 (15±4%)
3	DDA	E	1	3	12,14,14	1.92±0.24	3±2 (25±12%)
3	DDA	E	2	3	12,12,14	1.45±0.12	2±0 (18±3%)
3	ERI	E	3	3	17,19,21	1.38±0.08	2±1 (12±6%)
3	DDA	F	1	3	12,14,14	1.64±0.09	2±1 (20±6%)
3	DDA	F	2	3	12,12,14	1.52±0.13	2±0 (19±3%)
3	ERI	F	3	3	17,19,21	1.37±0.03	2±0 (14±2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2GL	C	1	2	-	0±0,4,20,20	0±0,1,1,1
2	1GL	C	2	2	-	0±0,2,16,18	0±0,1,1,1
2	2GL	D	1	2	-	0±0,4,20,20	0±0,1,1,1
2	1GL	D	2	2	-	0±0,2,16,18	0±0,1,1,1
3	DDA	E	1	3	-	-	0±0,1,1,1
3	DDA	E	2	3	-	-	0±0,1,1,1
3	ERI	E	3	3	-	0±0,4,21,23	0±0,1,1,1
3	DDA	F	1	3	-	-	1±0,1,1,1
3	DDA	F	2	3	-	-	0±0,1,1,1
3	ERI	F	3	3	-	0±0,4,21,23	0±0,1,1,1

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
3	E	1	DDA	C2-C3	4.52	1.44	1.52	2	5
3	F	1	DDA	C2-C3	4.45	1.45	1.52	4	5
2	D	1	2GL	O5-C1	3.46	1.37	1.44	6	1
3	E	1	DDA	C3-C4	3.22	1.57	1.52	6	2
2	D	1	2GL	C3-C4	3.15	1.45	1.52	6	1
3	E	1	DDA	C4-C5	2.97	1.59	1.52	5	1
3	F	1	DDA	O1-C1	2.72	1.46	1.39	3	1
3	E	2	DDA	C2-C3	2.63	1.48	1.52	3	1
2	D	1	2GL	C2-C3	2.46	1.48	1.52	6	1
2	C	2	1GL	O4-C4	2.33	1.48	1.42	5	6
2	D	2	1GL	O4-C4	2.30	1.48	1.42	1	6
3	F	1	DDA	C3-C4	2.17	1.55	1.52	6	2
3	E	1	DDA	C2-C1	2.16	1.45	1.51	3	1
2	C	1	2GL	O4-C4	2.15	1.48	1.44	2	5
2	D	1	2GL	O4-C4	2.14	1.48	1.44	1	3

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
3	E	1	DDA	O3-C3-C2	5.94	95.20	109.94	3	6
2	D	1	2GL	O5-C5-C6	5.49	118.56	106.70	6	3
3	F	1	DDA	O3-C3-C2	4.74	98.18	109.94	6	6
3	E	3	ERI	O5-C1-C2	4.55	104.97	111.38	5	6
2	D	1	2GL	C6-C5-C4	4.54	120.33	113.41	6	1
3	E	1	DDA	O5-C5-C4	4.26	117.16	109.52	5	2
2	D	1	2GL	O4-CO4-CME	4.01	103.71	111.09	6	6
3	F	3	ERI	O5-C1-C2	3.88	105.92	111.38	6	6
3	F	1	DDA	O1-C1-O5	3.78	117.14	108.36	3	4
3	E	2	DDA	O3-C3-C4	3.67	102.79	110.14	5	5
2	D	1	2GL	C4-O4-CO4	3.52	123.16	117.72	5	4
3	E	1	DDA	C2-C3-C4	3.51	105.58	110.69	5	2
3	F	2	DDA	O3-C3-C4	3.36	103.41	110.14	5	6
2	C	1	2GL	O4-CO4-CME	3.35	104.93	111.09	6	6
2	C	1	2GL	C4-O4-CO4	3.28	122.80	117.72	4	6
3	E	2	DDA	O5-C5-C4	3.11	103.95	109.52	5	6
3	F	2	DDA	O5-C5-C4	3.10	103.96	109.52	3	6
2	D	1	2GL	O4-C4-C3	3.06	113.36	107.99	6	5
3	E	1	DDA	O1-C1-C2	3.00	96.82	111.42	2	2
3	F	1	DDA	O1-C1-C2	2.94	97.10	111.42	4	2
3	F	3	ERI	C6-C5-C4	2.91	107.72	113.52	1	2
3	F	1	DDA	O5-C5-C6	2.78	112.69	106.70	1	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	C	1	2GL	O3-C3-C2	2.74	103.14	109.94	6	6
3	F	1	DDA	C1-C2-C3	2.74	104.61	110.91	5	1
2	D	1	2GL	O5-C5-C4	2.70	114.16	109.13	6	5
3	E	3	ERI	CC3-C3-C2	2.68	107.40	111.03	3	1
2	C	1	2GL	O5-C5-C4	2.66	104.17	109.13	2	6
2	C	1	2GL	C6-C5-C4	2.64	109.39	113.41	5	2
3	F	2	DDA	O5-C1-C2	2.60	105.75	110.73	1	1
3	E	1	DDA	O5-C5-C6	2.58	101.13	106.70	4	1
2	D	1	2GL	O3-C3-C2	2.57	103.56	109.94	3	5
2	C	1	2GL	O4-C4-C3	2.56	112.48	107.99	6	4
2	C	2	1GL	CME-O4-C4	2.43	108.14	114.52	4	6
3	E	3	ERI	C6-C5-C4	2.43	108.69	113.52	1	1
3	E	2	DDA	O4-C4-C3	2.40	114.59	109.99	3	1
2	D	2	1GL	CME-O4-C4	2.39	108.26	114.52	6	6
3	E	1	DDA	C3-C4-C5	2.37	102.88	110.02	6	2
2	D	2	1GL	O3-C3-C4	2.34	115.62	110.07	5	4
3	E	2	DDA	O3-C3-C2	2.31	104.42	109.96	3	1
3	F	3	ERI	O4-C4-C3	2.28	114.22	108.04	1	6
2	D	1	2GL	C1-C2-C3	2.24	116.06	110.91	6	1
3	E	1	DDA	C6-C5-C4	2.24	117.21	113.07	4	1
2	C	2	1GL	O3-C3-C4	2.21	115.31	110.07	5	2
3	F	1	DDA	O4-C4-C3	2.20	114.21	109.99	3	1
2	D	1	2GL	O4-CO4-OC4	2.19	127.32	122.96	5	1
2	C	1	2GL	O4-CO4-OC4	2.18	127.29	122.96	1	4
3	E	3	ERI	O4-C4-C3	2.17	113.92	108.04	6	3
3	E	3	ERI	O4-CO4-CME	2.17	107.09	111.09	3	1
3	E	3	ERI	C1-O5-C5	2.10	104.22	110.95	3	1
3	E	1	DDA	C1-C2-C3	2.10	106.08	110.91	5	1
3	E	1	DDA	O4-C4-C3	2.09	114.00	109.99	3	1
2	C	1	2GL	O4-C4-C5	2.08	111.98	107.83	2	1
3	F	3	ERI	O4-CO4-OC4	2.06	127.06	122.96	3	1
3	F	2	DDA	C2-C3-C4	2.00	108.86	111.16	6	1

All unique chiral outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.


Mol	Chain	Res	Type	Atoms	Models (Total)
2	D	1	2GL	C5	2
3	E	1	DDA	C4	2
3	F	1	DDA	C4	2



There are no torsion outliers.


All unique ring outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
3	F	1	DDA	C1-C2-C3-C4-C5-O5	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







6.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
5	CPH	B	23	4	29,30,32	1.39 ± 0.16	4 ± 2 ($13 \pm 5\%$)
5	CPH	A	33	4	29,30,32	1.51 ± 0.19	5 ± 1 ($16 \pm 4\%$)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Counts	Bond angles		
						RMSZ	#Z>2	
5	CPH	B	23	4	37,45,49	2.36±0.23	11±2 (29±4%)	
5	CPH	A	33	4	37,45,49	2.94±0.86	10±2 (28±5%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CPH	A	33	4	-	0±0,18,30,34	0±0,3,3,3
5	CPH	B	23	4	-	0±0,18,30,34	0±0,3,3,3

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
5	A	33	CPH	C4-C3	5.35	1.44	1.52	6	2
5	A	33	CPH	C2-C1	5.14	1.58	1.50	1	5
5	B	23	CPH	C2-C1	4.85	1.58	1.50	3	5
5	B	23	CPH	C5-C5A	3.67	1.33	1.41	5	5
5	A	33	CPH	C9A-C9	3.65	1.45	1.39	6	4
5	A	33	CPH	C5-C5A	3.41	1.33	1.41	5	4
5	B	23	CPH	C9A-C9	3.38	1.44	1.39	6	4
5	A	33	CPH	C5-C6	2.84	1.42	1.36	3	4
5	B	23	CPH	C4-C3	2.71	1.48	1.52	6	2
5	A	33	CPH	C9A-C1	2.57	1.52	1.46	3	4
5	A	33	CPH	C10-C5A	2.53	1.36	1.42	6	1
5	B	23	CPH	C9A-C1	2.46	1.52	1.46	6	2
5	A	33	CPH	C3'-C2'	2.45	1.60	1.52	3	2
5	B	23	CPH	C5-C6	2.41	1.41	1.36	5	4
5	B	23	CPH	C10-C5A	2.17	1.37	1.42	3	1
5	A	33	CPH	C4-C4A	2.16	1.47	1.51	6	1
5	A	33	CPH	O8-C8	2.12	1.42	1.35	4	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
5	A	33	CPH	C9-C9A-C1	20.54	112.03	119.50	4	4
5	A	33	CPH	C9-C9A-C4A	10.83	126.38	119.77	4	5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
5	A	33	CPH	C4A-C9A-C1	9.08	107.56	119.53	5	2
5	A	33	CPH	C4-C4A-C10	7.94	107.38	121.05	6	4
5	B	23	CPH	C4A-C9A-C1	7.86	109.18	119.53	4	3
5	A	33	CPH	C4-C4A-C9A	6.91	129.81	118.03	6	5
5	B	23	CPH	C9-C9A-C4A	6.56	123.78	119.77	4	6
5	B	23	CPH	C2-C1-C9A	6.42	105.90	117.12	4	5
5	B	23	CPH	C9A-C9-C8A	6.36	116.06	121.09	4	6
5	A	33	CPH	C9A-C9-C8A	6.32	116.09	121.09	5	6
5	A	33	CPH	O1-C1-C2	6.12	129.40	120.79	5	4
5	B	23	CPH	C4-C4A-C10	5.61	111.40	121.05	3	6
5	B	23	CPH	O1-C1-C2	5.56	128.60	120.79	4	3
5	B	23	CPH	C4-C4A-C9A	5.37	127.18	118.03	3	6
5	B	23	CPH	C9-C9A-C1	5.14	121.37	119.50	6	5
5	A	33	CPH	C2-C1-C9A	5.08	108.25	117.12	6	5
5	B	23	CPH	O1-C1-C9A	4.94	114.56	122.38	5	5
5	A	33	CPH	O1-C1-C9A	4.94	114.57	122.38	3	6
5	B	23	CPH	O9-C9-C8A	4.52	108.11	119.00	6	1
5	A	33	CPH	C5-C5A-C10	4.45	114.16	122.02	6	5
5	A	33	CPH	O3'-C3'-C2'	4.21	118.18	110.85	4	4
5	B	23	CPH	C5-C5A-C10	3.74	115.41	122.02	4	6
5	A	33	CPH	C3-C2-C1	3.70	104.83	113.37	3	1
5	B	23	CPH	C5'-C4'-C3'	3.68	102.65	111.55	6	2
5	B	23	CPH	C4A-C4-C3	3.64	103.84	112.23	5	1
5	B	23	CPH	O2'-C2'-C1'	3.50	129.28	119.87	5	1
5	B	23	CPH	O3'-C3'-C2'	3.42	116.80	110.85	2	2
5	A	33	CPH	C10-C5A-C8A	3.05	124.78	118.38	5	3
5	B	23	CPH	C10-C5A-C8A	2.91	124.50	118.38	5	4
5	A	33	CPH	C5'-C4'-C3'	2.91	104.51	111.55	6	1
5	A	33	CPH	O9-C9-C8A	2.83	112.19	119.00	6	4
5	B	23	CPH	CME-O1'-C1'	2.78	106.47	114.00	2	2
5	B	23	CPH	C4A-C10-C5A	2.51	116.83	121.28	5	2
5	A	33	CPH	C4A-C4-C3	2.33	106.85	112.23	4	1
5	A	33	CPH	C10-C4A-C9A	2.32	122.81	118.63	6	1
5	A	33	CPH	O9-C9-C9A	2.12	125.53	121.29	5	1
5	A	33	CPH	C4A-C10-C5A	2.08	117.58	121.28	6	1

All unique chiral outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Models (Total)
5	B	23	CPH	C1'	3
5	A	33	CPH	C4'	2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms	Models (Total)
5	A	33	CPH	C1'	2


All unique torsion outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
5	A	33	CPH	C3-C1'-O1'-CME	1

There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





6.7 Other polymers [\(i\)](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

7 Chemical shift validation [\(i\)](#)

No chemical shift data were provided