

Full wwPDB X-ray Structure Validation Report (i)

Nov 21, 2023 – 02:58 AM JST

PDB ID : 7EL6

Title: Structure of SMCR8 bound FEM1B

Authors : Zhao, S.; Xu, C.

Deposited on : 2021-04-08

Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467 Xtriage (Phenix) : 1.13

EDS : 2.36

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	357	82%	16%	•
1	В	357	78%	16%	5%



2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 5229 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Protein fem-1 homolog B,Guanine nucleotide exchange protein SMCR8.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	348	Total 2665	C 1684	N 485	O 485	S 11	0	3	0
1	В	338	Total 2564	C 1619	N 455	O 478	S 12	0	1	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	338	GLY	-	linker	UNP Q9UK73
A	339	GLY	-	linker	UNP Q9UK73
A	340	GLY	-	linker	UNP Q9UK73
A	341	SER	-	linker	UNP Q9UK73
A	342	GLY	-	linker	UNP Q9UK73
A	343	GLY	-	linker	UNP Q9UK73
A	344	GLY	-	linker	UNP Q9UK73
A	345	SER	-	linker	UNP Q9UK73
A	346	GLY	-	linker	UNP Q9UK73
A	347	SER	-	linker	UNP Q9UK73
В	338	GLY	-	linker	UNP Q9UK73
В	339	GLY	-	linker	UNP Q9UK73
В	340	GLY	-	linker	UNP Q9UK73
В	341	SER	-	linker	UNP Q9UK73
В	342	GLY	-	linker	UNP Q9UK73
В	343	GLY	-	linker	UNP Q9UK73
В	344	GLY	-	linker	UNP Q9UK73
В	345	SER	-	linker	UNP Q9UK73
В	346	GLY	-	linker	UNP Q9UK73
В	347	SER	-	linker	UNP Q9UK73



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Protein fem-1 homolog B, Guanine nucleotide exchange protein SMCR8





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	83.31Å 91.50Å 112.15Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.38 - 2.80	Depositor
Resolution (A)	37.38 - 2.80	EDS
% Data completeness	96.8 (37.38-2.80)	Depositor
(in resolution range)	96.8 (37.38-2.80)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	13.24 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
D D.	0.203 , 0.266	Depositor
R, R_{free}	0.203 , 0.265	DCC
R_{free} test set	1998 reflections (9.52%)	wwPDB-VP
Wilson B-factor (Å ²)	55.2	Xtriage
Anisotropy	0.868	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32, 39.1	EDS
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5229	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	$\mathbf{lengths}$	Bond angles		
WIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.30	0/2719	0.45	0/3687	
1	В	0.31	0/2605	0.48	0/3529	
All	All	0.31	0/5324	0.47	0/7216	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	2665	0	2633	30	0
1	В	2564	0	2505	33	0
All	All	5229	0	5138	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-1 Atom-2		Clash overlap (Å)	
1:B:239:LEU:HD22	1:B:249:ARG:HG3	1.73	0.71	
1:A:5:ALA:HA	1:A:8:VAL:HG12	1.74	0.70	

Continued on next page...



 $Continued\ from\ previous\ page...$

Continued from pred		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	$\text{overlap } (\mathring{\mathbf{A}})$
1:B:307:GLU:OE1	1:B:334:ARG:NH1	2.23	0.66
1:A:305:ARG:NH2	1:A:327:GLU:OE1	2.30	0.64
1:A:29:GLU:HA	1:A:32:ILE:HB	1.80	0.63
1:A:268:ASP:OD2	1:A:271:LYS:HD2	1.98	0.62
1:A:170:VAL:HG11	1:A:202:ILE:HG23	1.85	0.56
1:B:221:THR:O	1:B:225:VAL:HG23	2.06	0.55
1:B:133:ARG:O	1:B:137:VAL:HG23	2.07	0.54
1:B:195:ALA:HB1	1:B:226:ALA:HB2	1.88	0.54
1:A:118:THR:OG1	1:A:120:THR:OG1	2.21	0.54
1:A:195:ALA:HB1	1:A:226:ALA:HB2	1.90	0.53
1:A:20:LEU:HD21	1:A:65:LEU:HD21	1.93	0.51
1:B:137:VAL:O	1:B:141:VAL:HG23	2.11	0.51
1:A:29:GLU:HB3	1:A:33[B]:ARG:HH21	1.76	0.51
1:B:22:ALA:HA	1:B:25:LEU:HD23	1.94	0.50
1:B:226:ALA:HB1	1:B:234:VAL:HG13	1.95	0.49
1:B:198:GLY:CA	1:B:231:LYS:HG2	2.42	0.49
1:B:200:ILE:HD11	1:B:237:LEU:HD13	1.95	0.49
1:A:100:HIS:O	1:A:104:VAL:HG23	2.13	0.48
1:B:254:GLU:OE2	1:B:283:ARG:NH1	2.47	0.48
1:A:249:ARG:NH1	1:A:282:GLU:OE1	2.46	0.47
1:A:137:VAL:O	1:A:141:VAL:HG12	2.14	0.47
1:A:133:ARG:O	1:A:137:VAL:HG23	2.14	0.47
1:B:102:GLU:OE1	1:B:105:LYS:HE2	2.14	0.47
1:B:192:HIS:NE2	1:B:222:PRO:HD3	2.30	0.47
1:A:65:LEU:O	1:A:71:VAL:HG23	2.15	0.47
1:B:260:PHE:CE1	1:B:271:LYS:HE3	2.50	0.47
1:B:192:HIS:CE1	1:B:222:PRO:HD3	2.50	0.46
1:B:198:GLY:HA3	1:B:231:LYS:HG2	1.97	0.46
1:B:257:GLY:O	1:B:272:THR:HG23	2.15	0.46
1:B:279:ALA:O	1:B:283:ARG:HG3	2.16	0.46
1:A:260:PHE:O	1:A:267:TYR:HB2	2.15	0.46
1:B:20:LEU:HD23	1:B:64:LEU:HD23	1.98	0.46
1:B:16:LYS:HB3	1:B:19:THR:CG2	2.46	0.46
1:A:277:TYR:OH	1:A:311:PRO:HB3	2.17	0.45
1:B:233:ASP:N	1:B:233:ASP:OD1	2.48	0.45
1:B:168:ASP:N	1:B:168:ASP:OD1	2.48	0.45
1:A:320:ASP:HB3	1:A:323:ALA:HB3	1.99	0.44
1:B:280:MET:HE1	1:B:311:PRO:HA	1.98	0.44
1:A:63:ARG:NH2	1:A:67:GLU:OE1	2.50	0.44
1:B:247:ARG:HA	1:B:250:ILE:HG12	1.99	0.44
1:A:293:LYS:HE2	1:A:307:GLU:OE1	2.18	0.44

Continued on next page...



Continued from previous page...

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:A:296:LEU:HD23	1:A:296:LEU:HA	1.87	0.43
1:B:162:ALA:HA	1:B:202:ILE:HD13	2.00	0.43
1:A:174:LEU:HD13	1:A:209:TRP:CD1	2.54	0.43
1:B:232:ALA:HA	1:B:235:VAL:HG12	2.01	0.42
1:B:135:ASP:OD1	1:B:135:ASP:N	2.47	0.42
1:A:283:ARG:HD3	1:A:291:LEU:O	2.20	0.42
1:A:171:ARG:HG3	1:A:205:GLU:OE2	2.20	0.42
1:B:263:ASP:O	1:B:267:TYR:HB3	2.20	0.42
1:B:105:LYS:HB3	1:B:139:TYR:CZ	2.55	0.41
1:A:239:LEU:HD23	1:A:239:LEU:HA	1.89	0.41
1:A:284:PHE:CD2	1:A:290:ILE:HD13	2.54	0.41
1:A:293:LYS:O	1:A:309:ARG:NH2	2.53	0.41
1:B:235:VAL:HG11	1:B:256:LEU:HD22	2.02	0.41
1:A:299:ILE:HG13	1:A:300:HIS:CE1	2.55	0.41
1:B:141:VAL:HG13	1:B:147:ILE:HD11	2.02	0.41
1:B:264:ARG:HB2	1:B:325:HIS:ND1	2.36	0.41
1:A:257:GLY:O	1:A:272:THR:HG23	2.22	0.40
1:B:63:ARG:HB2	1:B:106:LEU:HD21	2.04	0.40
1:A:138:LYS:HG2	1:A:172:TYR:CZ	2.56	0.40
1:A:299:ILE:HG13	1:A:300:HIS:ND1	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	347/357 (97%)	333 (96%)	14 (4%)	0	100	100
1	В	331/357~(93%)	317 (96%)	14 (4%)	0	100	100
All	All	$678/714 \ (95\%)$	650 (96%)	28 (4%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	$263/285 \ (92\%)$	256 (97%)	7 (3%)	44 78
1	В	250/285~(88%)	244 (98%)	6 (2%)	49 81
All	All	513/570 (90%)	500 (98%)	13 (2%)	49 80

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33[A]	ARG
1	A	33[B]	ARG
1	A	55	ARG
1	A	64	LEU
1	A	246	ARG
1	A	255	LEU
1	A	349	LYS
1	В	55	ARG
1	В	105	LYS
1	В	171	ARG
1	В	244	CYS
1	В	246	ARG
1	В	357	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(Å^2)$	Q < 0.9
1	A	348/357 (97%)	-0.34	0 100 100	34, 56, 90, 140	0
1	В	338/357 (94%)	-0.24	2 (0%) 89 86	34, 63, 107, 172	0
All	All	686/714 (96%)	-0.29	2 (0%) 94 93	34, 59, 101, 172	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	241	HIS	2.6
1	В	26	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

