



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 02:15 PM BST

PDB ID : 4ENO
Title : Crystal structure of oxidized human nm23-H1
Authors : Kim, M.-S.; Shin, D.-H.
Deposited on : 2012-04-13
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2437 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoside diphosphate kinase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	151	Total 1196	C 765	N 206	O 217	S 8	0	0	0
1	B	149	Total 1176	C 750	N 204	O 214	S 8	0	0	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	Total 5	O 4	P 1	0	0
2	B	1	Total 5	O 4	P 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	25	Total 25	O 25	0	0
3	B	30	Total 30	O 30	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nucleoside diphosphate kinase A



- Molecule 1: Nucleoside diphosphate kinase A



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	106.77Å 106.77Å 106.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.75 – 2.80 47.75 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.75-2.80) 99.9 (47.75-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.14 (at 2.81Å)	Xtrriage
Refinement program	CNS 1.3, PHENIX 1.8.1_1168	Depositor
R, R_{free}	0.182 , 0.256 0.195 , 0.270	Depositor DCC
R_{free} test set	1021 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å ²)	34.9	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 35.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.057 for l,-k,h	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2437	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/1223	0.64	0/1649
1	B	0.46	0/1202	0.66	1/1620 (0.1%)
All	All	0.47	0/2425	0.65	1/3269 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	81	LEU	CA-CB-CG	5.12	127.09	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1196	0	1188	14	0
1	B	1176	0	1168	19	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	25	0	0	2	0
3	B	30	0	0	1	0
All	All	2437	0	2356	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:109:CYS:HB3	1:B:116:ILE:HG12	1.56	0.87
1:A:109:CYS:HB3	1:A:116:ILE:HG12	1.55	0.87
1:B:88:ARG:HD2	1:B:121:ASP:HA	1.71	0.73
1:A:38:LEU:HB3	1:B:40:PHE:HB2	1.73	0.69
1:B:149:TRP:O	3:B:329:HOH:O	2.13	0.66
1:B:141:ASP:HB3	1:B:146:ALA:HB2	1.86	0.58
1:A:72:PRO:HB3	1:B:140:VAL:HG11	1.87	0.56
1:A:17:GLN:NE2	3:A:314:HOH:O	2.37	0.56
1:A:96:PRO:HG2	1:A:112:VAL:HA	1.88	0.55
1:B:6:ARG:NH2	1:B:139:LEU:HD13	2.23	0.54
1:A:140:VAL:HG11	1:B:72:PRO:HB3	1.91	0.53
1:A:39:LYS:HG2	1:A:75:ALA:HB3	1.90	0.53
1:A:88:ARG:HD2	1:A:121:ASP:HA	1.93	0.51
1:B:116:ILE:HG22	1:B:117:ILE:HB	1.93	0.50
1:B:7:THR:HB	1:B:84:VAL:HG22	1.93	0.50
1:B:67:TYR:CD1	1:B:114:ARG:HD2	2.47	0.49
1:A:40:PHE:HB2	1:B:38:LEU:HB3	1.94	0.48
1:A:1:MET:SD	1:A:3:ASN:HB2	2.53	0.48
1:B:35:LEU:HD13	1:B:78:TRP:CZ2	2.49	0.47
1:A:18:ARG:HH11	1:A:116:ILE:HD11	1.79	0.47
1:B:94:THR:O	1:B:96:PRO:HD3	2.15	0.46
1:A:124:GLU:H	1:A:124:GLU:CD	2.19	0.46
1:A:43:ALA:HB1	1:A:47:LEU:HD23	1.99	0.44
1:B:142:TYR:N	1:B:142:TYR:CD1	2.86	0.43
1:B:56:LYS:HA	1:B:61:PHE:CD1	2.53	0.42
1:B:137:GLU:N	1:B:137:GLU:OE1	2.52	0.42
1:B:3:ASN:N	1:B:3:ASN:OD1	2.52	0.42
1:A:13:PRO:HD3	1:A:73:VAL:HG12	2.02	0.41
3:A:314:HOH:O	1:B:34:ARG:NH2	2.53	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/152 (98%)	141 (95%)	7 (5%)	1 (1%)	22	53
1	B	147/152 (97%)	140 (95%)	7 (5%)	0	100	100
All	All	296/304 (97%)	281 (95%)	14 (5%)	1 (0%)	41	72

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	142	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	128/129 (99%)	122 (95%)	6 (5%)	26	59
1	B	126/129 (98%)	115 (91%)	11 (9%)	10	30
All	All	254/258 (98%)	237 (93%)	17 (7%)	16	43

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	ARG
1	A	55	LEU
1	A	81	LEU
1	A	85	LYS
1	A	93	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	112	VAL
1	B	3	ASN
1	B	34	ARG
1	B	57	ASP
1	B	64	LEU
1	B	81	LEU
1	B	93	GLU
1	B	94	THR
1	B	114	ARG
1	B	143	THR
1	B	145	CYS
1	B	149	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	A	201	-	4,4,4	0.94	0	6,6,6	0.49	0
2	PO4	B	201	-	4,4,4	0.87	0	6,6,6	0.58	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	151/152 (99%)	-0.35	1 (0%) 87 84	9, 24, 53, 68	0
1	B	149/152 (98%)	-0.22	7 (4%) 31 22	10, 23, 61, 73	0
All	All	300/304 (98%)	-0.29	8 (2%) 54 44	9, 23, 60, 73	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	3.7
1	B	2	ALA	3.2
1	B	148	ASN	3.1
1	B	142	TYR	3.0
1	B	94	THR	2.9
1	B	149	TRP	2.7
1	A	1	MET	2.7
1	B	96	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	A	201	5/5	0.85	0.21	31,35,53,65	0
2	PO4	B	201	5/5	0.94	0.14	25,25,54,57	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.