

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 16, 2023 – 10:44 AM EDT

PDB ID : 8ERJ

Title: Crystal structure of Fub7 in complex with E-2-aminocrotonate

 $\begin{array}{cccc} \text{Authors} & : & \text{Hai, Y.} \\ \text{Deposited on} & : & 2022\text{-}10\text{-}12 \end{array}$ 

Resolution : 2.16 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
$R_{free}$	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	450	86%	10% 5%
1	В	450	90%	• • 5%
1	С	450	86%	8% • 5%
1	D	450	90%	5% 5%
1	Е	450	89%	6% 5%



	J	1	I = J	
$\mathbf{Mol}$	Chain	Length	Quality of chain	
1	F	450	89%	6% 5%
1	G	450	88%	5% 5%
1	Н	450	89%	5% • 5%



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 27599 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Sulfhydrylase FUB7.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace			
1	A	428	Total	С	N	О	S	0	0	0
1	A	420	3292	2091	567	627	7	0	0	
1	В	428	Total	С	N	О	S	0	1	0
1	Б	420	3300	2096	570	627	7	0		
1	С	428	Total	С	N	О	S	0	1	0
1		420	3300	2096	570	627	7		1	
1	D	428	Total	С	N	О	S	0	1	0
1	D	420	3300	2096	570	627	7			
1	Е	428	Total	С	N	О	S	0	1	0
1	12	420	3300	2096	570	627	7		1	
1	F	428	Total	С	N	О	S	0	1	0
1	I.	420	3300	2096	570	627	7		1	
1	G	427	Total	С	Ν	О	S	0	0	0
1	G	421	3285	2086	566	626	7		U	
1	Н	497	Total	С	N	О	S	0	0	0
1	1   H	427	3285	2086	566	626	7		U	U

There are 144 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-16	MET	-	initiating methionine	UNP S0DUX5
A	-15	GLY	-	expression tag	UNP S0DUX5
A	-14	SER	-	expression tag	UNP S0DUX5
A	-13	SER	-	expression tag	UNP S0DUX5
A	-12	HIS	-	expression tag	UNP S0DUX5
A	-11	HIS	-	expression tag	UNP S0DUX5
A	-10	HIS	-	expression tag	UNP S0DUX5
A	-9	HIS	ı	expression tag	UNP S0DUX5
A	-8	HIS	-	expression tag	UNP S0DUX5
A	-7	HIS	-	expression tag	UNP S0DUX5
A	-6	GLU		expression tag	UNP S0DUX5
A	-5	ASN	-	expression tag	UNP S0DUX5
A	-4	LEU	-	expression tag	UNP S0DUX5



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	TYR	-	expression tag	UNP S0DUX5
A	-2	PHE	-	expression tag	UNP S0DUX5
A	-1	GLN	-	expression tag	UNP S0DUX5
A	0	SER	-	expression tag	UNP S0DUX5
A	1	ASN	-	expression tag	UNP S0DUX5
В	-16	MET	ı	initiating methionine	UNP S0DUX5
В	-15	GLY	-	expression tag	UNP S0DUX5
В	-14	SER	-	expression tag	UNP S0DUX5
В	-13	SER	-	expression tag	UNP S0DUX5
В	-12	HIS	-	expression tag	UNP S0DUX5
В	-11	HIS	-	expression tag	UNP S0DUX5
В	-10	HIS	-	expression tag	UNP S0DUX5
В	-9	HIS	-	expression tag	UNP S0DUX5
В	-8	HIS	-	expression tag	UNP S0DUX5
В	-7	HIS	-	expression tag	UNP S0DUX5
В	-6	GLU	-	expression tag	UNP S0DUX5
В	-5	ASN	-	expression tag	UNP S0DUX5
В	-4	LEU	-	expression tag	UNP S0DUX5
В	-3	TYR	-	expression tag	UNP S0DUX5
В	-2	PHE	-	expression tag	UNP S0DUX5
В	-1	GLN	-	expression tag	UNP S0DUX5
В	0	SER	-	expression tag	UNP S0DUX5
В	1	ASN	-	expression tag	UNP S0DUX5
С	-16	MET	-	initiating methionine	UNP S0DUX5
С	-15	GLY	-	expression tag	UNP S0DUX5
С	-14	SER	-	expression tag	UNP S0DUX5
С	-13	SER	-	expression tag	UNP S0DUX5
С	-12	HIS	-	expression tag	UNP S0DUX5
С	-11	HIS	-	expression tag	UNP S0DUX5
С	-10	HIS	ı	expression tag	UNP S0DUX5
С	-9	HIS	ı	expression tag	UNP S0DUX5
С	-8	HIS	-	expression tag	UNP S0DUX5
С	-7	HIS	ı	expression tag	UNP S0DUX5
С	-6	GLU	<u> </u>	expression tag	UNP S0DUX5
С	-5	ASN	-	expression tag	UNP S0DUX5
С	-4	LEU	-	expression tag	UNP S0DUX5
С	-3	TYR		expression tag	UNP S0DUX5
С	-2	PHE	-	expression tag	UNP S0DUX5
С	-1	GLN		expression tag	UNP S0DUX5
С	0	SER	-	expression tag	UNP S0DUX5
С	1	ASN	-	expression tag	UNP S0DUX5
D	-16	MET	-	initiating methionine	UNP S0DUX5



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-15	GLY	-	expression tag	UNP S0DUX5
D	-14	SER	-	expression tag	UNP S0DUX5
D	-13	SER	-	expression tag	UNP S0DUX5
D	-12	HIS	_	expression tag	UNP S0DUX5
D	-11	HIS	-	expression tag	UNP S0DUX5
D	-10	HIS	-	expression tag	UNP S0DUX5
D	-9	HIS	-	expression tag	UNP S0DUX5
D	-8	HIS	-	expression tag	UNP S0DUX5
D	-7	HIS	-	expression tag	UNP S0DUX5
D	-6	GLU	-	expression tag	UNP S0DUX5
D	-5	ASN	_	expression tag	UNP S0DUX5
D	-4	LEU	-	expression tag	UNP S0DUX5
D	-3	TYR	-	expression tag	UNP S0DUX5
D	-2	PHE	-	expression tag	UNP S0DUX5
D	-1	GLN	-	expression tag	UNP S0DUX5
D	0	SER	-	expression tag	UNP S0DUX5
D	1	ASN	-	expression tag	UNP S0DUX5
Е	-16	MET	-	initiating methionine	UNP S0DUX5
Е	-15	GLY	-	expression tag	UNP S0DUX5
Е	-14	SER	-	expression tag	UNP S0DUX5
Е	-13	SER	-	expression tag	UNP S0DUX5
Е	-12	HIS	-	expression tag	UNP S0DUX5
Е	-11	HIS	-	expression tag	UNP S0DUX5
Е	-10	HIS	-	expression tag	UNP S0DUX5
Е	-9	HIS	-	expression tag	UNP S0DUX5
Е	-8	HIS	-	expression tag	UNP S0DUX5
Е	-7	HIS	-	expression tag	UNP S0DUX5
Е	-6	GLU	-	expression tag	UNP S0DUX5
Е	-5	ASN	-	expression tag	UNP S0DUX5
Е	-4	LEU	-	expression tag	UNP S0DUX5
Е	-3	TYR	-	expression tag	UNP S0DUX5
Е	-2	PHE	-	expression tag	UNP S0DUX5
Е	-1	GLN	-	expression tag	UNP S0DUX5
Е	0	SER	-	expression tag	UNP S0DUX5
Е	1	ASN	-	expression tag	UNP S0DUX5
F	-16	MET		initiating methionine	UNP S0DUX5
F	-15	GLY	-	expression tag	UNP S0DUX5
F	-14	SER		expression tag	UNP S0DUX5
F	-13	SER		expression tag	UNP S0DUX5
F	-12	HIS		expression tag	UNP S0DUX5
F	-11	HIS	-	expression tag	UNP S0DUX5
F	-10	HIS	-	expression tag	UNP S0DUX5



Continued from previous page...


Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
F	-9	HIS	-	expression tag	UNP S0DUX5
F	-8	HIS	-	expression tag	UNP S0DUX5
F	-7	HIS	-	expression tag	UNP S0DUX5
F	-6	GLU	-	expression tag	UNP S0DUX5
F	-5	ASN	-	expression tag	UNP S0DUX5
F	-4	LEU	-	expression tag	UNP S0DUX5
F	-3	TYR	-	expression tag	UNP S0DUX5
F	-2	PHE	-	expression tag	UNP S0DUX5
F	-1	GLN	-	expression tag	UNP S0DUX5
F	0	SER	-	expression tag	UNP S0DUX5
F	1	ASN	-	expression tag	UNP S0DUX5
G	-16	MET	-	initiating methionine	UNP S0DUX5
G	-15	GLY	-	expression tag	UNP S0DUX5
G	-14	SER	-	expression tag	UNP S0DUX5
G	-13	SER	-	expression tag	UNP S0DUX5
G	-12	HIS	-	expression tag	UNP S0DUX5
G	-11	HIS	-	expression tag	UNP S0DUX5
G	-10	HIS	-	expression tag	UNP S0DUX5
G	-9	HIS	-	expression tag	UNP S0DUX5
G	-8	HIS	-	expression tag	UNP S0DUX5
G	-7	HIS	-	expression tag	UNP S0DUX5
G	-6	GLU	-	expression tag	UNP S0DUX5
G	-5	ASN	_	expression tag	UNP S0DUX5
G	-4	LEU	-	expression tag	UNP S0DUX5
G	-3	TYR	-	expression tag	UNP S0DUX5
G	-2	PHE	_	expression tag	UNP S0DUX5
G	-1	GLN	-	expression tag	UNP S0DUX5
G	0	SER	_	expression tag	UNP S0DUX5
G	1	ASN	-	expression tag	UNP S0DUX5
Н	-16	MET	-	initiating methionine	UNP S0DUX5
Н	-15	GLY	-	expression tag	UNP S0DUX5
Н	-14	SER	-	expression tag	UNP S0DUX5
Н	-13	SER	-	expression tag	UNP S0DUX5
Н	-12	HIS	-	expression tag	UNP S0DUX5
Н	-11	HIS	-	expression tag	UNP S0DUX5
Н	-10	HIS	-	expression tag	UNP S0DUX5
Н	-9	HIS	-	expression tag	UNP S0DUX5
Н	-8	HIS		expression tag	UNP S0DUX5
Н	-7	HIS	-	expression tag	UNP S0DUX5
Н	-6	GLU	-	expression tag	UNP S0DUX5
Н	-5	ASN	-	expression tag	UNP S0DUX5
Н	-4	LEU		expression tag	UNP S0DUX5



n previous	paae
	n previous

Chain	Residue	Modelled	Actual Comment		Reference
Н	-3	TYR	-	expression tag	UNP S0DUX5
Н	-2	PHE	-	expression tag	UNP S0DUX5
Н	-1	GLN	-	expression tag	UNP S0DUX5
Н	0	SER	-	expression tag	UNP S0DUX5
Н	1	ASN	=	expression tag	UNP S0DUX5


• Molecule 2 is (2E)-2-{[(1E)-{3-hydroxy-2-methyl-5-[(phosphonooxy)methyl]pyridin-4-y l}methylidene]amino}but-2-enoic acid (three-letter code: 4LM) (formula:  $C_{12}H_{15}N_2O_7P$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total 22	_				0	0
2	F	1	Total 22				P 1	0	0

• Molecule 3 is (2S)-2-[({3-hydroxy-2-methyl-5-[(phosphonooxy)methyl]pyridin-4-yl}methyl)amino]but-3-enoic acid (three-letter code: WQF) (formula:  $C_{12}H_{17}N_2O_7P$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues		Ato	ms			ZeroOcc	AltConf
3	В	1	Total	С	N	О	Р	0	0
3	Б	1	22	12	2	7	1	U	
3	C	1	Total	С	N	О	Р	0	0
9		1	22	12	2	7	1	0	0
3	D	1	Total	С	N	О	Р	0	0
9	D	1	22	12	2	7	1	U	U
3	E	1	Total	С	N	О	Р	0	0
9	<u> 1</u> 2	1	22	12	2	7	1	0	0
3	G	1	Total	С	N	О	Р	0	0
9	G	1	22	12	2	7	1	U	0
3	Н	1	Total	С	N	О	Р	0	0
	11	1	22	12	2	7	1	U	0

### • Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	106	Total O 106 106	0	0
4	В	117	Total O 117 117	0	0
4	С	122	Total O 122 122	0	0
4	D	111	Total O 111 111	0	0
4	E	140	Total O 140 140	0	0
4	F	141	Total O 141 141	0	0




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	G	154	Total O 154 154	0	0
4	Н	170	Total O 170 170	0	0



# 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Sulfhydrylase FUB7



• Molecule 1: Sulfhydrylase FUB7











## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	190.96Å 194.75Å 146.34Å	D t
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $129.32^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	49.22 - 2.16	Depositor
Resolution (A)	49.22 - 2.16	EDS
% Data completeness	98.8 (49.22-2.16)	Depositor
(in resolution range)	98.9 (49.22-2.16)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sum}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.84 (at 2.16Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
рρ.	0.177 , $0.205$	Depositor
$R, R_{free}$	0.180 , $0.207$	DCC
$R_{free}$ test set	11280 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.2	Xtriage
Anisotropy	0.299	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.37 , 47.6	EDS
L-test for twinning <sup>2</sup>	$< L >=0.49, < L^2>=0.32$	Xtriage
	0.000  for k,h,-1/2*h-1/2*k-l	
Estimated twinning fraction	0.000  for -k,-h,-1/2*h+1/2*k-l	Xtriage
	0.017  for  h,-k,-h-l	
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	27599	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 48.54 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.4733e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 4LM, WQF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bo	Bond angles	
Moi   Chain		RMSZ	# Z  > 5	RMSZ	# Z >5	
1	A	0.41	0/3372	0.58	0/4580	
1	В	0.41	0/3383	0.59	1/4595 (0.0%)	
1	С	0.42	1/3383~(0.0%)	0.61	1/4595 (0.0%)	
1	D	0.41	0/3383	0.58	0/4595	
1	Е	0.44	0/3383	0.60	0/4595	
1	F	0.42	0/3383	0.60	2/4595~(0.0%)	
1	G	0.46	0/3365	0.61	1/4570 (0.0%)	
1	Н	0.44	0/3365	0.59	0/4570	
All	All	0.42	$1/27017 \ (0.0\%)$	0.60	5/36695 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	С	0	1
1	D	0	1
1	Н	0	1
All	All	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\text{\AA})$
1	С	354	ASP	CG-OD2	5.50	1.38	1.25

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}(^{o})$
1	С	354	ASP	CB-CG-OD2	-13.12	106.50	118.30



Mol	Chain	Res	Type	Atoms	${f Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	G	428	GLN	CA-CB-CG	-7.87	96.08	113.40
1	В	428	GLN	CA-CB-CG	6.86	128.50	113.40
1	F	21	ASP	CB-CG-OD2	-6.49	112.46	118.30
1	F	428	GLN	CA-CB-CG	5.44	125.38	113.40

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	С	354	ASP	Sidechain
1	D	303	GLN	Sidechain
1	Н	373	ASN	Sidechain

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3292	0	3215	33	0
1	В	3300	0	3226	14	0
1	С	3300	0	3226	25	0
1	D	3300	0	3226	14	0
1	Е	3300	0	3226	16	0
1	F	3300	0	3228	19	0
1	G	3285	0	3204	18	0
1	Н	3285	0	3204	17	0
2	A	22	0	12	1	0
2	F	22	0	12	0	0
3	В	22	0	0	0	0
3	С	22	0	0	0	0
3	D	22	0	0	0	0
3	Ε	22	0	0	0	0
3	G	22	0	0	0	0
3	Н	22	0	0	0	0
4	A	106	0	0	0	0
4	В	117	0	0	0	0
4	С	122	0	0	0	0
4	D	111	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	Ε	140	0	0	0	0
4	F	141	0	0	0	0
4	G	154	0	0	0	0
4	Н	170	0	0	0	0
All	All	27599	0	25779	142	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 142 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:F:158:MET:HE1	1:F:339:LEU:HD21	1.60	0.83
1:B:383:HIS:CE1	1:B:385:TRP:HB3	2.25	0.71
1:F:196[A]:ARG:NH1	1:F:199:GLU:OE2	2.22	0.71
1:E:158:MET:HE1	1:E:339:LEU:HD21	1.75	0.69
1:E:383:HIS:CE1	1:E:385:TRP:HB3	2.29	0.68

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	426/450~(95%)	414 (97%)	11 (3%)	1 (0%)	47	46
1	В	$427/450 \ (95\%)$	413 (97%)	13 (3%)	1 (0%)	47	46
1	С	427/450 (95%)	416 (97%)	10 (2%)	1 (0%)	47	46
1	D	$427/450 \ (95\%)$	418 (98%)	8 (2%)	1 (0%)	47	46
1	Е	427/450 (95%)	411 (96%)	15 (4%)	1 (0%)	47	46
1	F	$427/450 \ (95\%)$	414 (97%)	13 (3%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	425/450 (94%)	414 (97%)	10 (2%)	1 (0%)	47 46	
1	Н	425/450 (94%)	416 (98%)	8 (2%)	1 (0%)	47 46	
All	All	3411/3600 (95%)	3316 (97%)	88 (3%)	7 (0%)	47 46	

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	161	PRO
1	D	161	PRO
1	Е	211	LYS
1	В	161	PRO
1	G	161	PRO

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	346/366 (94%)	346 (100%)	0	100 100
1	В	347/366 (95%)	345 (99%)	2 (1%)	86 90
1	C	347/366 (95%)	345 (99%)	2 (1%)	86 90
1	D	347/366~(95%)	345 (99%)	2 (1%)	86 90
1	E	347/366 (95%)	347 (100%)	0	100 100
1	F	347/366~(95%)	347 (100%)	0	100 100
1	G	345/366 (94%)	345 (100%)	0	100 100
1	Н	345/366 (94%)	343 (99%)	2 (1%)	86 90
All	All	2771/2928 (95%)	2763 (100%)	8 (0%)	92 95

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Н	354	ASP
1	Н	62	MET



Mol	Chain	Res	Type
1	D	62	MET
1	С	211	LYS
1	D	211	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type	
1	A	7	GLN	
1	A	428	GLN	
1	Е	118	ASN	

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mo	Type	Chain	Res	Link	Bo	Bond lengths			Bond angles		
IVIO	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2	
3	WQF	D	501	1	22,22,22	2.05	6 (27%)	26,31,31	2.81	10 (38%)	
3	WQF	Н	501	1	22,22,22	2.17	7 (31%)	26,31,31	3.15	4 (15%)	



Mol	Mol Type Ch		Res	Link	Во	ond leng	$\overline{ ext{gths}}$	Bond angles		
MIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	WQF	С	501	1	22,22,22	2.04	6 (27%)	26,31,31	2.79	6 (23%)
3	WQF	В	501	1	22,22,22	2.10	6 (27%)	26,31,31	2.90	9 (34%)
2	4LM	F	501	-	21,22,22	2.27	7 (33%)	25,31,31	1.35	2 (8%)
3	WQF	Е	501	1	22,22,22	2.34	7 (31%)	26,31,31	3.02	6 (23%)
2	4LM	A	501	-	21,22,22	2.36	6 (28%)	25,31,31	1.99	6 (24%)
3	WQF	G	501	-	22,22,22	2.18	7 (31%)	26,31,31	3.05	5 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	WQF	D	501	1	-	2/17/17/17	0/1/1/1
3	WQF	Н	501	1	-	3/17/17/17	0/1/1/1
3	WQF	С	501	1	-	4/17/17/17	0/1/1/1
3	WQF	В	501	1	-	2/17/17/17	0/1/1/1
2	4LM	F	501	-	-	4/15/17/17	0/1/1/1
3	WQF	Е	501	1	-	2/17/17/17	0/1/1/1
2	4LM	A	501	-	-	5/15/17/17	0/1/1/1
3	WQF	G	501	-	-	1/17/17/17	0/1/1/1

The worst 5 of 52 bond length outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}( ext{\AA})$
3	Е	501	WQF	C13-C12	6.19	1.59	1.51
3	Н	501	WQF	P08-O07	5.45	1.77	1.60
3	В	501	WQF	C13-C12	5.18	1.58	1.51
3	Е	501	WQF	P08-O07	5.09	1.76	1.60
3	G	501	WQF	C13-C12	4.81	1.58	1.51

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	Н	501	WQF	C13-C12-C05	11.41	132.39	119.71
3	G	501	WQF	C13-C12-C05	11.02	131.96	119.71
3	Е	501	WQF	C13-C12-C05	10.81	131.73	119.71
3	В	501	WQF	C13-C12-C05	10.03	130.87	119.71
3	С	501	WQF	C13-C12-C05	9.78	130.59	119.71



There are no chirality outliers.

5 of 23 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	4LM	C-CA-CB-CG
2	F	501	4LM	O1-C-CA-CB
2	F	501	4LM	O2-C-CA-CB
2	F	501	4LM	C-CA-CB-CG
3	В	501	WQF	C28-C25-C26-C27


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	4LM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







































# 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	428/450 (95%)	-0.07	15 (3%) 44 52	17, 30, 50, 62	0
1	В	428/450 (95%)	0.01	21 (4%) 29 38	17, 30, 51, 64	0
1	С	428/450 (95%)	0.01	17 (3%) 38 47	17, 30, 51, 69	0
1	D	428/450 (95%)	-0.08	14 (3%) 46 55	18, 30, 51, 74	0
1	E	428/450 (95%)	-0.28	5 (1%) 79 83	13, 27, 46, 66	0
1	F	428/450 (95%)	-0.29	6 (1%) 75 80	16, 27, 46, 65	0
1	G	427/450 (94%)	-0.33	4 (0%) 84 88	14, 23, 42, 67	0
1	Н	427/450 (94%)	-0.33	2 (0%) 91 93	14, 24, 42, 67	0
All	All	3422/3600 (95%)	-0.17	84 (2%) 57 65	13, 28, 48, 74	0

The worst 5 of 84 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	432	GLY	7.6
1	A	355	ALA	5.3
1	С	432	GLY	5.3
1	D	355	ALA	4.5
1	В	355	ALA	4.5

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	4LM	A	501	22/22	0.96	0.12	21,27,40,46	0
2	4LM	F	501	22/22	0.96	0.13	18,24,39,41	0
3	WQF	С	501	22/22	0.96	0.14	19,30,43,47	0
3	WQF	В	501	22/22	0.97	0.14	20,28,48,51	0
3	WQF	D	501	22/22	0.97	0.15	22,31,46,47	0
3	WQF	Е	501	22/22	0.97	0.13	17,27,40,40	0
3	WQF	Н	501	22/22	0.97	0.13	15,26,39,43	0
3	WQF	G	501	22/22	0.98	0.11	16,25,37,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







































# 6.5 Other polymers (i)

There are no such residues in this entry.

