Full wwPDB X-ray Structure Validation Report (i)

May 15, 2020 – 08:07 pm BST

PDB ID	:	1ET0
Title	:	CRYSTAL STRUCTURE OF AMINODEOXYCHORISMATE LYASE FROM ESCHERICHIA COLI
Authors	:	Nakai, T.; Mizutani, H.; Miyahara, I.; Hirotsu, K.; Takeda, S.; Jhee, K.H.; Yoshimura, T.; Esaki, N.
Deposited on	:	2000-04-12
Resolution	:	2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	269	70% 22% • 6%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2088 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-AMINO-4-DEOXYCHORISMATE LYASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	254	1947	1217	345	368	17	0	0	0

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	15	8	1	5	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	126	126	126	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: 4-AMINO-4-DEOXYCHORISMATE LYASE

Chain A: 70% 22% • 6%



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α, β, γ	40.50Å 73.93Å 83.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.20	Depositor
% Data completeness (in resolution range)	91.4 (8.00-2.20)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.196 , 0.279	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2088	wwPDB-VP
Average B, all atoms (Å^2)	18.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.40	0/1976	0.68	1/2676 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	121	LEU	CA-CB-CG	5.87	128.81	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1947	0	1951	34	0
2	A	15	0	7	0	0
3	A	126	0	0	1	0
All	All	2088	0	1958	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:VAL:HG22	1:A:121:LEU:HD13	1.68	0.74
1:A:139:ILE:HG13	1:A:262:MET:HG2	1.69	0.73
1:A:139:ILE:HG13	1:A:262:MET:CG	2.26	0.65
1:A:229:LEU:HD21	1:A:280:LEU:HD21	1.84	0.60
1:A:199:LEU:O	1:A:209:THR:HG22	2.03	0.58
1:A:254:ILE:HD13	1:A:264:VAL:HG21	1.86	0.57
1:A:261:VAL:HG21	1:A:285:ALA:HA	1.87	0.56
1:A:57:ILE:O	1:A:61:GLN:HG3	2.04	0.56
1:A:219:GLY:O	1:A:223:GLN:HG3	2.07	0.55
1:A:80:GLN:O	1:A:84:THR:HG23	2.07	0.54
1:A:8:PHE:HB2	1:A:15:GLN:O	2.07	0.53
1:A:139:ILE:N	1:A:265:CYS:HG	2.07	0.53
1:A:182:ALA:O	1:A:194:CYS:HA	2.09	0.53
1:A:199:LEU:HD11	1:A:252:MET:HG2	1.92	0.52
1:A:200:PHE:CD1	1:A:209:THR:HG23	2.47	0.49
1:A:203:LYS:HA	1:A:250:ASP:OD2	2.13	0.48
1:A:63:ALA:HA	1:A:158:ILE:HD12	1.95	0.47
1:A:55:ALA:HB1	1:A:220:ILE:HG13	1.96	0.47
1:A:89:GLN:NE2	1:A:125:ALA:HA	2.29	0.47
1:A:244:GLU:H	1:A:244:GLU:CD	2.18	0.46
1:A:36:PHE:O	1:A:159:LYS:HD3	2.15	0.46
1:A:254:ILE:HD13	1:A:264:VAL:CG2	2.47	0.45
1:A:102:ARG:HD3	1:A:111:THR:HG23	1.99	0.44
1:A:10:ILE:O	1:A:13:HIS:HD2	2.00	0.44
1:A:89:GLN:O	1:A:90:GLN:HB2	2.18	0.43
1:A:185:LEU:HD23	1:A:191:VAL:HA	1.99	0.43
1:A:14:LYS:HD3	1:A:14:LYS:H	1.83	0.42
1:A:106:GLY:HA2	1:A:112:LEU:O	2.20	0.42
1:A:147:ARG:HE	1:A:187:SER:HA	1.85	0.41
1:A:251:GLU:HG3	3:A:517:HOH:O	2.20	0.41
1:A:58:GLN:HA	1:A:61:GLN:HE21	1.85	0.41
1:A:178:ASN:HD22	1:A:178:ASN:C	2.24	0.41
1:A:78:LEU:HG	1:A:82:MET:CE	2.51	0.40
1:A:285:ALA:HB3	1:A:286:PRO:HD3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	250/269 (93%)	240 (96%)	10 (4%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	214/227 (94%)	196 (92%)	18 (8%)	11 11

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	MET
1	A	13	HIS
1	A	14	LYS
1	A	40	ARG
1	A	49	LYS
1	A	59	ARG
1	A	74	PHE
1	A	121	LEU
1	A	178	ASN
1	A	181	GLU
1	A	202	ARG
1	A	203	LYS
1	A	209	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	233	SER
1	A	243	LEU
1	A	259	MET
1	A	284	LEU
1	A	289	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	61	GLN
1	A	80	GLN
1	A	89	GLN
1	A	162	ASN
1	A	176	GLN
1	A	178	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	PLP	A	413	1	15,15,16	1.71	4 (26%)	20,22,23	1.78	4 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	413	1	-	2/6/6/8	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	413	PLP	C3-C2	-4.16	1.36	1.40
2	A	413	PLP	C2A-C2	2.30	1.54	1.50
2	A	413	PLP	C2-N1	2.20	1.38	1.33
2	A	413	PLP	P-O3P	-2.17	1.46	1.54

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	413	PLP	O4P-C5A-C5	4.89	118.67	109.35
2	A	413	PLP	O2P-P-O4P	-3.02	98.69	106.73
2	A	413	PLP	C5-C6-N1	-2.64	119.42	123.82
2	A	413	PLP	C6-C5-C4	2.02	119.75	118.16

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	413	PLP	C4-C5-C5A-O4P
2	A	413	PLP	C6-C5-C5A-O4P

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.