



wwPDB X-ray Structure Validation Summary Report i

Aug 26, 2023 – 08:57 PM EDT

PDB ID : 3ETB
Title : Crystal structure of the engineered neutralizing antibody M18 complexed with anthrax protective antigen domain 4
Authors : Monzingo, A.F.; Leysath, C.E.; Barnett, J.; Iverson, B.L.; Georgiou, G.; Robertus, J.D.
Deposited on : 2008-10-07
Resolution : 3.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see references ①) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1212 (4.00-3.60)
Clashscore	141614	1288 (4.00-3.60)
Ramachandran outliers	138981	1243 (4.00-3.60)
Sidechain outliers	138945	1237 (4.00-3.60)
RSRZ outliers	127900	1121 (4.00-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain			
2	K	144	2%	42%	47%	10% •
2	L	144	.%	41%	50%	8% ..
2	M	144		44%	48%	7% •

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 11546 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antibody M18 light chain and antibody M18 heavy chain linked with a synthetic (GGGGS)4 linker.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	F	228	Total	C 1742	N 1091	O 296	S 347	8	0	0
1	G	229	Total	C 1743	N 1091	O 296	S 348	8	0	0
1	H	229	Total	C 1743	N 1091	O 296	S 348	8	0	0
1	I	228	Total	C 1742	N 1091	O 296	S 347	8	0	0

There are 140 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-4	MET	-	expression tag	PDB 3ETB
F	-3	ALA	-	expression tag	PDB 3ETB
F	-2	ASP	-	expression tag	PDB 3ETB
F	-1	TYR	-	expression tag	PDB 3ETB
F	0	LYS	-	expression tag	PDB 3ETB
F	21	VAL	ILE	engineered mutation	PDB 3ETB
F	46	PHE	LEU	engineered mutation	PDB 3ETB
F	56	PRO	SER	engineered mutation	PDB 3ETB
F	76	ASN	SER	engineered mutation	PDB 3ETB
F	78	LEU	GLN	engineered mutation	PDB 3ETB
F	94	PRO	LEU	engineered mutation	PDB 3ETB
F	109	GLY	-	linker	PDB 3ETB
F	110	GLY	-	linker	PDB 3ETB
F	111	GLY	-	linker	PDB 3ETB
F	112	GLY	-	linker	PDB 3ETB
F	113	SER	-	linker	PDB 3ETB
F	114	GLY	-	linker	PDB 3ETB
F	115	GLY	-	linker	PDB 3ETB
F	116	GLY	-	linker	PDB 3ETB
F	117	GLY	-	linker	PDB 3ETB

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	118	SER	-	linker	PDB 3ETB
F	119	GLY	-	linker	PDB 3ETB
F	120	GLY	-	linker	PDB 3ETB
F	121	GLY	-	linker	PDB 3ETB
F	122	GLY	-	linker	PDB 3ETB
F	123	SER	-	linker	PDB 3ETB
F	124	GLY	-	linker	PDB 3ETB
F	125	GLY	-	linker	PDB 3ETB
F	126	GLY	-	linker	PDB 3ETB
F	127	GLY	-	linker	PDB 3ETB
F	128	SER	-	linker	PDB 3ETB
F	1030	ASN	SER	engineered mutation	PDB 3ETB
F	1057	SER	THR	engineered mutation	PDB 3ETB
F	1064	GLU	LYS	engineered mutation	PDB 3ETB
F	1068	ILE	THR	engineered mutation	PDB 3ETB
G	-4	MET	-	expression tag	PDB 3ETB
G	-3	ALA	-	expression tag	PDB 3ETB
G	-2	ASP	-	expression tag	PDB 3ETB
G	-1	TYR	-	expression tag	PDB 3ETB
G	0	LYS	-	expression tag	PDB 3ETB
G	21	VAL	ILE	engineered mutation	PDB 3ETB
G	46	PHE	LEU	engineered mutation	PDB 3ETB
G	56	PRO	SER	engineered mutation	PDB 3ETB
G	76	ASN	SER	engineered mutation	PDB 3ETB
G	78	LEU	GLN	engineered mutation	PDB 3ETB
G	94	PRO	LEU	engineered mutation	PDB 3ETB
G	109	GLY	-	linker	PDB 3ETB
G	110	GLY	-	linker	PDB 3ETB
G	111	GLY	-	linker	PDB 3ETB
G	112	GLY	-	linker	PDB 3ETB
G	113	SER	-	linker	PDB 3ETB
G	114	GLY	-	linker	PDB 3ETB
G	115	GLY	-	linker	PDB 3ETB
G	116	GLY	-	linker	PDB 3ETB
G	117	GLY	-	linker	PDB 3ETB
G	118	SER	-	linker	PDB 3ETB
G	119	GLY	-	linker	PDB 3ETB
G	120	GLY	-	linker	PDB 3ETB
G	121	GLY	-	linker	PDB 3ETB
G	122	GLY	-	linker	PDB 3ETB
G	123	SER	-	linker	PDB 3ETB
G	124	GLY	-	linker	PDB 3ETB

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	125	GLY	-	linker	PDB 3ETB
G	126	GLY	-	linker	PDB 3ETB
G	127	GLY	-	linker	PDB 3ETB
G	128	SER	-	linker	PDB 3ETB
G	1030	ASN	SER	engineered mutation	PDB 3ETB
G	1057	SER	THR	engineered mutation	PDB 3ETB
G	1064	GLU	LYS	engineered mutation	PDB 3ETB
G	1068	ILE	THR	engineered mutation	PDB 3ETB
H	-4	MET	-	expression tag	PDB 3ETB
H	-3	ALA	-	expression tag	PDB 3ETB
H	-2	ASP	-	expression tag	PDB 3ETB
H	-1	TYR	-	expression tag	PDB 3ETB
H	0	LYS	-	expression tag	PDB 3ETB
H	21	VAL	ILE	engineered mutation	PDB 3ETB
H	46	PHE	LEU	engineered mutation	PDB 3ETB
H	56	PRO	SER	engineered mutation	PDB 3ETB
H	76	ASN	SER	engineered mutation	PDB 3ETB
H	78	LEU	GLN	engineered mutation	PDB 3ETB
H	94	PRO	LEU	engineered mutation	PDB 3ETB
H	109	GLY	-	linker	PDB 3ETB
H	110	GLY	-	linker	PDB 3ETB
H	111	GLY	-	linker	PDB 3ETB
H	112	GLY	-	linker	PDB 3ETB
H	113	SER	-	linker	PDB 3ETB
H	114	GLY	-	linker	PDB 3ETB
H	115	GLY	-	linker	PDB 3ETB
H	116	GLY	-	linker	PDB 3ETB
H	117	GLY	-	linker	PDB 3ETB
H	118	SER	-	linker	PDB 3ETB
H	119	GLY	-	linker	PDB 3ETB
H	120	GLY	-	linker	PDB 3ETB
H	121	GLY	-	linker	PDB 3ETB
H	122	GLY	-	linker	PDB 3ETB
H	123	SER	-	linker	PDB 3ETB
H	124	GLY	-	linker	PDB 3ETB
H	125	GLY	-	linker	PDB 3ETB
H	126	GLY	-	linker	PDB 3ETB
H	127	GLY	-	linker	PDB 3ETB
H	128	SER	-	linker	PDB 3ETB
H	1030	ASN	SER	engineered mutation	PDB 3ETB
H	1057	SER	THR	engineered mutation	PDB 3ETB
H	1064	GLU	LYS	engineered mutation	PDB 3ETB

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	1068	ILE	THR	engineered mutation	PDB 3ETB
I	-4	MET	-	expression tag	PDB 3ETB
I	-3	ALA	-	expression tag	PDB 3ETB
I	-2	ASP	-	expression tag	PDB 3ETB
I	-1	TYR	-	expression tag	PDB 3ETB
I	0	LYS	-	expression tag	PDB 3ETB
I	21	VAL	ILE	engineered mutation	PDB 3ETB
I	46	PHE	LEU	engineered mutation	PDB 3ETB
I	56	PRO	SER	engineered mutation	PDB 3ETB
I	76	ASN	SER	engineered mutation	PDB 3ETB
I	78	LEU	GLN	engineered mutation	PDB 3ETB
I	94	PRO	LEU	engineered mutation	PDB 3ETB
I	109	GLY	-	linker	PDB 3ETB
I	110	GLY	-	linker	PDB 3ETB
I	111	GLY	-	linker	PDB 3ETB
I	112	GLY	-	linker	PDB 3ETB
I	113	SER	-	linker	PDB 3ETB
I	114	GLY	-	linker	PDB 3ETB
I	115	GLY	-	linker	PDB 3ETB
I	116	GLY	-	linker	PDB 3ETB
I	117	GLY	-	linker	PDB 3ETB
I	118	SER	-	linker	PDB 3ETB
I	119	GLY	-	linker	PDB 3ETB
I	120	GLY	-	linker	PDB 3ETB
I	121	GLY	-	linker	PDB 3ETB
I	122	GLY	-	linker	PDB 3ETB
I	123	SER	-	linker	PDB 3ETB
I	124	GLY	-	linker	PDB 3ETB
I	125	GLY	-	linker	PDB 3ETB
I	126	GLY	-	linker	PDB 3ETB
I	127	GLY	-	linker	PDB 3ETB
I	128	SER	-	linker	PDB 3ETB
I	1030	ASN	SER	engineered mutation	PDB 3ETB
I	1057	SER	THR	engineered mutation	PDB 3ETB
I	1064	GLU	LYS	engineered mutation	PDB 3ETB
I	1068	ILE	THR	engineered mutation	PDB 3ETB

- Molecule 2 is a protein called Anthrax Protective Antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	J	144	Total	C	N	O	S	0	0	0
			1146	722	192	231	1			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	K	144	1146	722	192	231	1	0	0	0
2	L	143	1138	718	191	228	1	0	0	0
2	M	144	1146	722	192	231	1	0	0	0


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Antibody M18 light chain and antibody M18 heavy chain linked with a synthetic (GGGGS)₄ linker




- Molecule 1: Antibody M18 light chain and antibody M18 heavy chain linked with a synthetic (GGGGS)₄ linker



- Molecule 1: Antibody M18 light chain and antibody M18 heavy chain linked with a synthetic (GGGGS)₄ linker






- Molecule 1: Antibody M18 light chain and antibody M18 heavy chain linked with a synthetic (GGGGS)₄ linker



- Molecule 2: Anthrax Protective Antigen



- Molecule 2: Anthrax Protective Antigen



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.94Å 299.72Å 68.95Å 90.00° 94.45° 90.00°	Depositor
Resolution (Å)	20.00 – 3.80 43.84 – 3.81	Depositor EDS
% Data completeness (in resolution range)	83.8 (20.00-3.80) 89.8 (43.84-3.81)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.00 (at 3.77Å)	Xtriage
Refinement program	CNS	Depositor
R , R_{free}	0.232 , 0.276 0.223 , 0.225	Depositor DCC
R_{free} test set	942 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	39.3	Xtriage
Anisotropy	0.633	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 1.6	EDS
L-test for twinning ²	$< L > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	11546	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	F	0.48	0/1781	0.70	0/2413
1	G	0.49	0/1782	0.71	0/2416
1	H	0.49	0/1782	0.70	0/2416
1	I	0.49	0/1781	0.70	0/2413
2	J	0.44	0/1162	0.66	0/1566
2	K	0.47	0/1162	0.69	0/1566
2	L	0.47	0/1154	0.70	0/1555
2	M	0.47	0/1162	0.68	0/1566
All	All	0.48	0/11766	0.69	0/15911

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1742	0	1655	151	0
1	G	1743	0	1646	141	0
1	H	1743	0	1646	151	0
1	I	1742	0	1655	142	0
2	J	1146	0	1134	104	0
2	K	1146	0	1134	94	0
2	L	1138	0	1130	100	0
2	M	1146	0	1134	96	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	11546	0	11134	953	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 42.

The worst 5 of 953 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:1006:GLN:HE22	1:G:1091:PHE:HA	1.29	0.96
1:F:1006:GLN:HE22	1:F:1091:PHE:HA	1.27	0.95
1:G:1011:LEU:HD13	1:G:1110:THR:HG23	1.47	0.95
2:M:629:LEU:HD23	2:M:629:LEU:H	1.32	0.93
1:H:1050:ARG:HE	1:H:1058:ASN:HD22	1.18	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	F	224/252 (89%)	168 (75%)	36 (16%)	20 (9%)	1 12
1	G	225/252 (89%)	165 (73%)	37 (16%)	23 (10%)	0 9
1	H	225/252 (89%)	164 (73%)	40 (18%)	21 (9%)	0 11
1	I	224/252 (89%)	165 (74%)	39 (17%)	20 (9%)	1 12
2	J	142/144 (99%)	100 (70%)	28 (20%)	14 (10%)	0 10
2	K	142/144 (99%)	100 (70%)	28 (20%)	14 (10%)	0 10
2	L	141/144 (98%)	105 (74%)	26 (18%)	10 (7%)	1 17
2	M	142/144 (99%)	100 (70%)	30 (21%)	12 (8%)	1 12
All	All	1465/1584 (92%)	1067 (73%)	264 (18%)	134 (9%)	1 12

5 of 134 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	1033	TRP
1	F	1098	LEU
1	G	1002	VAL
1	G	1033	TRP
1	G	1098	LEU

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	F	188/201 (94%)	171 (91%)	17 (9%)	9 37
1	G	187/201 (93%)	171 (91%)	16 (9%)	10 40
1	H	187/201 (93%)	171 (91%)	16 (9%)	10 40
1	I	188/201 (94%)	170 (90%)	18 (10%)	8 34
2	J	128/131 (98%)	119 (93%)	9 (7%)	15 46
2	K	128/131 (98%)	120 (94%)	8 (6%)	18 49
2	L	127/131 (97%)	120 (94%)	7 (6%)	21 53
2	M	128/131 (98%)	122 (95%)	6 (5%)	26 56
All	All	1261/1328 (95%)	1164 (92%)	97 (8%)	13 43

5 of 97 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	97	THR
2	J	667	SER
1	I	1040	ARG
1	I	1102	TYR
2	J	730	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 52 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	1058	ASN
2	K	630	ASN
2	M	705	ASN
2	J	630	ASN
2	J	709	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	F	228/252 (90%)	-0.35	0	100	100	0
1	G	229/252 (90%)	-0.29	0	100	100	0
1	H	229/252 (90%)	-0.33	0	100	100	0
1	I	228/252 (90%)	-0.34	0	100	100	0
2	J	144/144 (100%)	0.15	0	100	100	0
2	K	144/144 (100%)	-0.21	3 (2%)	63	55	0
2	L	143/144 (99%)	0.30	2 (1%)	75	68	0
2	M	144/144 (100%)	-0.27	0	100	100	0
All	All	1489/1584 (94%)	-0.21	5 (0%)	94	91	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	K	707	ILE	3.0
2	L	629	LEU	2.9
2	L	616	HIS	2.5
2	K	708	ILE	2.3
2	K	712	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates i

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.