



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 16, 2023 – 09:27 PM EST

PDB ID : 3F0Y
Title : Crystal structure of the human Adenovirus type 14 fiber knob
Authors : Persson, B.D.; Reiter, D.M.; Arnberg, N.; Stehle, T.
Deposited on : 2008-10-27
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	208	 2% 89% 6% 5%
1	B	208	 % 90% 5% 5%
1	C	208	 % 92% . 5%
1	D	208	 % 88% 7% 5%
1	E	208	 2% 88% 7% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	208	 88% 7% 5% 2%
1	G	208	 89% 6% 5% 2%
1	H	208	 83% 12% 5% 2%
1	I	208	 89% 6% 5% 2%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 15886 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fiber protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	198	1570	986	259	315	10	0	4	0
1	B	198	1547	969	255	313	10	0	1	0
1	C	198	1552	974	255	313	10	0	2	0
1	D	198	1572	989	257	315	11	0	5	0
1	E	198	1561	982	255	314	10	0	3	0
1	F	198	1565	985	255	315	10	0	4	0
1	G	199	1569	988	256	315	10	0	3	0
1	H	198	1559	978	256	315	10	0	3	0
1	I	198	1549	970	256	313	10	0	1	0

There are 45 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	118	GLY	-	expression tag	UNP Q8V791
A	119	SER	-	expression tag	UNP Q8V791
A	120	HIS	-	expression tag	UNP Q8V791
A	121	MET	-	expression tag	UNP Q8V791
A	122	GLY	-	expression tag	UNP Q8V791
B	118	GLY	-	expression tag	UNP Q8V791
B	119	SER	-	expression tag	UNP Q8V791
B	120	HIS	-	expression tag	UNP Q8V791
B	121	MET	-	expression tag	UNP Q8V791
B	122	GLY	-	expression tag	UNP Q8V791
C	118	GLY	-	expression tag	UNP Q8V791

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	119	SER	-	expression tag	UNP Q8V791
C	120	HIS	-	expression tag	UNP Q8V791
C	121	MET	-	expression tag	UNP Q8V791
C	122	GLY	-	expression tag	UNP Q8V791
D	118	GLY	-	expression tag	UNP Q8V791
D	119	SER	-	expression tag	UNP Q8V791
D	120	HIS	-	expression tag	UNP Q8V791
D	121	MET	-	expression tag	UNP Q8V791
D	122	GLY	-	expression tag	UNP Q8V791
E	118	GLY	-	expression tag	UNP Q8V791
E	119	SER	-	expression tag	UNP Q8V791
E	120	HIS	-	expression tag	UNP Q8V791
E	121	MET	-	expression tag	UNP Q8V791
E	122	GLY	-	expression tag	UNP Q8V791
F	118	GLY	-	expression tag	UNP Q8V791
F	119	SER	-	expression tag	UNP Q8V791
F	120	HIS	-	expression tag	UNP Q8V791
F	121	MET	-	expression tag	UNP Q8V791
F	122	GLY	-	expression tag	UNP Q8V791
G	118	GLY	-	expression tag	UNP Q8V791
G	119	SER	-	expression tag	UNP Q8V791
G	120	HIS	-	expression tag	UNP Q8V791
G	121	MET	-	expression tag	UNP Q8V791
G	122	GLY	-	expression tag	UNP Q8V791
H	118	GLY	-	expression tag	UNP Q8V791
H	119	SER	-	expression tag	UNP Q8V791
H	120	HIS	-	expression tag	UNP Q8V791
H	121	MET	-	expression tag	UNP Q8V791
H	122	GLY	-	expression tag	UNP Q8V791
I	118	GLY	-	expression tag	UNP Q8V791
I	119	SER	-	expression tag	UNP Q8V791
I	120	HIS	-	expression tag	UNP Q8V791
I	121	MET	-	expression tag	UNP Q8V791
I	122	GLY	-	expression tag	UNP Q8V791

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	H	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is IMIDAZOLE (three-letter code: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N 5 3 2	0	0
3	A	1	Total C N 5 3 2	0	0
3	C	1	Total C N 5 3 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	183	Total O 183 183	0	0
4	B	225	Total O 225 225	0	0
4	C	185	Total O 185 185	0	0
4	D	216	Total O 216 216	0	0
4	E	199	Total O 199 199	0	0
4	F	223	Total O 223 223	0	0
4	G	218	Total O 218 218	0	0
4	H	183	Total O 183 183	0	0
4	I	171	Total O 171 171	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fiber protein




- Molecule 1: Fiber protein




- Molecule 1: Fiber protein




- Molecule 1: Fiber protein



- Molecule 1: Fiber protein




- Molecule 1: Fiber protein

Chain F:  88% 7% 5%




• Molecule 1: Fiber protein

Chain G:  2% 89% 6% 5%




• Molecule 1: Fiber protein

Chain H:  2% 83% 12% 5%



• Molecule 1: Fiber protein

Chain I:  2% 89% 6% 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 1 2	Depositor
Cell constants a, b, c, α , β , γ	106.47Å 106.47Å 311.71Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.40 – 1.80 47.38 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.40-1.80) 99.3 (47.38-1.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.44 (at 1.81Å)	Xtrriage
Refinement program	REFMAC 5.3.0037	Depositor
R, R_{free}	0.188 , 0.215 0.192 , 0.217	Depositor DCC
R_{free} test set	9272 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	17.5	Xtrriage
Anisotropy	0.014	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	0.046 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15886	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/1617	0.49	0/2205
1	B	0.29	0/1584	0.50	0/2160
1	C	0.28	0/1592	0.49	0/2172
1	D	0.29	0/1622	0.52	0/2212
1	E	0.29	0/1605	0.52	0/2190
1	F	0.29	0/1612	0.51	0/2200
1	G	0.28	0/1613	0.49	0/2201
1	H	0.29	0/1599	0.50	0/2182
1	I	0.29	0/1586	0.51	0/2163
All	All	0.28	0/14430	0.50	0/19685

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1570	0	1505	7	0
1	B	1547	0	1475	6	0
1	C	1552	0	1486	5	0
1	D	1572	0	1508	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1561	0	1494	15	0
1	F	1565	0	1502	10	0
1	G	1569	0	1506	10	0
1	H	1559	0	1492	18	0
1	I	1549	0	1476	10	0
2	A	12	0	16	0	0
2	C	6	0	8	0	0
2	H	6	0	8	0	0
3	A	10	0	10	0	0
3	C	5	0	5	0	0
4	A	183	0	0	0	0
4	B	225	0	0	2	0
4	C	185	0	0	2	0
4	D	216	0	0	0	0
4	E	199	0	0	0	0
4	F	223	0	0	1	0
4	G	218	0	0	1	0
4	H	183	0	0	1	0
4	I	171	0	0	0	0
All	All	15886	0	13491	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 76 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:281:ALA:HA	1:I:282:ILE:HG23	1.47	0.94
1:D:151:SER:HA	1:D:152:GLU:HB2	1.53	0.89
1:G:319[B]:TYR:OH	1:H:319:TYR:OH	1.88	0.88
1:E:319[A]:TYR:OH	1:F:319[A]:TYR:OH	1.93	0.85
1:E:319[B]:TYR:OH	1:F:319[B]:TYR:OH	1.99	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	200/208 (96%)	195 (98%)	5 (2%)	0	100	100
1	B	197/208 (95%)	192 (98%)	5 (2%)	0	100	100
1	C	198/208 (95%)	194 (98%)	4 (2%)	0	100	100
1	D	201/208 (97%)	192 (96%)	7 (4%)	2 (1%)	15	5
1	E	199/208 (96%)	194 (98%)	4 (2%)	1 (0%)	29	15
1	F	200/208 (96%)	196 (98%)	4 (2%)	0	100	100
1	G	200/208 (96%)	198 (99%)	2 (1%)	0	100	100
1	H	199/208 (96%)	192 (96%)	7 (4%)	0	100	100
1	I	197/208 (95%)	191 (97%)	5 (2%)	1 (0%)	29	15
All	All	1791/1872 (96%)	1744 (97%)	43 (2%)	4 (0%)	47	33

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	152	GLU
1	E	152	GLU
1	I	282	ILE
1	D	300	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/182 (98%)	176 (99%)	2 (1%)	73	68
1	B	175/182 (96%)	173 (99%)	2 (1%)	73	68
1	C	176/182 (97%)	174 (99%)	2 (1%)	73	68
1	D	179/182 (98%)	177 (99%)	2 (1%)	73	68
1	E	177/182 (97%)	176 (99%)	1 (1%)	86	84
1	F	178/182 (98%)	176 (99%)	2 (1%)	73	68
1	G	178/182 (98%)	177 (99%)	1 (1%)	86	84
1	H	177/182 (97%)	174 (98%)	3 (2%)	60	51
1	I	175/182 (96%)	173 (99%)	2 (1%)	73	68
All	All	1593/1638 (97%)	1576 (99%)	17 (1%)	73	68

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	297	ASN
1	I	279	GLN
1	D	279	GLN
1	E	135	TRP
1	F	135	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	297	ASN
1	I	218	HIS
1	I	253	ASN
1	E	218	HIS
1	D	279	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	IMD	A	326	-	3,5,5	0.41	0	4,5,5	0.58	0
2	GOL	A	2	-	5,5,5	0.34	0	5,5,5	0.30	0
2	GOL	H	1	-	5,5,5	0.40	0	5,5,5	0.22	0
2	GOL	A	3	-	5,5,5	0.36	0	5,5,5	0.22	0
3	IMD	C	2	-	3,5,5	0.42	0	4,5,5	0.59	0
3	IMD	A	1	-	3,5,5	0.43	0	4,5,5	0.57	0
2	GOL	C	4	-	5,5,5	0.38	0	5,5,5	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IMD	A	326	-	-	-	0/1/1/1
2	GOL	A	2	-	-	4/4/4/4	-
2	GOL	H	1	-	-	0/4/4/4	-
2	GOL	A	3	-	-	2/4/4/4	-
3	IMD	C	2	-	-	-	0/1/1/1
3	IMD	A	1	-	-	-	0/1/1/1
2	GOL	C	4	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2	GOL	O1-C1-C2-C3
2	A	2	GOL	C1-C2-C3-O3
2	A	3	GOL	C1-C2-C3-O3
2	C	4	GOL	O1-C1-C2-C3
2	C	4	GOL	C1-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	198/208 (95%)	-0.03	4 (2%) 65 61	16, 22, 26, 30	0
1	B	198/208 (95%)	-0.25	2 (1%) 82 80	12, 16, 23, 26	0
1	C	198/208 (95%)	-0.12	3 (1%) 73 70	15, 21, 27, 30	0
1	D	198/208 (95%)	-0.24	3 (1%) 73 70	12, 17, 24, 26	0
1	E	198/208 (95%)	-0.16	4 (2%) 65 61	12, 18, 25, 28	0
1	F	198/208 (95%)	-0.34	1 (0%) 91 89	10, 16, 23, 25	0
1	G	199/208 (95%)	-0.21	5 (2%) 57 52	13, 17, 25, 29	0
1	H	198/208 (95%)	-0.16	4 (2%) 65 61	14, 20, 25, 29	0
1	I	198/208 (95%)	-0.11	5 (2%) 57 52	15, 21, 27, 31	0
All	All	1783/1872 (95%)	-0.18	31 (1%) 70 66	10, 18, 26, 31	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	127	ILE	7.6
1	G	305	GLN	4.6
1	C	282	ILE	4.3
1	I	282	ILE	4.2
1	D	298	THR	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	2	6/6	0.42	0.28	42,42,42,42	0
2	GOL	A	3	6/6	0.42	0.23	47,47,47,48	0
2	GOL	C	4	6/6	0.67	0.26	63,63,63,63	0
3	IMD	A	1	5/5	0.78	0.15	55,55,55,55	0
3	IMD	C	2	5/5	0.80	0.21	47,47,47,47	0
2	GOL	H	1	6/6	0.84	0.25	32,32,32,32	6
3	IMD	A	326	5/5	0.88	0.17	63,63,63,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.