



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 07:37 pm BST

PDB ID : 6F97
Title : Crystal structure of the V465T mutant of 5-(Hydroxymethyl)furfural Oxidase (HMFO)
Authors : Pickl, M.; Swoboda, A.; Romero, E.; Winkler, C.K.; Binda, C.; Mattevi, A.; Faber, K.; Fraaije, M.W.
Deposited on : 2017-12-14
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	531	 3% 90% 8%
1	B	531	 3% 84% 7% 8%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8097 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 5-(hydroxymethyl)furfural oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	520	Total	C	N	O	S	0	1	0
			3957	2504	710	740	3			
1	B	487	Total	C	N	O	S	0	1	0
			3695	2341	662	690	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	465	THR	VAL	engineered mutation	UNP E4QP00
B	465	THR	VAL	engineered mutation	UNP E4QP00

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	B	1	53	27	9	15	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	157	Total	O	0	0
			157	157		
3	B	182	Total	O	0	0
			182	182		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5-(hydroxymethyl)furfural oxidase



- Molecule 1: 5-(hydroxymethyl)furfural oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.19Å 122.00Å 72.70Å 90.00° 91.63° 90.00°	Depositor
Resolution (Å)	72.67 – 1.90 46.72 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (72.67-1.90) 99.5 (46.72-1.90)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.172 , 0.231 0.185 , 0.243	Depositor DCC
R_{free} test set	3334 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	35.7	Xtrriage
Anisotropy	0.141	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.069 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8097	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.77	0/4057	0.95	12/5540 (0.2%)
1	B	0.78	0/3783	0.96	12/5161 (0.2%)
All	All	0.78	0/7840	0.96	24/10701 (0.2%)

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	108	ARG	NE-CZ-NH1	14.05	127.32	120.30
1	A	108	ARG	NE-CZ-NH2	-11.06	114.77	120.30
1	B	84	ARG	NE-CZ-NH1	10.48	125.54	120.30
1	A	473	ARG	NE-CZ-NH2	-8.76	115.92	120.30
1	B	519	LEU	CA-CB-CG	8.75	135.42	115.30
1	B	84	ARG	NE-CZ-NH2	-7.92	116.34	120.30
1	B	497	ARG	NE-CZ-NH2	-7.83	116.39	120.30
1	A	473	ARG	NE-CZ-NH1	7.54	124.07	120.30
1	A	485	ASP	CB-CG-OD1	7.50	125.05	118.30
1	B	519	LEU	CB-CG-CD2	7.11	123.09	111.00
1	B	497	ARG	NE-CZ-NH1	6.81	123.71	120.30
1	A	497	ARG	NE-CZ-NH1	6.38	123.49	120.30
1	A	277	ARG	NE-CZ-NH1	6.07	123.33	120.30
1	B	485	ASP	CB-CG-OD1	5.90	123.61	118.30
1	B	529	THR	N-CA-CB	5.57	120.88	110.30
1	B	115	ASP	CB-CG-OD1	5.45	123.20	118.30
1	A	479	ASP	CB-CG-OD1	5.41	123.17	118.30
1	A	75	ARG	NE-CZ-NH2	-5.31	117.65	120.30
1	B	277	ARG	NE-CZ-NH2	-5.26	117.67	120.30
1	B	405	ARG	NE-CZ-NH2	-5.24	117.68	120.30
1	A	497	ARG	NE-CZ-NH2	-5.15	117.72	120.30
1	A	220	ARG	NE-CZ-NH1	5.11	122.86	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	405	ARG	NE-CZ-NH1	5.11	122.85	120.30
1	A	405	ARG	NE-CZ-NH1	5.08	122.84	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3957	0	3912	21	0
1	B	3695	0	3664	17	0
2	A	53	0	31	4	0
2	B	53	0	31	0	0
3	A	157	0	0	6	1
3	B	182	0	0	4	1
All	All	8097	0	7638	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297:ARG:NH1	3:A:801:HOH:O	2.06	0.79
1:B:123:ASP:HB3	3:B:914:HOH:O	1.89	0.72
1:A:64:ASP:OD1	3:A:802:HOH:O	2.11	0.68
1:B:84:ARG:HD3	3:B:845:HOH:O	1.93	0.67
1:A:102:ASN:HB2	2:A:700:FAD:C4X	2.25	0.67
1:A:102:ASN:HB2	2:A:700:FAD:N5	2.10	0.66
1:A:117:TRP:HE1	1:A:506:THR:HG22	1.68	0.57
1:A:102:ASN:HB2	2:A:700:FAD:C5X	2.36	0.55
1:B:433:ARG:HE	1:B:461:ASN:HD22	1.53	0.55
1:A:60:ARG:HD2	3:A:898:HOH:O	2.08	0.53
1:B:164:HIS:HB2	3:B:879:HOH:O	2.07	0.53
1:A:102:ASN:HA	2:A:700:FAD:C6	2.39	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:433:ARG:HE	1:A:461:ASN:HD22	1.56	0.52
1:A:40:ASP:HB2	1:A:229:THR:HG21	1.90	0.52
1:B:187:GLN:HE21	1:B:199:ALA:H	1.58	0.52
1:A:108:ARG:HD3	3:A:832:HOH:O	2.12	0.50
1:B:146:ASP:HB3	3:B:829:HOH:O	2.12	0.50
1:B:148:LEU:HD12	1:B:189:ALA:HB1	1.93	0.49
1:B:30:ASN:O	1:B:224:ASN:ND2	2.41	0.49
1:A:117:TRP:NE1	1:A:506:THR:HG22	2.27	0.49
1:A:139:ARG:NH1	1:A:153:GLY:O	2.46	0.48
1:B:187:GLN:NE2	1:B:199:ALA:H	2.11	0.48
1:A:108:ARG:HD2	1:A:134:PHE:CD1	2.49	0.47
1:B:403:LEU:HD21	1:B:455:GLU:HG2	1.97	0.47
1:B:435:ALA:O	1:B:437:PRO:HD3	2.16	0.46
1:B:516:THR:HA	1:B:519:LEU:HD13	1.97	0.45
1:B:485:ASP:OD2	1:B:489:ARG:NH1	2.41	0.45
1:A:372:LYS:NZ	3:A:808:HOH:O	2.49	0.44
1:A:316:LEU:HD21	1:A:357:ALA:HB1	2.00	0.44
1:A:229:THR:CG2	3:A:872:HOH:O	2.65	0.44
1:A:187:GLN:NE2	1:A:199:ALA:H	2.17	0.43
1:B:166:SER:H	1:B:334:GLN:HE22	1.66	0.43
1:A:187:GLN:HE21	1:A:199:ALA:H	1.66	0.43
1:A:470:GLY:HA2	1:A:504:MET:O	2.19	0.42
1:B:371:ASN:HA	1:B:371:ASN:HD22	1.62	0.42
1:A:55:ARG:HG2	1:A:57:TRP:CZ2	2.55	0.42
1:B:470:GLY:HA2	1:B:504:MET:O	2.20	0.42
1:B:467:HIS:CE1	1:B:511:ASN:HA	2.56	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:946:HOH:O	3:B:946:HOH:O[2_646]	1.91	0.29

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	517/531 (97%)	503 (97%)	13 (2%)	1 (0%)	47	38
1	B	478/531 (90%)	465 (97%)	12 (2%)	1 (0%)	47	38
All	All	995/1062 (94%)	968 (97%)	25 (2%)	2 (0%)	47	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	104	VAL
1	B	436	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	409/416 (98%)	397 (97%)	12 (3%)	42	35
1	B	380/416 (91%)	367 (97%)	13 (3%)	37	28
All	All	789/832 (95%)	764 (97%)	25 (3%)	39	30

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	ASN
1	A	78	GLU
1	A	85	GLU
1	A	94	LEU
1	A	95	LEU
1	A	112	ARG
1	A	227	LEU
1	A	229	THR
1	A	254	THR
1	A	319	GLN
1	A	421	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	447	LEU
1	B	20	LEU
1	B	27	ARG
1	B	31	ARG
1	B	84	ARG
1	B	112	ARG
1	B	118	GLN
1	B	139	ARG
1	B	286	LEU
1	B	371	ASN
1	B	372	LYS
1	B	451	GLU
1	B	519	LEU
1	B	529	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	GLN
1	A	187	GLN
1	A	461	ASN
1	B	91	GLN
1	B	187	GLN
1	B	256	GLN
1	B	319	GLN
1	B	334	GLN
1	B	371	ASN
1	B	395	ASN
1	B	461	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	700	-	51,58,58	1.32	6 (11%)	60,89,89	1.92	11 (18%)
2	FAD	B	700	-	51,58,58	1.29	9 (17%)	60,89,89	1.81	12 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	700	-	-	6/30/50/50	0/6/6/6
2	FAD	B	700	-	-	4/30/50/50	0/6/6/6

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	700	FAD	C9A-C5X	3.76	1.50	1.42
2	A	700	FAD	C9A-N10	3.58	1.43	1.38
2	A	700	FAD	C4X-N5	3.46	1.38	1.33
2	B	700	FAD	C5A-C4A	3.38	1.49	1.40
2	B	700	FAD	C9A-N10	3.13	1.42	1.38
2	B	700	FAD	C10-N1	2.98	1.37	1.33
2	A	700	FAD	C4X-C10	2.61	1.41	1.38
2	B	700	FAD	C4X-C10	2.56	1.41	1.38
2	B	700	FAD	C9A-C5X	2.40	1.47	1.42
2	B	700	FAD	C4-N3	2.31	1.37	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	700	FAD	C5A-C4A	2.31	1.47	1.40
2	B	700	FAD	C2A-N3A	2.25	1.35	1.32
2	B	700	FAD	C8A-N7A	2.16	1.38	1.34
2	A	700	FAD	C5'-C4'	2.13	1.54	1.51
2	B	700	FAD	C2B-C1B	-2.01	1.50	1.53

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	700	FAD	C4-N3-C2	7.29	121.29	115.14
2	A	700	FAD	C1'-N10-C9A	6.41	123.34	118.29
2	B	700	FAD	C4-N3-C2	5.56	119.84	115.14
2	A	700	FAD	N3A-C2A-N1A	-4.89	121.04	128.68
2	B	700	FAD	N3A-C2A-N1A	-4.84	121.12	128.68
2	B	700	FAD	C4A-C5A-N7A	-4.60	104.60	109.40
2	B	700	FAD	C4X-N5-C5X	4.44	121.21	116.77
2	B	700	FAD	C1'-N10-C9A	3.77	121.26	118.29
2	A	700	FAD	O2P-P-O1P	3.54	129.75	112.24
2	A	700	FAD	C2A-N1A-C6A	2.93	123.77	118.75
2	A	700	FAD	C4X-C4-N3	-2.87	119.51	123.43
2	A	700	FAD	C4-C4X-C10	-2.76	118.12	119.95
2	A	700	FAD	C1B-N9A-C4A	-2.72	121.86	126.64
2	B	700	FAD	C2A-N1A-C6A	2.71	123.40	118.75
2	B	700	FAD	C6-C5X-N5	2.69	122.01	119.05
2	A	700	FAD	C9A-N10-C10	-2.33	118.85	121.91
2	B	700	FAD	C8M-C8-C9	-2.27	114.91	120.34
2	B	700	FAD	C4-C4X-C10	-2.23	118.47	119.95
2	B	700	FAD	C4X-C4-N3	-2.22	120.40	123.43
2	A	700	FAD	C6-C5X-N5	2.09	121.35	119.05
2	B	700	FAD	O3'-C3'-C2'	2.07	113.81	108.81
2	B	700	FAD	O4'-C4'-C5'	-2.02	105.38	109.92
2	A	700	FAD	C4A-C5A-N7A	-2.01	107.31	109.40

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	700	FAD	N10-C1'-C2'-O2'
2	A	700	FAD	O4B-C4B-C5B-O5B
2	B	700	FAD	P-O3P-PA-O1A
2	A	700	FAD	PA-O3P-P-O5'
2	B	700	FAD	PA-O3P-P-O5'

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	700	FAD	C3B-C4B-C5B-O5B
2	B	700	FAD	O4B-C4B-C5B-O5B
2	A	700	FAD	P-O3P-PA-O1A
2	A	700	FAD	P-O3P-PA-O2A
2	B	700	FAD	P-O3P-PA-O2A


There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	700	FAD	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	520/531 (97%)	0.07	18 (3%) 44 47	27, 43, 73, 104	0
1	B	487/531 (91%)	-0.05	15 (3%) 49 51	25, 40, 65, 99	0
All	All	1007/1062 (94%)	0.01	33 (3%) 46 49	25, 41, 71, 104	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	54	LEU	5.5
1	B	210	ALA	4.6
1	B	211	ALA	4.3
1	B	82	ILE	4.3
1	A	5	ILE	4.0
1	A	58	LEU	4.0
1	A	57	TRP	3.7
1	A	252	ARG	3.5
1	B	5	ILE	3.5
1	A	164	HIS	3.4
1	A	53	GLY	3.3
1	B	98	GLY	3.1
1	A	51	HIS	3.1
1	B	80	PRO	3.0
1	A	325	SER	3.0
1	A	81	GLY	3.0
1	B	203	LYS	2.9
1	B	164	HIS	2.9
1	B	202	LEU	2.9
1	B	83	THR	2.9
1	A	163	ARG	2.8
1	A	55	ARG	2.7
1	B	163	ARG	2.6
1	B	252	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	45	ASN	2.4
1	A	78	GLU	2.3
1	A	80	PRO	2.3
1	A	61	LEU	2.3
1	B	251	GLY	2.2
1	B	326	THR	2.1
1	B	81	GLY	2.0
1	A	205	GLU	2.0
1	A	52	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FAD	A	700	53/53	0.98	0.09	26,30,36,37	0
2	FAD	B	700	53/53	0.98	0.08	24,31,39,41	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD A 700:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 700:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.