



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 27, 2023 – 11:35 AM EST

PDB ID : 8F9X
Title : Cyclase-PTE
Authors : Ji, D.; Frkic, R.L.; Jackson, C.J.
Deposited on : 2022-11-24
Resolution : 2.32 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	230	 2% 82% 18%
1	B	230	 % 86% 13%
1	C	230	 4% 83% 16%
1	D	230	 % 77% 20%
1	E	230	 4% 77% 21%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	230	 3% 80% 17% **
1	G	230	 2% 80% 19% .
1	H	230	 4% 83% 16% **
1	I	230	 2% 77% 22% **
1	J	230	 % 80% 20% .

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 17461 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclase family protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	230	1708	1076	296	326	10	0	0	0
1	B	228	1695	1068	294	324	9	0	0	0
1	C	226	1683	1060	292	322	9	0	0	0
1	D	225	1679	1058	291	321	9	0	0	0
1	E	227	1691	1066	293	323	9	0	0	0
1	F	225	1679	1058	291	321	9	0	0	0
1	G	228	1695	1068	294	324	9	0	0	0
1	H	228	1695	1068	294	324	9	0	0	0
1	I	228	1695	1068	294	324	9	0	0	0
1	J	230	1708	1076	296	326	10	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q5LVE1
B	1	MET	-	initiating methionine	UNP Q5LVE1
C	1	MET	-	initiating methionine	UNP Q5LVE1
D	1	MET	-	initiating methionine	UNP Q5LVE1
E	1	MET	-	initiating methionine	UNP Q5LVE1
F	1	MET	-	initiating methionine	UNP Q5LVE1
G	1	MET	-	initiating methionine	UNP Q5LVE1
H	1	MET	-	initiating methionine	UNP Q5LVE1
I	1	MET	-	initiating methionine	UNP Q5LVE1

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
J	1	MET	-	initiating methionine	UNP Q5LVE1

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	3	Total Zn 3 3	0	0
2	B	3	Total Zn 3 3	0	0
2	C	3	Total Zn 3 3	0	0
2	D	3	Total Zn 3 3	0	0
2	E	3	Total Zn 3 3	0	0
2	F	3	Total Zn 3 3	0	0
2	G	3	Total Zn 3 3	0	0
2	H	3	Total Zn 3 3	0	0
2	I	3	Total Zn 3 3	0	0
2	J	3	Total Zn 3 3	0	0

- Molecule 3 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (three-letter code: BTB) (formula: C₈H₁₉NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	Total 14	8	1	5	0	0
3	A	1	Total 14	8	1	5	0	0
3	B	1	Total 14	8	1	5	0	0
3	C	1	Total 14	8	1	5	0	0
3	D	1	Total 14	8	1	5	0	0
3	F	1	Total 14	8	1	5	0	0
3	F	1	Total 14	8	1	5	0	0
3	G	1	Total 14	8	1	5	0	0
3	G	1	Total 14	8	1	5	0	0
3	H	1	Total 14	8	1	5	0	0
3	I	1	Total 14	8	1	5	0	0
3	I	1	Total 14	8	1	5	0	0
3	J	1	Total 14	8	1	5	0	0
3	J	1	Total 14	8	1	5	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	F	1	Total C O 4 2 2	0	0
4	F	1	Total C O 4 2 2	0	0
4	F	1	Total C O 4 2 2	0	0
4	G	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	G	1	Total	C	O	0	0
			4	2	2		
4	H	1	Total	C	O	0	0
			4	2	2		
4	H	1	Total	C	O	0	0
			4	2	2		
4	I	1	Total	C	O	0	0
			4	2	2		
4	I	1	Total	C	O	0	0
			4	2	2		
4	J	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			8	4	1	3		
5	D	1	Total	C	N	O	0	0
			8	4	1	3		
5	E	1	Total	C	N	O	0	0
			8	4	1	3		
5	H	1	Total	C	N	O	0	0
			8	4	1	3		
5	I	1	Total	C	N	O	0	0
			8	4	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
5	J	1	8	4	1	3	0	0

- Molecule 6 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	C	1	13	8	5	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	25	Total	O	0	0
			25	25		
7	B	25	Total	O	0	0
			25	25		
7	C	20	Total	O	0	0
			20	20		
7	D	16	Total	O	0	0
			16	16		
7	E	10	Total	O	0	0
			10	10		
7	F	17	Total	O	0	0
			17	17		
7	G	22	Total	O	0	0
			22	22		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	H	8	Total O 8 8	0	0
7	I	12	Total O 12 12	0	0
7	J	15	Total O 15 15	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyclase family protein




- Molecule 1: Cyclase family protein




- Molecule 1: Cyclase family protein





- Molecule 1: Cyclase family protein









● Molecule 1: Cyclase family protein





● Molecule 1: Cyclase family protein




● Molecule 1: Cyclase family protein




● Molecule 1: Cyclase family protein



- Molecule 1: Cyclase family protein



- Molecule 1: Cyclase family protein



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	68.08Å 68.08Å 446.50Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.04 – 2.32 34.04 – 2.32	Depositor EDS
% Data completeness (in resolution range)	99.9 (34.04-2.32) 99.9 (34.04-2.32)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 2.31Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.203 , 0.251 0.201 , 0.249	Depositor DCC
R_{free} test set	5012 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	49.2	Xtrriage
Anisotropy	0.328	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 37.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	0.042 for -h,-k,l 0.477 for h,-h-k,-l 0.047 for -k,-h,-l	Xtrriage
Reported twinning fraction	0.499 for H, K, L 0.501 for K, H, -L	Depositor
Outliers	0 of 100157 reflections	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	17461	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, BTB, TRS, EDO, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/1757	0.45	0/2396
1	B	0.24	0/1744	0.45	0/2379
1	C	0.24	0/1732	0.43	0/2363
1	D	0.24	0/1728	0.44	0/2358
1	E	0.25	0/1740	0.45	0/2374
1	F	0.24	0/1728	0.43	0/2358
1	G	0.24	0/1744	0.43	0/2379
1	H	0.25	0/1744	0.44	0/2379
1	I	0.25	0/1744	0.44	0/2379
1	J	0.24	0/1757	0.44	0/2396
All	All	0.24	0/17418	0.44	0/23761

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1708	0	1606	23	0
1	B	1695	0	1589	15	0
1	C	1683	0	1575	21	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1679	0	1572	35	0
1	E	1691	0	1586	24	0
1	F	1679	0	1572	23	0
1	G	1695	0	1589	29	0
1	H	1695	0	1589	24	0
1	I	1695	0	1589	28	0
1	J	1708	0	1606	23	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
2	C	3	0	0	0	0
2	D	3	0	0	0	0
2	E	3	0	0	0	0
2	F	3	0	0	0	0
2	G	3	0	0	0	0
2	H	3	0	0	0	0
2	I	3	0	0	0	0
2	J	3	0	0	0	0
3	A	28	0	38	3	0
3	B	14	0	19	1	0
3	C	14	0	19	3	0
3	D	14	0	19	0	0
3	F	28	0	38	4	0
3	G	28	0	38	4	0
3	H	14	0	19	0	0
3	I	28	0	38	3	0
3	J	28	0	38	5	0
4	A	8	0	12	0	0
4	C	12	0	18	0	0
4	D	4	0	6	0	0
4	E	12	0	18	0	0
4	F	12	0	18	0	0
4	G	8	0	12	0	0
4	H	8	0	12	0	0
4	I	8	0	12	0	0
4	J	4	0	6	0	0
5	B	8	0	12	0	0
5	D	8	0	12	1	0
5	E	8	0	12	0	0
5	H	8	0	12	0	0
5	I	8	0	12	0	0
5	J	8	0	12	0	0
6	C	13	0	18	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	25	0	0	0	0
7	B	25	0	0	0	0
7	C	20	0	0	0	0
7	D	16	0	0	0	0
7	E	10	0	0	0	0
7	F	17	0	0	0	0
7	G	22	0	0	0	0
7	H	8	0	0	0	0
7	I	12	0	0	0	0
7	J	15	0	0	0	0
All	All	17461	0	16343	237	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 237 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:26:GLY:HA3	1:G:51:THR:HG21	1.64	0.78
1:A:113:PRO:HG3	1:A:159:THR:HA	1.74	0.70
1:G:134:ARG:NH1	1:G:142:MET:SD	2.66	0.69
1:B:141:LYS:HE2	1:B:175:PRO:HB3	1.74	0.69
1:A:20:THR:O	1:A:134:ARG:NH1	2.26	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	228/230 (99%)	210 (92%)	17 (8%)	1 (0%)	34 41
1	B	226/230 (98%)	209 (92%)	17 (8%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	224/230 (97%)	204 (91%)	20 (9%)	0	100	100
1	D	223/230 (97%)	205 (92%)	18 (8%)	0	100	100
1	E	225/230 (98%)	200 (89%)	22 (10%)	3 (1%)	12	12
1	F	223/230 (97%)	209 (94%)	13 (6%)	1 (0%)	34	41
1	G	226/230 (98%)	208 (92%)	18 (8%)	0	100	100
1	H	226/230 (98%)	214 (95%)	11 (5%)	1 (0%)	34	41
1	I	226/230 (98%)	211 (93%)	13 (6%)	2 (1%)	17	19
1	J	228/230 (99%)	207 (91%)	20 (9%)	1 (0%)	34	41
All	All	2255/2300 (98%)	2077 (92%)	169 (8%)	9 (0%)	34	41

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	114	ASP
1	E	87	GLU
1	E	140	GLY
1	H	138	SER
1	I	208	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/172 (100%)	169 (98%)	3 (2%)	60	75
1	B	171/172 (99%)	165 (96%)	6 (4%)	36	49
1	C	170/172 (99%)	162 (95%)	8 (5%)	26	36
1	D	170/172 (99%)	166 (98%)	4 (2%)	49	65
1	E	171/172 (99%)	159 (93%)	12 (7%)	15	19
1	F	170/172 (99%)	163 (96%)	7 (4%)	30	43
1	G	171/172 (99%)	167 (98%)	4 (2%)	50	66
1	H	171/172 (99%)	166 (97%)	5 (3%)	42	57

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	I	171/172 (99%)	166 (97%)	5 (3%)	42 57
1	J	172/172 (100%)	167 (97%)	5 (3%)	42 57
All	All	1709/1720 (99%)	1650 (96%)	59 (4%)	36 49

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	87	GLU
1	J	7	VAL
1	F	112	ILE
1	J	1	MET
1	I	84	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	135	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 70 ligands modelled in this entry, 30 are monoatomic - leaving 40 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	H	306	-	3,3,3	0.45	0	2,2,2	0.34	0
3	BTB	J	306	-	13,13,13	0.80	0	7,16,16	0.68	0
5	TRS	B	304	-	7,7,7	0.34	0	9,9,9	0.34	0
4	EDO	E	305	-	3,3,3	0.46	0	2,2,2	0.34	0
4	EDO	C	307	-	3,3,3	0.46	0	2,2,2	0.34	0
3	BTB	I	305	-	13,13,13	0.76	0	7,16,16	0.69	0
3	BTB	H	305	-	13,13,13	0.78	0	7,16,16	0.66	0
5	TRS	I	304	-	7,7,7	0.33	0	9,9,9	0.34	0
3	BTB	F	304	-	13,13,13	0.76	0	7,16,16	0.76	0
5	TRS	D	304	-	7,7,7	0.33	0	9,9,9	0.34	0
3	BTB	G	304	-	13,13,13	0.79	0	7,16,16	0.64	0
4	EDO	F	308	-	3,3,3	0.46	0	2,2,2	0.33	0
3	BTB	G	305	-	13,13,13	0.78	0	7,16,16	0.67	0
4	EDO	C	305	-	3,3,3	0.46	0	2,2,2	0.33	0
4	EDO	D	306	-	3,3,3	0.46	0	2,2,2	0.34	0
4	EDO	I	307	-	3,3,3	0.46	0	2,2,2	0.34	0
3	BTB	D	305	-	13,13,13	0.79	0	7,16,16	0.68	0
3	BTB	J	305	-	13,13,13	0.79	0	7,16,16	0.62	0
4	EDO	F	307	-	3,3,3	0.46	0	2,2,2	0.34	0
3	BTB	B	305	-	13,13,13	0.81	0	7,16,16	0.60	0
4	EDO	G	307	-	3,3,3	0.45	0	2,2,2	0.35	0
3	BTB	I	308	-	13,13,13	0.78	0	7,16,16	0.67	0
5	TRS	H	304	-	7,7,7	0.33	0	9,9,9	0.34	0
4	EDO	A	306	-	3,3,3	0.46	0	2,2,2	0.34	0
6	PG4	C	308	-	12,12,12	0.11	0	11,11,11	0.68	0
4	EDO	I	306	-	3,3,3	0.46	0	2,2,2	0.34	0
3	BTB	A	305	-	13,13,13	0.77	0	7,16,16	0.63	0
3	BTB	A	304	-	13,13,13	0.79	0	7,16,16	0.64	0
4	EDO	E	306	-	3,3,3	0.45	0	2,2,2	0.33	0
4	EDO	J	307	-	3,3,3	0.45	0	2,2,2	0.34	0
4	EDO	F	306	-	3,3,3	0.46	0	2,2,2	0.34	0
5	TRS	E	304	-	7,7,7	0.33	0	9,9,9	0.34	0
3	BTB	F	305	-	13,13,13	0.78	0	7,16,16	0.65	0
4	EDO	G	306	-	3,3,3	0.46	0	2,2,2	0.34	0
4	EDO	A	307	-	3,3,3	0.46	0	2,2,2	0.34	0
5	TRS	J	304	-	7,7,7	0.33	0	9,9,9	0.34	0
4	EDO	E	307	-	3,3,3	0.46	0	2,2,2	0.32	0
4	EDO	H	307	-	3,3,3	0.46	0	2,2,2	0.35	0
4	EDO	C	306	-	3,3,3	0.45	0	2,2,2	0.34	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BTB	C	304	-	13,13,13	0.79	0	7,16,16	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	H	306	-	-	0/1/1/1	-
3	BTB	J	306	-	-	12/21/21/21	-
5	TRS	B	304	-	-	0/9/9/9	-
4	EDO	E	305	-	-	0/1/1/1	-
4	EDO	C	307	-	-	0/1/1/1	-
3	BTB	I	305	-	-	7/21/21/21	-
3	BTB	H	305	-	-	2/21/21/21	-
5	TRS	I	304	-	-	0/9/9/9	-
3	BTB	F	304	-	-	4/21/21/21	-
5	TRS	D	304	-	-	0/9/9/9	-
3	BTB	G	304	-	-	11/21/21/21	-
4	EDO	F	308	-	-	0/1/1/1	-
3	BTB	G	305	-	-	8/21/21/21	-
4	EDO	C	305	-	-	0/1/1/1	-
4	EDO	D	306	-	-	0/1/1/1	-
4	EDO	I	307	-	-	0/1/1/1	-
3	BTB	D	305	-	-	10/21/21/21	-
3	BTB	J	305	-	-	7/21/21/21	-
4	EDO	F	307	-	-	0/1/1/1	-
3	BTB	B	305	-	-	13/21/21/21	-
4	EDO	G	307	-	-	0/1/1/1	-
3	BTB	I	308	-	-	5/21/21/21	-
5	TRS	H	304	-	-	0/9/9/9	-
4	EDO	A	306	-	-	0/1/1/1	-
6	PG4	C	308	-	-	8/10/10/10	-
4	EDO	I	306	-	-	0/1/1/1	-
3	BTB	A	305	-	-	11/21/21/21	-
3	BTB	A	304	-	-	9/21/21/21	-
4	EDO	E	306	-	-	0/1/1/1	-
4	EDO	J	307	-	-	0/1/1/1	-
4	EDO	F	306	-	-	0/1/1/1	-
5	TRS	E	304	-	-	0/9/9/9	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BTB	F	305	-	-	10/21/21/21	-
4	EDO	G	306	-	-	0/1/1/1	-
4	EDO	A	307	-	-	0/1/1/1	-
5	TRS	J	304	-	-	0/9/9/9	-
4	EDO	E	307	-	-	0/1/1/1	-
4	EDO	H	307	-	-	0/1/1/1	-
4	EDO	C	306	-	-	0/1/1/1	-
3	BTB	C	304	-	-	11/21/21/21	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 128 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	304	BTB	C1-C2-C4-O4
3	A	304	BTB	C1-C2-N-C7
3	A	304	BTB	C3-C2-N-C7
3	A	304	BTB	C4-C2-N-C7
3	A	305	BTB	C4-C2-C3-O3

There are no ring outliers.

12 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	J	306	BTB	2	0
3	I	305	BTB	2	0
3	F	304	BTB	2	0
5	D	304	TRS	1	0
3	G	304	BTB	3	0
3	G	305	BTB	1	0
3	J	305	BTB	3	0
3	B	305	BTB	1	0
3	I	308	BTB	1	0
3	A	305	BTB	3	0
3	F	305	BTB	2	0
3	C	304	BTB	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	230/230 (100%)	0.03	4 (1%) 70 76	41, 56, 84, 200	0
1	B	228/230 (99%)	0.02	2 (0%) 84 88	33, 58, 88, 155	0
1	C	226/230 (98%)	0.12	9 (3%) 38 45	43, 63, 108, 139	0
1	D	225/230 (97%)	0.08	3 (1%) 77 81	45, 65, 96, 132	0
1	E	227/230 (98%)	0.17	9 (3%) 38 45	46, 69, 107, 162	0
1	F	225/230 (97%)	0.18	7 (3%) 49 56	45, 66, 95, 127	0
1	G	228/230 (99%)	0.01	4 (1%) 68 75	41, 64, 92, 149	0
1	H	228/230 (99%)	0.22	10 (4%) 34 41	45, 64, 100, 187	0
1	I	228/230 (99%)	0.04	5 (2%) 62 69	39, 62, 95, 152	0
1	J	230/230 (100%)	-0.07	2 (0%) 84 88	45, 61, 90, 114	0
All	All	2275/2300 (98%)	0.08	55 (2%) 59 66	33, 62, 98, 200	0

The worst 5 of 55 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	107	SER	7.7
1	C	43	PHE	6.0
1	E	93	ALA	5.9
1	I	230	VAL	4.9
1	B	217	ARG	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	D	303	1/1	0.09	0.13	126,126,126,126	0
4	EDO	E	307	4/4	0.55	0.23	102,107,110,118	0
4	EDO	G	306	4/4	0.62	0.16	62,70,75,77	0
2	ZN	D	302	1/1	0.77	0.15	73,73,73,73	0
2	ZN	C	301	1/1	0.83	0.18	63,63,63,63	0
4	EDO	F	306	4/4	0.83	0.11	62,64,66,69	0
4	EDO	E	306	4/4	0.83	0.29	67,79,86,91	0
4	EDO	H	307	4/4	0.83	0.16	98,99,101,102	0
6	PG4	C	308	13/13	0.83	0.14	58,63,66,70	0
2	ZN	I	303	1/1	0.85	0.07	109,109,109,109	0
3	BTB	I	308	14/14	0.86	0.12	60,67,74,75	0
5	TRS	I	304	8/8	0.87	0.14	54,62,64,66	0
5	TRS	D	304	8/8	0.88	0.10	69,74,80,80	0
3	BTB	F	305	14/14	0.88	0.12	62,71,75,80	0
5	TRS	J	304	8/8	0.88	0.20	62,69,79,83	0
3	BTB	H	305	14/14	0.88	0.15	59,67,78,79	0
4	EDO	C	305	4/4	0.90	0.17	55,69,76,84	0
4	EDO	C	307	4/4	0.90	0.12	60,64,66,72	0
4	EDO	E	305	4/4	0.90	0.14	49,52,53,55	0
2	ZN	E	303	1/1	0.90	0.09	98,98,98,98	0
3	BTB	F	304	14/14	0.91	0.12	66,75,83,88	0
4	EDO	J	307	4/4	0.91	0.14	47,51,58,61	0
3	BTB	J	305	14/14	0.91	0.16	49,54,68,69	0
4	EDO	A	306	4/4	0.91	0.11	56,58,59,62	0
2	ZN	D	301	1/1	0.91	0.14	71,71,71,71	0
3	BTB	A	304	14/14	0.91	0.09	60,68,72,73	0
4	EDO	F	307	4/4	0.92	0.10	52,53,57,60	0
3	BTB	C	304	14/14	0.92	0.13	51,61,68,70	0
4	EDO	C	306	4/4	0.92	0.11	55,61,70,72	0
3	BTB	I	305	14/14	0.92	0.13	57,61,69,69	0
3	BTB	D	305	14/14	0.92	0.13	53,60,62,67	0
5	TRS	E	304	8/8	0.92	0.09	74,85,92,93	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	F	301	1/1	0.92	0.11	66,66,66,66	0
3	BTB	J	306	14/14	0.92	0.11	60,70,79,79	0
3	BTB	B	305	14/14	0.92	0.10	55,63,66,67	0
3	BTB	A	305	14/14	0.93	0.10	52,56,66,67	0
3	BTB	G	305	14/14	0.93	0.09	51,59,64,66	0
4	EDO	I	307	4/4	0.93	0.12	48,55,58,60	0
5	TRS	B	304	8/8	0.94	0.13	51,59,67,74	0
4	EDO	D	306	4/4	0.94	0.10	45,50,51,60	0
3	BTB	G	304	14/14	0.94	0.14	57,64,67,68	0
4	EDO	I	306	4/4	0.95	0.13	51,55,56,58	0
4	EDO	A	307	4/4	0.95	0.12	49,49,51,52	0
2	ZN	C	303	1/1	0.95	0.16	58,58,58,58	0
4	EDO	G	307	4/4	0.96	0.12	49,56,60,60	0
2	ZN	A	301	1/1	0.96	0.14	51,51,51,51	0
5	TRS	H	304	8/8	0.96	0.12	60,63,67,68	0
4	EDO	F	308	4/4	0.97	0.08	42,43,45,55	0
2	ZN	F	302	1/1	0.97	0.12	59,59,59,59	0
2	ZN	C	302	1/1	0.97	0.15	53,53,53,53	0
4	EDO	H	306	4/4	0.97	0.12	56,59,61,68	0
2	ZN	E	301	1/1	0.97	0.11	65,65,65,65	0
2	ZN	E	302	1/1	0.98	0.12	51,51,51,51	0
2	ZN	J	303	1/1	0.98	0.17	42,42,42,42	0
2	ZN	B	301	1/1	0.98	0.13	38,38,38,38	0
2	ZN	B	303	1/1	0.98	0.14	57,57,57,57	0
2	ZN	A	303	1/1	0.98	0.13	53,53,53,53	0
2	ZN	G	303	1/1	0.98	0.11	60,60,60,60	0
2	ZN	H	301	1/1	0.98	0.15	46,46,46,46	0
2	ZN	I	302	1/1	0.99	0.20	53,53,53,53	0
2	ZN	G	302	1/1	0.99	0.14	51,51,51,51	0
2	ZN	J	301	1/1	0.99	0.14	38,38,38,38	0
2	ZN	J	302	1/1	0.99	0.12	40,40,40,40	0
2	ZN	A	302	1/1	0.99	0.12	49,49,49,49	0
2	ZN	F	303	1/1	0.99	0.12	45,45,45,45	0
2	ZN	H	302	1/1	0.99	0.10	57,57,57,57	0
2	ZN	H	303	1/1	0.99	0.11	71,71,71,71	0
2	ZN	I	301	1/1	0.99	0.13	52,52,52,52	0
2	ZN	G	301	1/1	1.00	0.13	50,50,50,50	0
2	ZN	B	302	1/1	1.00	0.12	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.