



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 11, 2023 – 10:04 AM EST

PDB ID : 8FBW  
Title : Crystal structure of SIV-1 V2 antibody NCI05 in complex with a V2 peptide  
Authors : Zhou, T.; Kwong, P.D.; Van Wazer, D.  
Deposited on : 2022-11-30  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.31.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	
1	C	231	
2	B	216	
2	D	216	
3	E	17	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	17	<p>76% 59% 24% 6% 12%</p>
4	G	4	<p>50% 25% 25%</p>
5	H	2	<p>100%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	H	2	-	-	-	X

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 7319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heavy chain of anti-SIV V2 antibody NCI05.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	231	Total	C	N	O	S	0	0	0
			1745	1103	290	347	5			
1	A	231	Total	C	N	O	S	0	0	0
			1745	1103	290	347	5			


- Molecule 2 is a protein called Light chain of anti-SIV V2 antibody NCI05.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	216	Total	C	N	O	S	0	0	0
			1596	991	268	332	5			
2	B	214	Total	C	N	O	S	0	0	0
			1583	985	266	328	4			

- Molecule 3 is a protein called SIV V2 peptide.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	16	Total	C	N	O	0	0	0
			145	93	24	28			
3	F	15	Total	C	N	O	0	0	0
			134	87	20	27			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	4	Total	C	N	O	0	0	0
			49	28	2	19			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	H	2	28	16	2	10	0	0	0

- Molecule 6 is CITRIC ACID (three-letter code: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	C	1	13	6	7	0	0

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	D	1	Total	O	P	0	0
			5	4	1		
8	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	C	109	Total O 109 109	0	0
9	D	69	Total O 69 69	0	0
9	E	9	Total O 9 9	0	0
9	A	48	Total O 48 48	0	0
9	B	28	Total O 28 28	0	0
9	F	1	Total O 1 1	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heavy chain of anti-SIV V2 antibody NCI05





- Molecule 1: Heavy chain of anti-SIV V2 antibody NCI05



- Molecule 2: Light chain of anti-SIV V2 antibody NCI05





- Molecule 2: Light chain of anti-SIV V2 antibody NCI05





- Molecule 3: SIV V2 peptide







- Molecule 3: SIV V2 peptide



- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	130.65Å 130.65Å 216.95Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.91 – 2.35 48.91 – 2.35	Depositor EDS
% Data completeness (in resolution range)	98.6 (48.91-2.35) 98.6 (48.91-2.35)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.15 (at 2.34Å)	Xtrriage
Refinement program	PHENIX 1.20.1-4487-000	Depositor
R, $R_{free}$	0.194 , 0.246 0.191 , 0.245	Depositor DCC
$R_{free}$ test set	2000 reflections (4.39%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.1	Xtrriage
Anisotropy	0.058	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 40.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7319	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, CIT, FUC, PO4, NAG, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/1790	0.52	0/2441
1	C	0.26	0/1790	0.51	0/2441
2	B	0.26	0/1620	0.47	0/2208
2	D	0.27	0/1633	0.48	0/2224
3	E	0.26	0/148	0.47	0/197
3	F	0.22	0/137	0.38	0/183
All	All	0.26	0/7118	0.49	0/9694

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1745	0	1684	20	0
1	C	1745	0	1684	9	0
2	B	1583	0	1524	22	0
2	D	1596	0	1534	10	0
3	E	145	0	140	2	0
3	F	134	0	127	5	0
4	G	49	0	43	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	H	28	0	25	1	0
6	C	13	0	5	0	0
7	C	7	0	10	0	0
8	A	5	0	0	0	0
8	D	5	0	0	0	0
9	A	48	0	0	0	0
9	B	28	0	0	2	0
9	C	109	0	0	0	0
9	D	69	0	0	0	0
9	E	9	0	0	0	0
9	F	1	0	0	0	0
All	All	7319	0	6776	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:52:SER:OG	3:F:182:GLU:OE2	2.05	0.74
1:A:39:GLN:HE22	2:B:39:GLN:HE22	1.34	0.72
1:A:29:PHE:O	1:A:71:ARG:NH2	2.24	0.70
2:B:40:LEU:HD23	2:B:85:ALA:HB2	1.75	0.69
2:B:36:TRP:HB2	2:B:49:ILE:HB	1.76	0.68
1:A:135:PRO:HB3	1:A:161:TYR:HB3	1.77	0.66
2:B:175:LYS:NZ	9:B:302:HOH:O	2.31	0.64
1:A:17:SER:HA	1:A:82:MET:O	1.97	0.62
2:B:112:GLN:HG2	2:B:113:PRO:HD2	1.82	0.62
2:B:136:LEU:HD12	2:B:182:LEU:HD23	1.83	0.61
2:B:191:SER:OG	2:B:192:HIS:N	2.35	0.57
1:C:142:PRO:HD2	1:C:229:ILE:HG12	1.85	0.56
1:A:90:THR:HG23	1:A:126:THR:HA	1.87	0.56
2:B:140:ILE:HG12	2:B:199:VAL:HG21	1.87	0.55
1:A:51:ILE:HG12	1:A:71:ARG:HB3	1.87	0.55
1:A:93:TYR:O	1:A:122:GLY:HA2	2.08	0.54
2:D:40:LEU:HD23	2:D:85:ALA:HB2	1.90	0.53
4:G:1:NAG:H83	4:G:1:NAG:H3	1.89	0.53
2:B:28:ASN:OD1	2:B:29:ILE:N	2.28	0.53
1:A:40:ALA:HB3	1:A:43:LYS:HG3	1.90	0.53
2:B:13:ALA:HB3	2:B:16:GLN:HG3	1.91	0.53
2:D:1:GLN:N	2:D:93:ASP:OD2	2.44	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:36:TRP:CZ3	2:B:89:CYS:HB3	2.47	0.50
2:B:17:SER:HA	2:B:76:ILE:O	2.12	0.50
2:D:38:GLN:HB2	2:D:48:LEU:HD22	1.94	0.49
2:B:196:SER:HG	2:B:209:THR:HG1	1.59	0.49
1:C:70:SER:O	1:C:78:LEU:HD12	2.13	0.49
3:E:178:ASP:OD1	3:E:180:LYS:HE2	2.13	0.49
2:B:186:SER:O	2:B:190:LYS:HG3	2.12	0.49
2:B:187:ASP:O	2:B:191:SER:HB3	2.13	0.49
2:B:136:LEU:HD21	2:B:189:TRP:CZ3	2.48	0.49
2:B:136:LEU:HB2	2:B:182:LEU:HB3	1.96	0.48
5:H:1:NAG:H62	5:H:2:NAG:N2	2.28	0.48
1:A:58:TYR:HE2	3:F:176:LYS:HD2	1.77	0.48
1:A:58:TYR:CE2	3:F:176:LYS:HD2	2.51	0.46
2:B:133:LYS:HE2	2:B:133:LYS:HB3	1.80	0.45
1:A:96:ALA:HB1	1:A:116:PHE:HB3	1.98	0.45
1:C:104:TYR:OH	1:C:107:GLY:HA2	2.17	0.45
2:B:189:TRP:CZ2	2:B:212:PRO:HA	2.52	0.45
1:C:3:GLN:NE2	1:A:151:THR:OG1	2.50	0.44
1:C:209:THR:HB	1:C:226:ARG:HD2	1.99	0.44
1:A:38:ARG:HG2	1:A:48:VAL:HG22	1.98	0.44
1:A:58:TYR:CE1	3:F:179:LYS:HA	2.52	0.44
1:C:82:MET:HB3	1:C:85:LEU:HD21	1.98	0.44
1:A:109:TRP:CD1	3:F:187:TRP:HB3	2.53	0.44
2:B:34:VAL:HG22	2:B:52:ASN:HA	1.99	0.44
2:D:124:PRO:HG3	2:D:134:ALA:HB1	2.00	0.44
2:D:193:LYS:HD2	2:D:193:LYS:HA	1.77	0.44
1:A:71:ARG:NE	1:A:73:ASN:OD1	2.38	0.43
1:A:78:LEU:HD23	1:A:79:SER:N	2.33	0.43
1:C:201:PRO:O	1:C:204:SER:OG	2.32	0.43
1:A:229:ILE:HD12	1:A:229:ILE:HA	1.78	0.43
1:A:51:ILE:HD11	1:A:55:GLY:HA2	2.01	0.42
1:C:17:SER:HA	1:C:82:MET:O	2.19	0.42
2:D:93:ASP:HB2	2:D:100:LEU:HD22	2.02	0.42
2:D:17:SER:HA	2:D:76:ILE:O	2.20	0.42
3:E:180:LYS:HE3	3:E:180:LYS:HB2	1.78	0.42
2:B:39:GLN:NE2	9:B:308:HOH:O	2.51	0.42
1:C:183:PRO:HG2	2:D:169:SER:OG	2.21	0.41
2:D:36:TRP:HB2	2:D:49:ILE:HB	2.03	0.41
2:B:93:ASP:OD1	2:B:95:SER:OG	2.31	0.41
2:D:214:GLU:C	2:D:216:SER:H	2.25	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/231 (99%)	221 (96%)	6 (3%)	2 (1%)	17	17
1	C	229/231 (99%)	229 (100%)	0	0	100	100
2	B	212/216 (98%)	203 (96%)	9 (4%)	0	100	100
2	D	214/216 (99%)	208 (97%)	5 (2%)	1 (0%)	29	32
3	E	14/17 (82%)	13 (93%)	1 (7%)	0	100	100
3	F	13/17 (76%)	12 (92%)	1 (8%)	0	100	100
All	All	911/928 (98%)	886 (97%)	22 (2%)	3 (0%)	41	47

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	145	ARG
1	A	106	GLY
2	D	2	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/195 (100%)	186 (95%)	9 (5%)	27	32
1	C	195/195 (100%)	192 (98%)	3 (2%)	65	76
2	B	177/179 (99%)	172 (97%)	5 (3%)	43	53
2	D	179/179 (100%)	173 (97%)	6 (3%)	37	46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	E	16/17 (94%)	16 (100%)	0	100	100
3	F	15/17 (88%)	13 (87%)	2 (13%)	4	4
All	All	777/782 (99%)	752 (97%)	25 (3%)	39	47

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	25	SER
1	C	147	THR
1	C	187	GLN
2	D	8	PRO
2	D	41	SER
2	D	160	ASN
2	D	170	LYS
2	D	174	ASN
2	D	215	CYS
1	A	25	SER
1	A	28	THR
1	A	30	SER
1	A	31	ARG
1	A	45	LEU
1	A	64	GLU
1	A	180	HIS
1	A	213	ASN
1	A	222	LYS
2	B	41	SER
2	B	66	SER
2	B	71	SER
2	B	179	SER
2	B	191	SER
3	F	176	LYS
3	F	177	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	3	GLN
1	C	187	GLN
2	B	32	ASN
2	B	39	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	G	1	4,1	14,14,15	0.44	0	17,19,21	1.33	2 (11%)
4	NAG	G	2	4	14,14,15	0.25	0	17,19,21	0.37	0
4	BMA	G	3	4	11,11,12	0.62	0	15,15,17	0.74	0
4	FUC	G	4	4	10,10,11	0.65	0	14,14,16	1.00	1 (7%)
5	NAG	H	1	1,5	14,14,15	0.29	0	17,19,21	0.42	0
5	NAG	H	2	5	14,14,15	0.25	0	17,19,21	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	4,1	-	3/6/23/26	0/1/1/1
4	NAG	G	2	4	-	1/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
4	FUC	G	4	4	-	-	0/1/1/1
5	NAG	H	1	1,5	-	3/6/23/26	0/1/1/1
5	NAG	H	2	5	-	2/6/23/26	0/1/1/1

There are no bond length outliers.



All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	1	NAG	C2-N2-C7	4.25	128.95	122.90
4	G	4	FUC	C1-O5-C5	2.21	117.78	112.78
4	G	1	NAG	C1-C2-N2	2.00	113.91	110.49

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	G	1	NAG	C8-C7-N2-C2
4	G	1	NAG	O7-C7-N2-C2
5	H	1	NAG	C8-C7-N2-C2
5	H	1	NAG	O7-C7-N2-C2
5	H	2	NAG	C8-C7-N2-C2
5	H	2	NAG	O7-C7-N2-C2
4	G	2	NAG	C4-C5-C6-O6
4	G	1	NAG	C3-C2-N2-C7
5	H	1	NAG	C4-C5-C6-O6


There are no ring outliers.

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	1	NAG	1	0
5	H	1	NAG	1	0
5	H	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	CIT	C	301	-	12,12,12	1.06	0	17,17,17	1.57	2 (11%)
8	PO4	A	301	-	4,4,4	0.90	0	6,6,6	0.42	0
8	PO4	D	301	-	4,4,4	0.92	0	6,6,6	0.42	0
7	PEG	C	302	-	6,6,6	0.10	0	5,5,5	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	CIT	C	301	-	-	1/16/16/16	-
7	PEG	C	302	-	-	2/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	301	CIT	O6-C6-C3	3.90	119.83	113.05
6	C	301	CIT	O4-C5-C4	2.11	121.13	114.35

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	C	302	PEG	O2-C3-C4-O4
7	C	302	PEG	C1-C2-O2-C3
6	C	301	CIT	C4-C3-C6-O5

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	231/231 (100%)	0.55	17 (7%) 14 22	31, 58, 112, 160	0
1	C	231/231 (100%)	0.23	4 (1%) 70 78	26, 37, 70, 142	0
2	B	214/216 (99%)	0.67	18 (8%) 11 16	36, 69, 110, 146	0
2	D	216/216 (100%)	0.14	6 (2%) 53 64	30, 46, 64, 148	0
3	E	16/17 (94%)	1.70	4 (25%) 0 1	42, 53, 118, 123	0
3	F	15/17 (88%)	5.46	13 (86%) 0 0	108, 119, 137, 144	0
All	All	923/928 (99%)	0.50	62 (6%) 17 26	26, 49, 112, 160	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	216	SER	13.3
1	C	231	THR	12.7
2	B	1	GLN	12.5
2	D	215	CYS	11.8
3	F	175	LEU	11.6
3	F	188	TYR	10.6
1	A	108	TYR	9.8
1	A	107	GLY	9.1
3	F	185	GLU	8.6
3	F	189	SER	8.4
1	A	231	THR	7.7
3	E	190	ARG	7.6
2	B	2	SER	7.1
3	F	186	THR	7.1
3	E	175	LEU	6.9
2	D	1	GLN	6.6
3	E	189	SER	6.6
1	A	147	THR	5.8
2	D	2	SER	5.7

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	106	GLY	5.3
2	B	67	LYS	5.2
3	F	184	ASN	5.1
2	B	3	VAL	5.1
1	A	143	SER	4.8
3	F	177	SER	4.6
2	B	214	GLU	4.4
1	A	104	TYR	4.4
3	F	187	TRP	4.3
3	F	183	TYR	4.3
1	A	109	TRP	4.2
3	F	178	ASP	4.0
1	A	144	SER	4.0
3	F	179	LYS	3.9
2	B	213	ALA	3.9
1	A	110	TYR	3.7
3	F	181	ILE	3.5
2	B	61	ASP	3.4
1	C	144	SER	3.4
1	A	111	VAL	3.2
2	D	3	VAL	3.1
2	B	95	SER	3.1
2	B	34	VAL	3.0
2	B	94	SER	2.9
1	A	56	SER	2.9
3	F	176	LYS	2.9
2	B	27	SER	2.8
1	A	58	TYR	2.8
1	C	145	ARG	2.8
3	E	185	GLU	2.8
1	A	102	TYR	2.7
2	B	160	ASN	2.6
2	B	65	GLY	2.5
1	A	103	TYR	2.5
1	A	114	TYR	2.5
2	B	96	LEU	2.5
2	B	100	LEU	2.2
1	C	146	SER	2.2
2	B	63	PHE	2.2
2	D	213	ALA	2.2
2	B	35	TYR	2.1
2	B	68	SER	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	118	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAG	H	2	14/15	0.60	0.41	103,117,122,123	0
4	BMA	G	3	11/12	0.64	0.19	75,83,87,92	0
5	NAG	H	1	14/15	0.68	0.22	99,111,116,121	0
4	NAG	G	2	14/15	0.86	0.17	52,63,70,80	0
4	NAG	G	1	14/15	0.89	0.15	45,53,65,65	0
4	FUC	G	4	10/11	0.92	0.18	48,56,58,59	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
8	PO4	A	301	5/5	0.82	0.55	52,75,103,121	0
6	CIT	C	301	13/13	0.87	0.25	45,54,63,67	0
8	PO4	D	301	5/5	0.88	0.36	77,80,91,106	0
7	PEG	C	302	7/7	0.92	0.15	52,58,70,74	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.