

Full wwPDB NMR Structure Validation Report (i)

Feb 12, 2022 – 03:10 PM EST

PDB ID : 1FH3

Title : NMR STRUCTURES OF LQH III ALPHA-LIKE SCORPION TOXIN FROM

LEIURUS QUINQUESTRIATUS CORRESPONDING TO THE MAJOR

CONFORMER IN SOLUTION

Authors: Krimm, I.; Trivelli, X.; Lancelin, J.M.

Deposited on : 2000-07-31

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

ShiftChecker : 2.26

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.26

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$rac{ m NMR~archive}{ m (\#Entries)}$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain			
1	A	68	65%	24%	12%	



2 Ensemble composition and analysis (i)

This entry contains 42 models. Model 30 is the overall representative, medoid model (most similar to other models). The authors have identified model 25 as representative, based on the following criterion: closest to the average.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues							
Well-defined core	Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model						
1	1 A:1-A:60 (60) 0.50 30						

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 8 single-model clusters were found.

Cluster number	Models
1	1, 3, 5, 6, 8, 9, 10, 11, 12, 17, 19, 23, 25, 30, 34,
1	36, 38, 40, 42
2	16, 20, 26, 27, 32, 39
3	2, 13, 21, 37
4	7, 14, 35
5	4, 24
Single-model clusters	15; 18; 22; 28; 29; 31; 33; 41



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 945 atoms, of which 455 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called LQH III ALPHA-LIKE TOXIN.

Mol	Chain	Residues	Atoms					Trace	
1	Λ	69	Total	С	Н	N	О	S	1
	A	68	945	301	455	88	93	8	1



4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: LQH III ALPHA-LIKE TOXIN



4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.2 Score per residue for model 2






4.2.3 Score per residue for model 3

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.4 Score per residue for model 4

• Molecule 1: LQH III ALPHA-LIKE TOXIN



4.2.5 Score per residue for model 5

• Molecule 1: LQH III ALPHA-LIKE TOXIN



4.2.6 Score per residue for model 6

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.7 Score per residue for model 7






4.2.8 Score per residue for model 8

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.9 Score per residue for model 9

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.10 Score per residue for model 10

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.11 Score per residue for model 11

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.12 Score per residue for model 12





4.2.13 Score per residue for model 13

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.14 Score per residue for model 14

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.15 Score per residue for model 15

• Molecule 1: LQH III ALPHA-LIKE TOXIN



4.2.16 Score per residue for model 16

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.17 Score per residue for model 17






4.2.18 Score per residue for model 18

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.19 Score per residue for model 19

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.20 Score per residue for model 20

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.21 Score per residue for model 21

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.22 Score per residue for model 22






4.2.23 Score per residue for model 23

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.24 Score per residue for model 24

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.25 Score per residue for model 25

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.26 Score per residue for model 26

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.27 Score per residue for model 27




4.2.28 Score per residue for model 28

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.29 Score per residue for model 29

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.30 Score per residue for model 30 (medoid)

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.31 Score per residue for model 31

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.32 Score per residue for model 32




4.2.33 Score per residue for model 33

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.34 Score per residue for model 34

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.35 Score per residue for model 35

• Molecule 1: LQH III ALPHA-LIKE TOXIN



4.2.36 Score per residue for model 36

• Molecule 1: LQH III ALPHA-LIKE TOXIN



4.2.37 Score per residue for model 37






4.2.38 Score per residue for model 38

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.39 Score per residue for model 39

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.40 Score per residue for model 40

• Molecule 1: LQH III ALPHA-LIKE TOXIN




4.2.41 Score per residue for model 41

• Molecule 1: LQH III ALPHA-LIKE TOXIN



4.2.42 Score per residue for model 42





5 Refinement protocol and experimental data overview (i)



The models were refined using the following method: distance geometry simulated annealing molecular dynamics.

Of the 80 calculated structures, 42 were deposited, based on the following criterion: structures with acceptable covalent geometry, structures with favorable non-bond energy, structures with the least restraint violations, structures with the lowest energy.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	3.851
X-PLOR	refinement	3.851

No chemical shift data was provided.



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles		
MIOI	Chain	RMSZ	#Z>5	RMSZ	#Z>5	
1	A	0.95 ± 0.01	$0\pm0/448~(~0.0\pm~0.0\%)$	1.43 ± 0.04	$10\pm2/608~(~1.6\pm~0.3\%)$	
All	All	0.95	0/18816 (0.0%)	1.43	407/25536 (1.6%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0 ± 0.0	0.7 ± 0.7
All	All	0	28

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Chain Res Type Atoms		$\mathbf{Z} = \mathbf{Observed}(^{o})$		Ideal(0)	Models		
IVIOI	Chain	nes	Type	Atoms	L	$\mathrm{Observed}(^{o})$	$\mathrm{Ideal}(^{o})$	Worst	Total
1	A	11	ASN	N-CA-C	-7.94	89.57	111.00	10	31
1	A	19	GLY	N-CA-C	-7.80	93.61	113.10	25	27
1	A	60	VAL	N-CA-C	-7.65	90.33	111.00	18	11
1	A	42	GLY	N-CA-C	-7.30	94.86	113.10	21	5
1	A	44	GLY	N-CA-C	-7.23	95.02	113.10	31	13
1	A	14	TYR	N-CA-C	-7.01	92.06	111.00	31	4
1	A	39	PHE	CB-CG-CD1	6.90	125.63	120.80	4	3
1	A	38	GLY	N-CA-C	-6.90	95.85	113.10	16	26
1	A	17	PHE	N-CA-C	-6.90	92.38	111.00	17	15
1	A	20	SER	N-CA-CB	6.80	120.70	110.50	1	12
1	A	39	PHE	CB-CG-CD2	-6.70	116.11	120.80	4	4
1	A	8	GLN	N-CA-C	-6.69	92.94	111.00	37	29

Continued on next page...



 $Continued\ from\ previous\ page...$

	Clasies				\mathbf{Z}	Ob 22222 d(0)	T-11(0)	Mod	dels
Mol	Chain	Res	Type	Atoms	L	$Observed(^o)$	$\operatorname{Ideal}(^{o})$	Worst	Total
1	A	15	HIS	N-CA-C	-6.64	93.08	111.00	14	25
1	A	3	ASP	N-CA-C	-6.54	93.33	111.00	30	35
1	A	51	ALA	N-CA-C	-6.44	93.61	111.00	37	4
1	A	14	TYR	CB-CG-CD1	6.41	124.85	121.00	30	1
1	A	16	CYS	N-CA-C	-6.36	93.82	111.00	31	24
1	A	14	TYR	CB-CG-CD2	-6.31	117.21	121.00	30	1
1	A	40	LYS	N-CA-C	-6.24	94.16	111.00	24	8
1	A	57	GLY	N-CA-C	-6.23	97.53	113.10	36	13
1	A	43	HIS	N-CA-C	-6.12	94.47	111.00	9	3
1	A	35	GLY	N-CA-C	-6.06	97.94	113.10	2	29
1	A	45	LEU	CB-CA-C	5.98	121.56	110.20	4	3
1	A	56	VAL	N-CA-C	-5.88	95.13	111.00	1	1
1	A	52	LEU	N-CA-C	-5.81	95.32	111.00	2	3
1	A	39	PHE	N-CA-C	-5.80	95.33	111.00	3	15
1	A	36	HIS	N-CA-C	-5.80	95.34	111.00	42	12
1	A	17	PHE	CB-CG-CD1	5.76	124.83	120.80	29	2
1	A	45	LEU	N-CA-C	-5.75	95.48	111.00	2	5
1	A	12	CYS	N-CA-C	-5.74	95.52	111.00	39	5
1	A	13	VAL	N-CA-C	-5.71	95.59	111.00	1	16
1	A	41	VAL	N-CA-C	-5.68	95.67	111.00	35	3
1	A	37	CYS	N-CA-C	-5.61	95.85	111.00	42	6
1	A	2	ARG	N-CA-C	-5.55	96.03	111.00	8	2
1	A	12	CYS	N-CA-CB	5.51	120.51	110.60	36	2
1	A	17	PHE	CB-CG-CD2	-5.47	116.97	120.80	29	2
1	A	10	GLU	N-CA-C	-5.43	96.33	111.00	23	1
1	A	4	GLY	N-CA-C	-5.38	99.65	113.10	36	1
1	A	46	ALA	N-CA-C	-5.35	96.55	111.00	41	2
1	A	7	ALA	N-CA-C	-5.20	96.95	111.00	18	1
1	A	34	SER	N-CA-C	-5.07	97.32	111.00	32	2

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	5	TYR	Sidechain	13
1	A	14	TYR	Sidechain	5
1	A	2	ARG	Sidechain	3
1	A	51	ALA	Peptide	2
1	A	57	GLY	Peptide	1
1	A	17	PHE	Peptide	1

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	19	GLY	Peptide	1
1	A	58	ILE	Peptide	1
1	A	20	SER	Peptide	1

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	436	408	405	1±1
All	All	18312	17136	17010	33

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2		Distance(A)	Worst	Total
1:A:7:ALA:HB2	1:A:59:ILE:HD13	0.67	1.65	5	14
1:A:41:VAL:HG12	1:A:42:GLY:H	0.57	1.59	24	2
1:A:13:VAL:HG12	1:A:59:ILE:HD12	0.52	1.80	2	12
1:A:7:ALA:HB2	1:A:59:ILE:HD12	0.49	1.84	26	1
1:A:36:HIS:CE1	1:A:48:TRP:CD1	0.49	3.01	19	3
1:A:13:VAL:HG12	1:A:59:ILE:HD11	0.45	1.88	26	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	59/68 (87%)	49±2 (82±4%)	8±2 (14±4%)	2±1 (4±2%)	5	32
All	All	2478/2856 (87%)	2042 (82%)	339 (14%)	97 (4%)	5	32



All 16 unique Ramachandran outliers are listed below.	They are sorted by the frequency of
occurrence in the ensemble.	

Mol	Chain	Res	Type	Models (Total)
1	A	58	ILE	27
1	A	12	CYS	21
1	A	41	VAL	13
1	A	20	SER	9
1	A	18	PRO	5
1	A	40	LYS	4
1	A	31	GLY	3
1	A	43	HIS	3
1	A	51	ALA	3
1	A	44	GLY	2
1	A	38	GLY	2
1	A	11	ASN	1
1	A	56	VAL	1
1	A	19	GLY	1
1	A	36	HIS	1
1	A	42	GLY	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Per	centiles
1	A	47/53 (89%)	42±2 (89±4%)	5±2 (11±4%)	10	54
All	All	1974/2226 (89%)	1756 (89%)	218 (11%)	10	54

All 30 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	14	TYR	42
1	A	2	ARG	28
1	A	13	VAL	16
1	A	12	CYS	15
1	A	25	THR	13
1	A	45	LEU	11
1	A	41	VAL	11
1	A	60	VAL	10

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	21	SER	9
1	A	50	ASN	7
1	A	59	ILE	7
1	A	47	CYS	5
1	A	34	SER	5
1	A	11	ASN	4
1	A	23	CYS	4
1	A	40	LYS	4
1	A	30	LYS	4
1	A	33	THR	3
1	A	16	CYS	3
1	A	20	SER	3
1	A	39	PHE	2
1	A	52	LEU	2
1	A	18	PRO	2
1	A	28	LYS	2
1	A	56	VAL	1
1	A	27	CYS	1
1	A	8	GLN	1
1	A	43	HIS	1
1	A	17	PHE	1
1	A	26	LEU	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.



6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

No chemical shift data were provided

