



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 7, 2024 – 05:26 am GMT

PDB ID : 6FKW
Title : Europium-containing methanol dehydrogenase
Authors : Barends, T.; Dietl, A.
Deposited on : 2018-01-24
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	577	97%
1	B	577	95%
1	C	577	96%
1	D	577	20% (poor fit), 95%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 19596 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Methanol dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	576	4494	2879	761	836	18	0	1	0
1	B	576	4494	2881	759	835	19	0	2	0
1	C	576	4490	2876	760	836	18	0	1	0
1	D	576	4490	2876	759	837	18	0	1	0

- Molecule 2 is EUROPIUM ION (three-letter code: EU) (formula: Eu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Eu	0	0
			1	1		
2	B	1	Total	Eu	0	0
			1	1		
2	C	1	Total	Eu	0	0
			1	1		
2	D	1	Total	Eu	0	0
			1	1		

- Molecule 3 is PYRROLOQUINOLINE QUINONE (three-letter code: PQQ) (formula: C₁₄H₆N₂O₈).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	24	14	2	8	0	0
3	B	1	24	14	2	8	0	0
3	C	1	24	14	2	8	0	0
3	D	1	24	14	2	8	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	509	509	509	0	0
4	B	427	427	427	0	0
4	C	443	443	443	0	0
4	D	149	149	149	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Methanol dehydrogenase

Chain A:  97%



- Molecule 1: Methanol dehydrogenase

Chain B:  95%




- Molecule 1: Methanol dehydrogenase


Chain C:  96%



- Molecule 1: Methanol dehydrogenase

Chain D:  20% 95%





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	55.08Å 101.18Å 107.79Å 115.21° 97.53° 91.78°	Depositor
Resolution (Å)	96.27 – 1.40 48.80 – 1.40	Depositor EDS
% Data completeness (in resolution range)	90.6 (96.27-1.40) 90.6 (48.80-1.40)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 1.40Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.169 , 0.185 0.173 , 0.189	Depositor DCC
R_{free} test set	18419 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	21.7	Xtrriage
Anisotropy	0.070	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	19596	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PQQ, EU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/4625	0.58	0/6295
1	B	0.29	0/4631	0.56	0/6304
1	C	0.29	0/4624	0.56	0/6295
1	D	0.28	0/4624	0.49	0/6295
All	All	0.29	0/18504	0.55	0/25189

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4494	0	4337	12	0
1	B	4494	0	4343	13	0
1	C	4490	0	4331	13	0
1	D	4490	0	4329	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	24	0	3	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	24	0	3	1	0
3	C	24	0	3	2	0
3	D	24	0	3	2	0
4	A	509	0	0	2	0
4	B	427	0	0	1	0
4	C	443	0	0	0	0
4	D	149	0	0	1	0
All	All	19596	0	17352	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:570:GLU:OE2	1:D:572:TYR:OH	2.12	0.67
1:B:42:LEU:HD22	1:D:42:LEU:HD22	1.81	0.62
1:D:370:THR:HG21	1:D:412:TYR:CD1	2.37	0.59
1:A:42:LEU:HD22	1:C:42:LEU:HD22	1.85	0.57
1:C:426:VAL:HG11	1:C:550:LEU:HD21	1.87	0.56
1:D:384:MET:HG2	1:D:532:TRP:CE3	2.42	0.54
1:A:238:TRP:CZ2	3:A:602:PQQ:C6A	2.92	0.53
1:A:370:THR:HG21	1:A:412:TYR:CD1	2.45	0.52
1:D:44:TRP:CH2	1:D:573:VAL:HG21	2.46	0.50
1:C:370:THR:HG21	1:C:412:TYR:CD1	2.47	0.49
1:A:570:GLU:OE2	1:C:572:TYR:OH	2.30	0.49
1:A:532:TRP:CE3	3:A:602:PQQ:O4	2.66	0.49
1:D:238:TRP:CZ2	1:D:303:VAL:HG21	2.48	0.49
1:D:238:TRP:CE2	1:D:303:VAL:HG21	2.48	0.48
1:B:370:THR:HG21	1:B:412:TYR:CD1	2.48	0.48
1:D:379:ILE:HD13	4:D:751:HOH:O	2.11	0.48
1:B:426:VAL:HG11	1:B:550:LEU:HD21	1.95	0.48
1:D:44:TRP:CZ2	1:D:573:VAL:HG21	2.49	0.48
1:B:238:TRP:CZ2	3:B:602:PQQ:C6A	2.97	0.47
1:C:520:LYS:HE3	1:C:575:ALA:HB1	1.95	0.47
1:C:238:TRP:CZ2	3:C:602:PQQ:C6A	2.97	0.47
1:D:532:TRP:CE3	3:D:602:PQQ:O4	2.67	0.47
1:B:238:TRP:CE2	1:B:303:VAL:HG21	2.51	0.46
1:A:42:LEU:CD2	1:C:42:LEU:HD22	2.46	0.45
1:C:238:TRP:CE2	1:C:303:VAL:HG21	2.52	0.45
1:B:42:LEU:HD22	1:D:42:LEU:CD2	2.46	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:544:ASP:N	1:A:544:ASP:OD1	2.49	0.45
1:D:479:PHE:CE2	1:D:489:ALA:HB2	2.52	0.44
1:A:572:TYR:OH	1:C:570:GLU:OE2	2.34	0.44
1:A:42:LEU:HD22	1:C:42:LEU:CD2	2.48	0.43
1:B:384:MET:HG2	1:B:532:TRP:CE3	2.54	0.43
3:D:602:PQQ:O9A	3:D:602:PQQ:N1	2.48	0.43
1:D:115:ALA:CB	1:D:159:VAL:HG11	2.49	0.43
1:B:7:LEU:HB3	1:B:15:VAL:HG11	2.01	0.42
1:B:44:TRP:CH2	1:B:573:VAL:HG21	2.55	0.42
1:B:502:LYS:NZ	4:B:702:HOH:O	2.52	0.42
1:C:44:TRP:CH2	1:C:573:VAL:HG21	2.55	0.42
1:A:493:LYS:NZ	4:A:703:HOH:O	2.53	0.41
1:B:469:GLY:O	1:B:511:PRO:HD2	2.21	0.41
1:B:479:PHE:CE2	1:B:489:ALA:HB2	2.56	0.41
1:D:349:VAL:HG13	1:D:351:TRP:CD1	2.56	0.41
1:C:490:VAL:HG12	1:C:497:VAL:HG22	2.02	0.41
1:C:532:TRP:CE3	3:C:602:PQQ:O4	2.74	0.41
1:A:238:TRP:CE2	1:A:303:VAL:HG21	2.56	0.40
1:A:502:LYS:NZ	4:A:701:HOH:O	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	575/577 (100%)	553 (96%)	20 (4%)	2 (0%)	41	18
1	B	576/577 (100%)	558 (97%)	16 (3%)	2 (0%)	41	18
1	C	575/577 (100%)	554 (96%)	19 (3%)	2 (0%)	41	18
1	D	575/577 (100%)	553 (96%)	21 (4%)	1 (0%)	47	21
All	All	2301/2308 (100%)	2218 (96%)	76 (3%)	7 (0%)	41	18

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	569	GLY
1	A	107	VAL
1	A	569	GLY
1	B	107	VAL
1	B	569	GLY
1	C	107	VAL
1	D	569	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	462/462 (100%)	456 (99%)	6 (1%)	69	42
1	B	463/462 (100%)	457 (99%)	6 (1%)	69	42
1	C	462/462 (100%)	456 (99%)	6 (1%)	69	42
1	D	462/462 (100%)	455 (98%)	7 (2%)	65	37
All	All	1849/1848 (100%)	1824 (99%)	25 (1%)	69	40

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	196	GLN
1	A	314[A]	LYS
1	A	314[B]	LYS
1	A	371	LYS
1	A	412	TYR
1	A	544	ASP
1	B	39	ARG
1	B	83	LYS
1	B	196	GLN
1	B	371	LYS
1	B	412	TYR
1	B	544	ASP
1	C	98	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	196	GLN
1	C	358	GLU
1	C	371	LYS
1	C	412	TYR
1	C	544	ASP
1	D	35	LYS
1	D	172	GLU
1	D	196	GLN
1	D	316	LYS
1	D	371	LYS
1	D	416	GLU
1	D	544	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	558	GLN
1	C	82	GLN
1	C	229	GLN
1	D	82	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PQQ	A	602	2	23,26,26	1.89	2 (8%)	29,40,40	1.84	6 (20%)
3	PQQ	D	602	2	23,26,26	1.94	2 (8%)	29,40,40	1.84	7 (24%)
3	PQQ	B	602	2	23,26,26	1.87	2 (8%)	29,40,40	1.92	6 (20%)
3	PQQ	C	602	2	23,26,26	1.87	3 (13%)	29,40,40	1.95	7 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PQQ	A	602	2	-	0/10/28/28	0/3/3/3
3	PQQ	D	602	2	-	0/10/28/28	0/3/3/3
3	PQQ	B	602	2	-	0/10/28/28	0/3/3/3
3	PQQ	C	602	2	-	4/10/28/28	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	602	PQQ	C6A-C9A	6.20	1.50	1.42
3	A	602	PQQ	C6A-C9A	5.90	1.49	1.42
3	B	602	PQQ	C6A-C9A	5.77	1.49	1.42
3	C	602	PQQ	C6A-C9A	5.68	1.49	1.42
3	D	602	PQQ	C9-C9A	4.74	1.51	1.41
3	A	602	PQQ	C9-C9A	4.55	1.50	1.41
3	B	602	PQQ	C9-C9A	4.52	1.50	1.41
3	C	602	PQQ	C9-C9A	4.44	1.50	1.41
3	C	602	PQQ	O2B-C2X	-2.15	1.24	1.30

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	602	PQQ	C9-C9A-C1A	5.78	127.62	122.88
3	C	602	PQQ	C9-C9A-C1A	5.60	127.47	122.88
3	A	602	PQQ	C9-C9A-C1A	5.59	127.46	122.88

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	602	PQQ	C9-C9A-C1A	5.39	127.30	122.88
3	B	602	PQQ	C9-C9A-C6A	-4.50	115.66	121.68
3	C	602	PQQ	C9-C9A-C6A	-4.30	115.94	121.68
3	A	602	PQQ	C9-C9A-C6A	-4.10	116.20	121.68
3	D	602	PQQ	C9-C9A-C6A	-4.09	116.21	121.68
3	B	602	PQQ	C2-N1-C1A	3.34	110.75	103.90
3	A	602	PQQ	C2-N1-C1A	3.18	110.44	103.90
3	D	602	PQQ	C2-N1-C1A	3.14	110.35	103.90
3	C	602	PQQ	C2-N1-C1A	3.12	110.30	103.90
3	D	602	PQQ	C6A-N6-C7	2.66	122.37	117.91
3	C	602	PQQ	O9B-C9X-C9	2.64	121.88	114.39
3	D	602	PQQ	O2B-C2X-C2	2.46	120.28	114.69
3	C	602	PQQ	C6A-N6-C7	2.41	121.95	117.91
3	A	602	PQQ	C8-C9-C9A	-2.40	116.80	120.06
3	D	602	PQQ	O2A-C2X-C2	-2.36	116.45	121.24
3	B	602	PQQ	C6A-N6-C7	2.30	121.78	117.91
3	A	602	PQQ	C6A-N6-C7	2.26	121.71	117.91
3	A	602	PQQ	O7A-C7X-C7	-2.16	116.86	121.24
3	C	602	PQQ	O2B-C2X-C2	2.12	119.51	114.69
3	C	602	PQQ	C8-C9-C9A	-2.11	117.18	120.06
3	B	602	PQQ	C8-C9-C9A	-2.10	117.20	120.06
3	B	602	PQQ	O7A-C7X-C7	-2.09	117.00	121.24
3	D	602	PQQ	C8-C9-C9A	-2.08	117.23	120.06

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	602	PQQ	C9A-C9-C9X-O9B
3	C	602	PQQ	C9A-C9-C9X-O9A
3	C	602	PQQ	C8-C9-C9X-O9A
3	C	602	PQQ	C8-C9-C9X-O9B


There are no ring outliers.


4 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	PQQ	2	0
3	D	602	PQQ	2	0
3	B	602	PQQ	1	0
3	C	602	PQQ	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	576/577 (99%)	-0.04	2 (0%) 94 93	14, 20, 29, 42	0
1	B	576/577 (99%)	-0.11	2 (0%) 94 93	17, 23, 34, 45	0
1	C	576/577 (99%)	-0.16	3 (0%) 91 89	16, 22, 34, 46	0
1	D	576/577 (99%)	1.23	117 (20%) 1 0	23, 40, 63, 77	0
All	All	2304/2308 (99%)	0.23	124 (5%) 25 24	14, 24, 52, 77	0

All (124) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	230	TRP	7.5
1	D	202	VAL	6.9
1	D	418	THR	6.6
1	D	225	TRP	6.6
1	D	221	GLY	6.5
1	D	233	GLY	6.4
1	D	223	LYS	6.2
1	D	203	LEU	5.8
1	D	272	TRP	5.6
1	D	198	PRO	5.3
1	D	420	THR	5.3
1	D	267	GLY	5.3
1	D	206	SER	5.2
1	D	228	GLU	5.2
1	D	544	ASP	4.9
1	D	184	ILE	4.9
1	D	419	TYR	4.9
1	D	549	GLY	4.9
1	D	204	LEU	4.9
1	D	546	PRO	4.9
1	D	212	PHE	4.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	219	GLY	4.8
1	D	208	PHE	4.8
1	D	547	TYR	4.7
1	D	222	THR	4.7
1	D	286	ALA	4.6
1	D	190	VAL	4.6
1	D	197	GLY	4.5
1	D	207	ASP	4.5
1	D	266	LYS	4.5
1	D	232	LEU	4.5
1	D	295	TRP	4.4
1	D	226	PRO	4.4
1	D	283	THR	4.4
1	D	220	ASP	4.3
1	D	425	TYR	4.3
1	D	148	THR	4.2
1	D	98	GLN	4.2
1	D	257	PRO	4.2
1	D	188	ASN	4.1
1	D	213	PRO	4.1
1	D	193	ALA	4.0
1	D	199	ASP	4.0
1	D	218	PRO	3.9
1	D	173	PHE	3.9
1	D	224	THR	3.9
1	D	409	CYS	3.9
1	D	231	LYS	3.9
1	D	284	GLY	3.8
1	D	191	TRP	3.8
1	D	379	ILE	3.7
1	D	247	LEU	3.7
1	D	248	ASP	3.7
1	D	421	ALA	3.6
1	D	280	ARG	3.6
1	D	205	ASP	3.5
1	D	254	THR	3.4
1	D	181	ALA	3.4
1	D	200	GLU	3.3
1	D	192	LEU	3.2
1	D	376	THR	3.2
1	D	217	GLY	3.1
1	D	416	GLU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	83	LYS	3.1
1	D	96	ASN	3.0
1	D	162[A]	ASP	3.0
1	D	262	ALA	3.0
1	D	149	LYS	3.0
1	D	194	TYR	3.0
1	D	423	ALA	2.9
1	D	358	GLU	2.9
1	D	189	ARG	2.9
1	D	263	GLU	2.9
1	D	227	GLY	2.8
1	D	102	VAL	2.8
1	D	282	ASP	2.8
1	D	160	VAL	2.8
1	C	358	GLU	2.8
1	D	436	HIS	2.7
1	D	281	PRO	2.7
1	D	359	ASN	2.7
1	C	2	ASP	2.7
1	D	161	LYS	2.7
1	D	277	PHE	2.7
1	D	415	VAL	2.6
1	D	417	ALA	2.6
1	D	340	LEU	2.6
1	D	373	GLY	2.6
1	D	278	ALA	2.5
1	D	246	ALA	2.5
1	D	412	TYR	2.5
1	D	378	GLY	2.5
1	D	493	LYS	2.5
1	D	134	ASN	2.4
1	D	171	GLY	2.4
1	D	287	ARG	2.4
1	D	243	TYR	2.4
1	D	269	ASP	2.4
1	D	545	ASP	2.4
1	D	174	GLY	2.4
1	D	285	LYS	2.3
1	D	137	LYS	2.3
1	B	358	GLU	2.3
1	D	427	GLY	2.3
1	D	141	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	3	GLU	2.3
1	D	431	LEU	2.3
1	D	216	GLY	2.2
1	D	229	GLN	2.2
1	D	430	VAL	2.2
1	B	550	LEU	2.2
1	D	201	GLU	2.2
1	A	210	LYS	2.2
1	A	367	GLU	2.2
1	C	314	LYS	2.2
1	D	140	TRP	2.2
1	D	2	ASP	2.1
1	D	550	LEU	2.1
1	D	210	LYS	2.1
1	D	101	ALA	2.1
1	D	255	SER	2.1
1	D	103	ALA	2.1
1	D	428	ALA	2.1
1	D	294	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PQQ	D	602	24/24	0.85	0.11	32,41,45,47	0
3	PQQ	A	602	24/24	0.91	0.10	18,21,23,25	0
3	PQQ	C	602	24/24	0.92	0.09	19,21,22,23	0
3	PQQ	B	602	24/24	0.93	0.09	20,21,23,23	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	EU	A	601	1/1	0.97	0.05	24,24,24,24	1
2	EU	D	601	1/1	0.98	0.04	42,42,42,42	1
2	EU	C	601	1/1	0.99	0.07	20,20,20,20	1
2	EU	B	601	1/1	0.99	0.07	22,22,22,22	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around PQQ A 602:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around PQQ C 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.