



Full wwPDB X-ray Structure Validation Report i

Feb 18, 2024 – 05:35 PM EST

PDB ID : 4FMS
Title : Crystal structure of *Pseudomonas aeruginosa* OccK2 (OpdF) in complex with glucuronate
Authors : van den Berg, B.; Eren, E.
Deposited on : 2012-06-18
Resolution : 2.45 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	A	397	2%	84%	8%	•	7%
1	B	397	1%	80%	12%	•	7%

2 Entry composition [\(i\)](#)


There are 6 unique types of molecules in this entry. The entry contains 6237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable porin.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	B	368	Total	C 2875	N 1805	O 526	S 539	0	2	0
1	A	369	Total	C 2876	N 1810	O 521	S 540	0	2	0

- Molecule 2 is beta-D-glucopyranuronic acid (three-letter code: BDP) (formula: C₆H₁₀O₇).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	B	1	Total	C 13	O 6	O 7	0	0
2	A	1	Total	C 13	O 6	O 7	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0

- Molecule 4 is (HYDROXYETHYLOXY)TRI(ETHYLOXY)OCTANE (three-letter code: C8E) (formula: C₁₆H₃₄O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O		
			21	16	5	0	0
4	B	1	Total	C	O		
			17	14	3	0	0
4	B	1	Total	C	O		
			12	10	2	0	0
4	B	1	Total	C	O		
			17	12	5	0	0
4	A	1	Total	C	O		
			10	9	1	0	0
4	A	1	Total	C	O		
			21	16	5	0	0
4	A	1	Total	C	O		
			13	8	5	0	0

- Molecule 5 is (1R)-2-{{[(2S)-2,3-DIHYDROXYPROPYL]OXY}(HYDROXY)PHOSPHORYL]OXY}-1-[(PALMITOYLOXY)METHYL]ETHYL (11E)-OCTADEC-11-ENOATE (three-letter code: PGV) (formula: C₄₀H₇₇O₁₀P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	O	P		
			26	17	8	1	0	0
5	A	1	Total	C	O	P		
			26	16	9	1	0	0
5	A	1	Total	C	O	P		
			22	12	9	1	0	0
5	A	1	Total	C	O	P		
			24	15	8	1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	113	Total O 113 113	0	0
6	A	118	Total O 118 118	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable porin



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.44Å 206.80Å 51.75Å 90.00° 99.09° 90.00°	Depositor
Resolution (Å)	14.99 – 2.45 38.18 – 2.39	Depositor EDS
% Data completeness (in resolution range)	92.1 (14.99-2.45) 90.7 (38.18-2.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	3.09 (at 2.39Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R , R_{free}	0.191 , 0.242 0.186 , 0.234	Depositor DCC
R_{free} test set	2000 reflections (5.82%)	wwPDB-VP
Wilson B-factor (Å ²)	32.8	Xtriage
Anisotropy	0.458	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6237	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PGV, BDP, C8E, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/2946	0.59	1/3976 (0.0%)
1	B	0.41	0/2944	0.58	1/3972 (0.0%)
All	All	0.41	0/5890	0.58	2/7948 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	69	LEU	CA-CB-CG	5.87	128.80	115.30
1	B	69	LEU	CA-CB-CG	5.06	126.93	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2876	0	2734	19	0
1	B	2875	0	2743	29	0
2	A	13	0	9	0	0
2	B	13	0	9	0	0
3	A	10	0	0	0	0
3	B	10	0	0	1	0
4	A	44	0	68	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	67	0	105	5	0
5	A	72	0	57	2	0
5	B	26	0	25	0	0
6	A	118	0	0	3	0
6	B	113	0	0	11	0
All	All	6237	0	5750	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:291:ASN:OD1	6:B:593:HOH:O	1.91	0.88
1:B:248:ASP:OD1	1:B:250:ARG:NH2	2.09	0.86
1:B:307:GLN:OE1	6:B:593:HOH:O	1.96	0.82
1:A:95:PHE:HB2	1:A:164:ALA:HB2	1.66	0.78
1:B:25:HIS:HB2	4:B:405:C8E:H162	1.71	0.73
1:A:174:ARG:NH2	6:A:524:HOH:O	2.25	0.68
1:A:248:ASP:OD1	1:A:250:ARG:NH2	2.27	0.67
1:B:95:PHE:HB2	1:B:164:ALA:HB2	1.75	0.67
1:B:324:PHE:HZ	4:B:404:C8E:H141	1.60	0.67
1:B:72:ASP:O	6:B:570:HOH:O	2.13	0.66
1:B:64:LEU:HB3	1:B:99:ALA:HB3	1.85	0.59
1:A:263:HIS:HE2	4:A:404:C8E:H71	1.68	0.59
1:B:178:THR:O	1:B:243:ARG:NH1	2.36	0.58
1:A:269:LEU:HB3	4:A:406:C8E:H111	1.85	0.58
5:A:408:PGV:H042	6:A:614:HOH:O	2.04	0.56
1:B:161:ARG:NH2	3:B:402:SO4:O1	2.32	0.55
1:B:345:GLU:OE2	1:B:375:ARG:HD2	2.06	0.55
1:B:374:ARG:NH2	6:B:564:HOH:O	2.43	0.51
1:A:256:PHE:HB3	4:A:405:C8E:H42	1.92	0.51
1:B:362:LYS:O	1:B:363:ARG:HB2	2.11	0.50
1:B:188:GLU:OE1	1:B:190:ARG:NH2	2.45	0.48
1:A:66:LEU:HB3	1:A:131:LEU:HD13	1.95	0.48
1:B:162:ASN:HB3	6:B:595:HOH:O	2.13	0.48
1:A:248:ASP:HB3	1:A:275:ASP:HB2	1.95	0.48
1:A:374:ARG:O	1:A:375:ARG:HD3	2.14	0.48
1:B:259:ARG:NH1	6:B:600:HOH:O	2.29	0.47
1:A:250:ARG:HD3	5:A:409:PGV:H032	1.97	0.46
4:A:405:C8E:H72	4:A:405:C8E:H41	1.65	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:308:LEU:HD12	1:A:308:LEU:HA	1.80	0.46
1:B:314:PHE:HB2	1:B:322:LEU:HB3	1.98	0.46
1:B:158:ASN:HB2	1:B:169:MET:HG2	1.98	0.45
1:B:266:TYR:HB2	1:B:309:ARG:HB3	1.98	0.45
1:A:258:ALA:HB2	4:A:405:C8E:H22	1.97	0.45
1:B:25:HIS:CB	4:B:405:C8E:H162	2.44	0.44
1:B:386:ASN:ND2	4:B:405:C8E:O15	2.46	0.44
1:A:263:HIS:NE2	4:A:404:C8E:H71	2.30	0.44
1:A:382:ARG:HE	1:A:382:ARG:HB3	1.63	0.44
1:B:268:GLY:HA3	6:B:593:HOH:O	2.16	0.44
1:B:136:ARG:NH2	6:B:567:HOH:O	2.50	0.43
4:B:404:C8E:H41	4:A:404:C8E:H51	2.01	0.43
1:B:259:ARG:NE	6:B:569:HOH:O	2.52	0.42
1:B:348:ARG:HG3	1:A:317:LEU:HD13	2.02	0.42
1:B:374:ARG:O	1:B:375:ARG:HD3	2.20	0.42
1:B:7:GLN:HA	1:B:12:ASP:OD2	2.19	0.41
1:B:25:HIS:HB3	6:B:609:HOH:O	2.20	0.41
1:B:163:ASP:O	6:B:502:HOH:O	2.21	0.41
1:A:123:LEU:HD13	1:A:182:PHE:CE2	2.55	0.41
1:A:136:ARG:HE	1:A:136:ARG:HB3	1.57	0.41
1:A:149:THR:HG22	6:A:594:HOH:O	2.21	0.41
1:A:263:HIS:HE1	4:A:404:C8E:H21	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	363/397 (91%)	353 (97%)	10 (3%)	0	100 100
1	B	364/397 (92%)	352 (97%)	12 (3%)	0	100 100
All	All	727/794 (92%)	705 (97%)	22 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	288/314 (92%)	270 (94%)	18 (6%)	18 22
1	B	289/314 (92%)	271 (94%)	18 (6%)	18 23
All	All	577/628 (92%)	541 (94%)	36 (6%)	18 23

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	10	LEU
1	B	69	LEU
1	B	98	LEU
1	B	120	LEU
1	B	140	LEU
1	B	141	SER
1	B	171	LEU
1	B	174	ARG
1	B	200	LEU
1	B	231	LEU
1	B	246	LYS
1	B	248	ASP
1	B	250	ARG
1	B	336	LEU
1	B	375	ARG
1	B	387	ARG
1	B	394	LEU
1	B	396	LEU
1	A	29	VAL
1	A	69	LEU
1	A	70	LYS
1	A	98	LEU
1	A	120	LEU
1	A	171	LEU
1	A	200	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	237	ARG
1	A	250	ARG
1	A	253	SER
1	A	285	SER
1	A	308	LEU
1	A	372	SER
1	A	375	ARG
1	A	382	ARG
1	A	387	ARG
1	A	394	LEU
1	A	396	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	C8E	A	405	-	20,20,20	0.43	0	19,19,19	0.29	0
2	BDP	A	401	-	13,13,13	0.74	0	18,19,19	0.59	0
3	SO4	B	403	-	4,4,4	0.11	0	6,6,6	0.28	0
5	PGV	A	407	-	25,25,50	1.02	3 (12%)	28,30,56	1.51	6 (21%)
4	C8E	B	406	-	11,11,20	0.37	0	10,10,19	0.35	0
4	C8E	B	407	-	16,16,20	0.48	0	15,15,19	0.26	0
5	PGV	A	409	-	23,23,50	1.09	4 (17%)	27,28,56	1.41	5 (18%)
4	C8E	A	404	-	9,9,20	0.33	0	8,8,19	0.29	0
5	PGV	A	408	-	21,21,50	1.12	2 (9%)	24,26,56	1.62	6 (25%)
3	SO4	A	402	-	4,4,4	0.17	0	6,6,6	0.33	0
4	C8E	B	404	-	20,20,20	0.39	0	19,19,19	0.50	0
4	C8E	B	405	-	16,16,20	0.46	0	15,15,19	0.31	0
5	PGV	B	408	-	25,25,50	1.01	4 (16%)	29,30,56	1.73	5 (17%)
3	SO4	A	403	-	4,4,4	0.16	0	6,6,6	0.09	0
2	BDP	B	401	-	13,13,13	0.75	0	18,19,19	0.60	0
3	SO4	B	402	-	4,4,4	0.12	0	6,6,6	0.35	0
4	C8E	A	406	-	12,12,20	0.47	0	11,11,19	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	C8E	A	405	-	-	11/18/18/18	-
2	BDP	A	401	-	-	1/4/24/24	0/1/1/1
5	PGV	A	407	-	-	12/29/29/55	-
5	PGV	A	409	-	-	6/25/25/55	-
4	C8E	B	407	-	-	6/14/14/18	-
4	C8E	B	405	-	-	7/14/14/18	-
5	PGV	A	408	-	-	7/25/25/55	-
4	C8E	B	404	-	-	10/18/18/18	-
4	C8E	B	406	-	-	3/9/9/18	-
5	PGV	B	408	-	-	10/27/27/55	-
4	C8E	A	404	-	-	4/7/7/18	-
2	BDP	B	401	-	-	0/4/24/24	0/1/1/1
4	C8E	A	406	-	-	4/10/10/18	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	408	PGV	O03-C01	-2.81	1.38	1.45
5	B	408	PGV	O03-C19	2.63	1.41	1.33
5	A	408	PGV	O01-C1	2.62	1.41	1.34
5	A	407	PGV	O01-C1	2.61	1.41	1.34
5	A	409	PGV	O03-C01	-2.59	1.39	1.45
5	A	407	PGV	O03-C01	-2.47	1.39	1.45
5	A	409	PGV	O01-C1	2.39	1.41	1.34
5	B	408	PGV	O03-C01	-2.24	1.40	1.45
5	B	408	PGV	O01-C02	-2.24	1.41	1.46
5	A	407	PGV	O03-C19	2.22	1.39	1.33
5	A	409	PGV	O01-C02	-2.10	1.41	1.46
5	B	408	PGV	O01-C1	2.09	1.40	1.34
5	A	409	PGV	O03-C19	2.05	1.39	1.33

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	408	PGV	O03-C01-C02	4.77	122.32	108.43
5	A	408	PGV	O03-C01-C02	3.61	118.95	108.43
5	B	408	PGV	O01-C1-C2	3.56	119.16	111.50
5	A	408	PGV	O01-C1-C2	3.51	120.60	110.80
5	B	408	PGV	O01-C1-O02	-3.43	115.41	123.70
5	A	407	PGV	O03-C01-C02	3.17	117.67	108.43
5	B	408	PGV	O03-C19-C20	3.13	121.73	111.91
5	A	407	PGV	O01-C1-C2	3.11	119.48	110.80
5	A	407	PGV	O03-C19-C20	3.07	121.54	111.91
5	A	409	PGV	O01-C1-C2	2.85	117.65	111.50
5	A	409	PGV	O03-C01-C02	2.83	116.68	108.43
5	A	408	PGV	O03-C19-C20	2.78	118.68	111.38
5	A	408	PGV	O12-C04-C05	2.39	116.98	108.99
5	A	408	PGV	C03-C02-C01	-2.30	106.35	111.79
5	A	407	PGV	C02-O01-C1	2.29	123.44	117.79
5	A	409	PGV	O01-C1-O02	-2.28	118.20	123.70
5	A	409	PGV	C3-C2-C1	2.26	121.83	113.62
5	B	408	PGV	O04-C19-C20	-2.24	115.00	123.73
5	A	408	PGV	C02-O01-C1	2.12	123.02	117.79
5	A	409	PGV	O03-C19-C20	2.03	118.28	111.91
5	A	407	PGV	C22-C21-C20	2.02	120.45	113.19
5	A	407	PGV	O03-C19-O04	-2.01	118.53	123.59

There are no chirality outliers.

All (81) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	408	PGV	C03-O11-P-O13
5	A	408	PGV	C04-C05-C06-O06
5	A	408	PGV	C2-C1-O01-C02
5	A	407	PGV	O04-C19-O03-C01
5	A	408	PGV	O02-C1-O01-C02
5	A	407	PGV	C20-C19-O03-C01
4	B	406	C8E	C4-C5-C6-C7
4	A	405	C8E	O15-C16-C17-O18
4	B	404	C8E	O12-C13-C14-O15
5	A	408	PGV	O01-C02-C03-O11
4	B	406	C8E	C6-C7-C8-O9
4	A	406	C8E	C16-C17-O18-C19
4	A	406	C8E	O12-C13-C14-O15
4	B	404	C8E	O18-C19-C20-O21
4	A	406	C8E	O18-C19-C20-O21
4	A	404	C8E	C7-C8-O9-C10
4	A	404	C8E	C4-C5-C6-C7
5	A	407	PGV	C20-C21-C22-C23
4	A	405	C8E	C4-C5-C6-C7
5	A	407	PGV	C19-C20-C21-C22
4	B	404	C8E	C3-C4-C5-C6
4	B	405	C8E	C4-C5-C6-C7
5	B	408	PGV	C20-C21-C22-C23
5	B	408	PGV	C21-C22-C23-C24
4	A	405	C8E	O18-C19-C20-O21
5	A	407	PGV	O02-C1-O01-C02
4	A	404	C8E	C2-C3-C4-C5
4	B	405	C8E	O12-C13-C14-O15
5	A	408	PGV	C01-C02-C03-O11
4	B	404	C8E	C1-C2-C3-C4
4	A	404	C8E	C1-C2-C3-C4
4	A	405	C8E	C1-C2-C3-C4
4	B	404	C8E	C2-C3-C4-C5
5	B	408	PGV	O03-C01-C02-O01
5	B	408	PGV	O03-C01-C02-C03
4	B	405	C8E	C3-C4-C5-C6
5	A	407	PGV	O03-C01-C02-O01
4	A	405	C8E	C3-C4-C5-C6
5	A	407	PGV	C23-C24-C25-C26
4	B	406	C8E	C2-C3-C4-C5
4	A	405	C8E	C13-C14-O15-C16
4	B	407	C8E	C11-C10-O9-C8
4	A	405	C8E	C14-C13-O12-C11

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	A	407	PGV	O12-C04-C05-O05
5	B	408	PGV	C4-C5-C6-C7
4	A	405	C8E	C6-C7-C8-O9
5	A	409	PGV	C21-C22-C23-C24
5	A	407	PGV	C04-O12-P-O11
5	B	408	PGV	C3-C4-C5-C6
4	B	404	C8E	C4-C5-C6-C7
4	A	405	C8E	C17-C16-O15-C14
5	A	407	PGV	O03-C01-C02-C03
4	B	405	C8E	C1-C2-C3-C4
4	B	407	C8E	C7-C8-O9-C10
4	A	406	C8E	C20-C19-O18-C17
5	B	408	PGV	C19-C20-C21-C22
5	B	408	PGV	C5-C6-C7-C8
5	B	408	PGV	O01-C02-C03-O11
4	B	404	C8E	C16-C17-O18-C19
5	A	409	PGV	O04-C19-O03-C01
4	B	405	C8E	C5-C6-C7-C8
4	B	404	C8E	C13-C14-O15-C16
5	A	409	PGV	C01-C02-C03-O11
4	B	404	C8E	C17-C16-O15-C14
4	B	405	C8E	C10-C11-O12-C13
4	A	405	C8E	C16-C17-O18-C19
5	A	409	PGV	C02-C01-O03-C19
5	B	408	PGV	C03-O11-P-O12
4	B	405	C8E	C17-C16-O15-C14
5	A	409	PGV	O03-C19-C20-C21
4	B	407	C8E	C16-C17-O18-C19
5	A	409	PGV	C20-C21-C22-C23
4	B	407	C8E	C20-C19-O18-C17
5	A	407	PGV	O04-C19-C20-C21
4	B	407	C8E	C14-C13-O12-C11
5	A	408	PGV	C04-O12-P-O13
4	B	404	C8E	O15-C16-C17-O18
5	A	407	PGV	O03-C19-C20-C21
2	A	401	BDP	C4-C5-C6-O6B
4	B	407	C8E	O15-C16-C17-O18
4	A	405	C8E	O12-C13-C14-O15


There are no ring outliers.


8 monomers are involved in 15 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	405	C8E	3	0
5	A	409	PGV	1	0
4	A	404	C8E	4	0
5	A	408	PGV	1	0
4	B	404	C8E	2	0
4	B	405	C8E	3	0
3	B	402	SO4	1	0
4	A	406	C8E	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	369/397 (92%)	0.02	9 (2%) 59 54	21, 31, 48, 61	0
1	B	368/397 (92%)	0.04	3 (0%) 86 86	19, 30, 46, 54	0
All	All	737/794 (92%)	0.03	12 (1%) 72 69	19, 30, 47, 61	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	29	VAL	4.1
1	A	146	ALA	3.6
1	B	358	SER	3.2
1	A	222	LEU	3.0
1	A	360	ALA	3.0
1	A	358	SER	2.6
1	A	1	GLY	2.5
1	A	361	PHE	2.4
1	A	145	ILE	2.3
1	A	2	HIS	2.1
1	B	318	GLY	2.1
1	B	289	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)


There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	C8E	B	407	17/21	0.78	0.23	43,55,63,65	0
4	C8E	A	405	21/21	0.81	0.17	28,56,62,66	0
4	C8E	B	405	17/21	0.83	0.18	28,38,44,45	0
4	C8E	B	406	12/21	0.84	0.30	26,31,39,43	0
4	C8E	A	406	13/21	0.88	0.21	36,45,51,53	0
5	PGV	B	408	26/51	0.88	0.19	30,41,58,65	0
5	PGV	A	409	24/51	0.88	0.17	32,41,52,58	0
4	C8E	B	404	21/21	0.89	0.18	21,35,43,47	0
4	C8E	A	404	10/21	0.92	0.20	24,31,39,42	0
3	SO4	B	402	5/5	0.93	0.10	43,47,63,65	0
5	PGV	A	407	26/51	0.94	0.18	39,55,68,69	0
3	SO4	B	403	5/5	0.94	0.30	44,48,77,81	0
3	SO4	A	403	5/5	0.95	0.12	43,48,61,69	0
5	PGV	A	408	22/51	0.96	0.16	35,54,63,67	0
2	BDP	B	401	13/13	0.96	0.20	19,24,29,32	0
2	BDP	A	401	13/13	0.97	0.13	23,32,37,41	0
3	SO4	A	402	5/5	0.98	0.14	35,40,42,46	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.