Full wwPDB X-ray Structure Validation Report (i)

Jun 18, 2024 – 12:11 AM EDT

PDB ID	:	3FOM
Title	:	Crystal structure of the Class I MHC Molecule H-2Kwm7 with a Single Self Peptide IQQSIERL
Authors	:	Brims, D.R.; Qian, J.; Jarchum, I.; Yamada, T.; Mikesh, L.; Palmieri, E.; Lund, T.; Hattori, M.; Shabanowitz, J.; Hunt, D.F.; Ramagopal, U.A.; Malashkevich, V.N.; Almo, S.C.; Nathenson, S.G.; DiLorenzo, T.P.
Deposited on	:	2008-12-30
Resolution	:	2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	9% 70% 25% 5%
2	B	100	7% 74% 24% •
3	P	8	62% 38%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 3344 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	274	2227	1395	403	420	9	0	1	0

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	829	529	139	153	8	1	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	EXPRESSION TAG	UNP P01887

- Molecule 3 is a protein called 8 residue synthetic peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	P	8	69	42	13	14	0	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
4	A	1	1	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	218	218	218	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MHC

Chain A:



- Molecule 2: Beta-2-microglobulin

Chain B:



- Molecule 3: 8 residue synthetic peptide

Chain P:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	106.49Å 70.65Å 72.89Å 90.00° 104.21° 90.00°	Depositor
Resolution (Å)	19.82 – 2.10 19.82 – 2.10	Depositor EDS
% Data completeness (in resolution range)	90.3 (19.82-2.10) 90.3 (19.82-2.10)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.78 (at 2.09Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.222 , 0.289 0.220 , 0.277	Depositor DCC
R_{free} test set	1404 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å^2)	41.7	Xtriage
Anisotropy	0.074	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.31 , 48.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	3344	wwPDB-VP
Average B, all atoms (Å^2)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.80% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.56	0/2289	0.68	0/3107
2	B	0.83	1/855 (0.1%)	0.69	1/1158 (0.1%)
3	P	0.65	0/68	0.72	0/88
All	All	0.65	1/3212 (0.0%)	0.68	1/4353 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	99	MET	C-O	-16.70	0.91	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	99	MET	CA-C-O	8.21	137.35	120.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2227	0	2125	102	2
2	B	829	0	805	31	0
3	P	69	0	75	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	1	0	0	0	0
5	A	218	0	0	21	1
All	All	3344	0	3005	131	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All (131) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:48:LYS:N	2:B:48:LYS:HE2	1.29	1.41
2:B:48:LYS:H	2:B:48:LYS:CE	1.56	1.16
1:A:230:LEU:HD12	1:A:230:LEU:O	1.45	1.13
2:B:48:LYS:N	2:B:48:LYS:CE	2.10	1.13
1:A:98:MET:HB3	5:A:479:HOH:O	1.51	1.10
1:A:194:ARG:HH11	1:A:195:PRO:HD2	1.08	1.10
2:B:48:LYS:HE2	2:B:48:LYS:CA	1.87	1.04
1:A:154:GLU:N	1:A:154:GLU:OE1	1.95	0.99
1:A:222:GLU:HB2	5:A:382:HOH:O	1.61	0.98
1:A:56:GLU:H	1:A:56:GLU:CD	1.61	0.96
5:A:595:HOH:O	2:B:48:LYS:NZ	1.98	0.96
1:A:35:ARG:HG3	2:B:53:ASP:OD2	1.66	0.95
2:B:48:LYS:HE2	2:B:48:LYS:H	0.80	0.93
1:A:194:ARG:HB2	1:A:195:PRO:CD	2.03	0.88
1:A:194:ARG:NH1	1:A:195:PRO:HD2	1.89	0.87
1:A:111:ARG:NH2	1:A:128:GLU:HG3	1.90	0.86
5:A:371:HOH:O	2:B:38:GLN:HG2	1.75	0.86
1:A:131:LYS:NZ	5:A:640:HOH:O	1.64	0.86
5:A:606:HOH:O	3:P:4:SER:CB	2.23	0.85
1:A:23:MET:HE1	1:A:37:ASP:OD2	1.77	0.85
1:A:194:ARG:HH11	1:A:195:PRO:CD	1.89	0.85
1:A:194:ARG:HB2	1:A:195:PRO:HD2	1.59	0.83
1:A:9:VAL:HG22	1:A:24:GLU:HG2	1.60	0.82
5:A:411:HOH:O	2:B:53:ASP:HA	1.79	0.81
1:A:195:PRO:C	1:A:196:LYS:HD3	2.02	0.79
5:A:606:HOH:O	3:P:4:SER:HB2	1.81	0.79
1:A:21:ARG:HH11	1:A:21:ARG:HG3	1.48	0.78
1:A:195:PRO:HB2	1:A:196:LYS:HD3	1.63	0.78
1:A:21:ARG:NH1	1:A:39:ASP:OD2	2.18	0.77
1:A:35:ARG:HD3	2:B:53:ASP:OD2	1.84	0.77
1:A:169:ARG:O	1:A:173:GLU:HG2	1.84	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:230:LEU:HD12	1:A:230:LEU:C	2.06	0.76
1:A:35:ARG:CG	2:B:53:ASP:OD2	2.34	0.74
1:A:170[A]:ARG:HH21	1:A:170[A]:ARG:HG2	1.53	0.73
1:A:23:MET:CE	1:A:37:ASP:OD2	2.36	0.73
1:A:56:GLU:CD	1:A:56:GLU:N	2.41	0.72
1:A:103:VAL:HG13	1:A:108:ARG:O	1.88	0.72
5:A:563:HOH:O	2:B:54:MET:CB	2.37	0.71
1:A:78:LEU:HD13	1:A:95:ILE:HG13	1.74	0.69
1:A:35:ARG:CD	2:B:53:ASP:OD2	2.40	0.69
1:A:167:TRP:CE3	1:A:170[A]:ARG:HD2	2.28	0.68
1:A:3:HIS:ND1	1:A:29:ASP:OD2	2.22	0.68
5:A:299:HOH:O	2:B:95:TRP:O	2.13	0.67
1:A:230:LEU:O	1:A:230:LEU:CD1	2.35	0.67
2:B:38:GLN:HG3	2:B:81:ARG:HB3	1.77	0.66
1:A:219:LEU:N	1:A:224:LEU:HD21	2.11	0.66
1:A:194:ARG:HD3	1:A:195:PRO:HD2	1.78	0.65
1:A:230:LEU:C	1:A:230:LEU:CD1	2.64	0.65
1:A:198:LYS:HE2	1:A:198:LYS:HA	1.80	0.64
1:A:182:THR:HG21	1:A:265:GLY:HA2	1.79	0.63
1:A:51:TRP:HA	1:A:54:GLN:HE21	1.65	0.62
1:A:223:GLU:C	1:A:224:LEU:HD22	2.22	0.60
1:A:21:ARG:HG3	1:A:21:ARG:NH1	2.16	0.59
1:A:219:LEU:H	1:A:224:LEU:HD21	1.66	0.59
1:A:47:PRO:HG3	1:A:60:TRP:CZ2	2.37	0.59
1:A:84:TYR:OH	1:A:146:LYS:HE2	2.03	0.58
1:A:196:LYS:HE2	1:A:198:LYS:HG3	1.85	0.58
2:B:48:LYS:HE2	2:B:48:LYS:HA	1.81	0.58
1:A:195:PRO:CB	1:A:196:LYS:HD3	2.34	0.58
1:A:224:LEU:HD22	1:A:224:LEU:N	2.20	0.57
1:A:184:SER:HB3	1:A:265:GLY:O	2.03	0.57
1:A:170[A]:ARG:HH21	1:A:170[A]:ARG:CG	2.15	0.57
1:A:194:ARG:O	1:A:195:PRO:O	2.23	0.57
1:A:196:LYS:HD3	1:A:196:LYS:N	2.20	0.57
1:A:167:TRP:CZ3	1:A:170[A]:ARG:HD2	2.40	0.56
1:A:196:LYS:HG2	1:A:198:LYS:HG2	1.86	0.56
1:A:264:GLU:HG2	5:A:316:HOH:O	2.06	0.56
1:A:135:ALA:HB3	1:A:141:LEU:CD1	2.36	0.55
1:A:182:THR:HG21	1:A:265:GLY:CA	2.36	0.55
1:A:35:ARG:NH2	1:A:46:GLU:OE1	2.40	0.54
1:A:126:LEU:HB2	1:A:133:TRP:CZ3	2.42	0.54
1:A:21:ARG:HH22	1:A:23:MET:HE1	1.73	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:32:PRO:HB2	2:B:33:PRO:HD2	1.90	0.54
1:A:194:ARG:HG3	1:A:198:LYS:HB2	1.90	0.54
1:A:195:PRO:HB2	1:A:196:LYS:CD	2.37	0.54
5:A:390:HOH:O	2:B:34:HIS:HD2	1.91	0.54
5:A:397:HOH:O	2:B:77:THR:HG23	2.09	0.53
1:A:40:ALA:HB3	1:A:43:PRO:HA	1.91	0.52
1:A:224:LEU:N	1:A:224:LEU:CD2	2.72	0.51
1:A:29:ASP:O	1:A:30:ASP:CB	2.58	0.51
1:A:182:THR:CG2	1:A:265:GLY:HA3	2.40	0.51
1:A:170[A]:ARG:CG	1:A:170[A]:ARG:NH2	2.73	0.50
1:A:135:ALA:HB3	1:A:141:LEU:HD12	1.93	0.50
1:A:33:PHE:CD2	1:A:34:VAL:HG13	2.47	0.50
1:A:194:ARG:HB2	1:A:195:PRO:HD3	1.92	0.50
1:A:29:ASP:O	1:A:30:ASP:HB2	2.11	0.50
1:A:138:MET:HB2	5:A:349:HOH:O	2.12	0.49
1:A:56:GLU:N	1:A:56:GLU:OE2	2.35	0.49
1:A:111:ARG:HH21	1:A:128:GLU:HG3	1.71	0.49
1:A:195:PRO:O	1:A:196:LYS:HG2	2.14	0.48
2:B:32:PRO:CB	2:B:33:PRO:HD2	2.43	0.48
1:A:78:LEU:CD1	1:A:95:ILE:HG13	2.42	0.48
2:B:59:ASP:O	2:B:60:TRP:HB2	2.13	0.48
5:A:563:HOH:O	2:B:54:MET:HB2	2.05	0.48
1:A:99:ARG:HG3	1:A:114:GLN:HG2	1.96	0.47
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.49	0.47
1:A:194:ARG:CB	1:A:195:PRO:CD	2.79	0.47
1:A:135:ALA:CB	1:A:141:LEU:HD12	2.45	0.47
1:A:129:ASP:OD2	1:A:130:LEU:N	2.49	0.46
1:A:51:TRP:HA	1:A:54:GLN:NE2	2.30	0.46
1:A:21:ARG:HH22	1:A:23:MET:CE	2.29	0.45
5:A:374:HOH:O	2:B:0:MET:O	2.21	0.45
1:A:196:LYS:CG	1:A:198:LYS:HG2	2.46	0.45
1:A:99:ARG:HD3	1:A:114:GLN:OE1	2.15	0.45
1:A:1:GLY:O	1:A:105:SER:HB3	2.18	0.44
5:A:563:HOH:O	2:B:54:MET:HB3	2.07	0.44
1:A:194:ARG:HD2	1:A:198:LYS:HB2	1.99	0.44
1:A:182:THR:CG2	1:A:265:GLY:CA	2.96	0.44
1:A:194:ARG:HD2	1:A:198:LYS:CB	2.46	0.44
1:A:227:ASP:N	1:A:227:ASP:OD1	2.51	0.44
1:A:35:ARG:HE	1:A:35:ARG:HB3	1.47	0.44
2:B:21:ASN:HB3	2:B:70:PHE:CE1	2.52	0.43
5:A:575:HOH:O	2:B:0:MET:CE	2.67	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:73:THR:HG1	2:B:76:ASP:CG	2.18	0.43
1:A:203:CYS:HB2	1:A:217:TRP:CZ2	2.54	0.43
1:A:230:LEU:CD1	5:A:311:HOH:O	2.67	0.43
1:A:108:ARG:HB3	5:A:340:HOH:O	2.19	0.43
1:A:75:ARG:NH1	1:A:79:ARG:NH1	2.66	0.42
1:A:230:LEU:HD11	5:A:311:HOH:O	2.19	0.42
1:A:82:LEU:HD12	1:A:87:GLN:HB2	2.00	0.42
1:A:131:LYS:HE2	1:A:157:ARG:NH1	2.35	0.42
1:A:217:TRP:O	1:A:224:LEU:HD23	2.19	0.42
1:A:229:GLU:O	1:A:245:ALA:HA	2.20	0.42
1:A:55:VAL:CG1	1:A:174:LEU:HD11	2.49	0.42
1:A:43:PRO:O	1:A:44:ARG:HB3	2.20	0.41
1:A:195:PRO:C	1:A:196:LYS:CD	2.81	0.41
3:P:2:GLN:HG3	3:P:3:GLN:O	2.20	0.41
2:B:46:ILE:HG23	2:B:47:PRO:HD2	2.02	0.41
2:B:71:THR:HA	2:B:72:PRO:HD2	1.93	0.41
2:B:19:LYS:O	2:B:20:PRO:C	2.58	0.41
1:A:174:LEU:HA	1:A:174:LEU:HD23	1.86	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:GLU:OE1	1:A:61:GLU:OE1[2_555]	1.99	0.21
1:A:149:GLN:OE1	5:A:644:HOH:O[2_555]	2.16	0.04

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	273/274 (100%)	255 (93%)	16 (6%)	2 (1%)	22 18

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	98/100 (98%)	93 (95%)	5 (5%)	0	100	100
3	P	6/8 (75%)	6 (100%)	0	0	100	100
All	All	377/382 (99%)	354 (94%)	21 (6%)	2 (0%)	29	26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	195	PRO
1	A	221	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/230 (100%)	210 (91%)	21 (9%)	9	6
2	B	95/95 (100%)	91 (96%)	4 (4%)	30	30
3	P	8/8 (100%)	8 (100%)	0	100	100
All	All	334/333 (100%)	309 (92%)	25 (8%)	13	10

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	ARG
1	A	35	ARG
1	A	41	GLU
1	A	44	ARG
1	A	45	TYR
1	A	56	GLU
1	A	61	GLU
1	A	78	LEU
1	A	111	ARG
1	A	114	GLN
1	A	128	GLU
1	A	182	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	194	ARG
1	A	196	LYS
1	A	220	ASN
1	A	224	LEU
1	A	227	ASP
1	A	228	MET
1	A	230	LEU
1	A	268	GLU
1	A	270	LEU
2	B	48	LYS
2	B	49	VAL
2	B	58	LYS
2	B	70	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	A	72	GLN
1	A	115	GLN
1	A	145	HIS
1	A	149	GLN
1	A	220	ASN
1	A	256	ASN
2	B	34	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	274/274 (100%)	0.49	24 (8%) 10 12	17, 28, 48, 60	0
2	B	100/100 (100%)	0.43	7 (7%) 16 20	19, 29, 41, 44	3 (3%)
3	P	8/8 (100%)	-0.11	0 100 100	37, 40, 44, 44	0
All	All	382/382 (100%)	0.46	31 (8%) 12 15	17, 29, 45, 60	3 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	220	ASN	4.5
1	A	196	LYS	4.2
1	A	17	HIS	3.7
2	B	23	LEU	3.5
1	A	16	GLY	3.4
1	A	195	PRO	3.3
1	A	197	ASP	3.2
1	A	219	LEU	3.2
2	B	48	LYS	3.1
1	A	222	GLU	3.1
1	A	226	GLN	3.0
1	A	227	ASP	2.9
1	A	54	GLN	2.8
1	A	177	ALA	2.7
1	A	194	ARG	2.7
2	B	47	PRO	2.6
1	A	201	LEU	2.6
2	B	74	GLU	2.5
1	A	255	GLN	2.5
1	A	18	GLY	2.5
1	A	21	ARG	2.4
1	A	247	VAL	2.4
1	A	224	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	34	VAL	2.4
2	B	43	GLY	2.3
2	B	75	THR	2.3
1	A	230	LEU	2.3
1	A	225	THR	2.1
1	A	58	GLU	2.1
1	A	1	GLY	2.0
2	B	44	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	CL	A	275	1/1	0.86	0.53	72,72,72,72	0

6.5 Other polymers (i)

There are no such residues in this entry.