



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 7, 2020 – 10:10 am BST

PDB ID : 6FOY
Title : The crystal structure of *P.fluorescens* Kynurenine 3-monooxygenase (KMO) in complex with competitive inhibitor No. 9
Authors : Levy, C.W.; Leys, D.
Deposited on : 2018-02-08
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	
1	B	460	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 14997 atoms, of which 7048 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Kynurenine 3-monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	445	7004	2202	3490	655	639	18	0	10	0
1	B	447	6966	2191	3468	652	637	18	0	4	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	251	SER	CYS	engineered mutation	UNP Q84HF5
A	460	SER	-	expression tag	UNP Q84HF5
B	251	SER	CYS	engineered mutation	UNP Q84HF5
B	460	SER	-	expression tag	UNP Q84HF5

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	A	1	Total	C	H	N	O	P	0	0
			84	27	31	9	15	2		
2	B	1	Total	C	H	N	O	P	0	0
			84	27	31	9	15	2		

- Molecule 3 is 5-[2,3-bis(chloranyl)phenyl]furan-2-carboxylic acid (three-letter code: E0T) (formula: C₁₁H₆Cl₂O₃) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
3	A	1	Total	C	Cl	O	0	0
			16	11	2	3		
3	B	1	Total	C	Cl	O	0	0
			16	11	2	3		

- Molecule 4 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	24	6	14	4	0	0
4	B	1	24	6	14	4	0	0

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
5	A	1	1	1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	402	402	402	0	0
6	B	376	376	376	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kynurenine 3-monoxygenase



- Molecule 1: Kynurenine 3-monoxygenase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.80Å 52.98Å 135.76Å 90.00° 103.83° 90.00°	Depositor
Resolution (Å)	67.78 – 1.65 67.78 – 1.65	Depositor EDS
% Data completeness (in resolution range)	98.2 (67.78-1.65) 91.0 (67.78-1.65)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 1.65Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.164 , 0.213 0.164 , 0.214	Depositor DCC
R_{free} test set	1976 reflections (1.72%)	wwPDB-VP
Wilson B-factor (Å ²)	11.8	Xtrriage
Anisotropy	0.529	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 50.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14997	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.83 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.7054e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PGE, E0T, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/3616	0.67	3/4909 (0.1%)
1	B	0.47	0/3581	0.62	1/4858 (0.0%)
All	All	0.48	0/7197	0.65	4/9767 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	172	ARG	NE-CZ-NH2	-7.05	116.77	120.30
1	A	172	ARG	NE-CZ-NH1	6.05	123.33	120.30
1	A	165	ASP	CB-CG-OD1	5.87	123.58	118.30
1	B	165	ASP	CB-CG-OD1	5.04	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3514	3490	3475	32	0
1	B	3498	3468	3473	31	0
2	A	53	31	31	0	0
2	B	53	31	31	0	0
3	A	16	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	16	0	0	0	0
4	A	10	14	14	1	0
4	B	10	14	14	2	0
5	A	1	0	0	0	0
6	A	402	0	0	12	2
6	B	376	0	0	10	0
All	All	7949	7048	7038	63	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:38:ARG:NH2	6:B:601:HOH:O	2.02	0.93
1:A:281:GLN:OE1	6:A:601:HOH:O	1.88	0.91
1:A:271:ASP:OD1	6:A:602:HOH:O	2.08	0.72
1:B:421:GLN:OE1	6:B:602:HOH:O	2.11	0.68
1:A:448:GLU:OE2	1:A:451:ARG:NH2	2.24	0.68
1:B:383:LEU:HD23	1:B:431:VAL:HG13	1.74	0.68
1:A:47:ALA:O	6:A:603:HOH:O	2.12	0.68
1:A:417:MET:O	1:A:421:GLN:HG3	1.93	0.67
1:B:194:GLU:OE2	6:B:603:HOH:O	2.14	0.65
1:A:6:ALA:N	6:A:607:HOH:O	2.31	0.64
1:B:76:GLU:OE2	1:B:112:ARG:NH2	2.35	0.59
1:B:196:GLN:NE2	6:B:609:HOH:O	2.35	0.59
1:A:147:SER:HB3	1:A:153:ARG:HD3	1.85	0.58
1:A:437:LEU:HD12	1:A:440:ILE:HD12	1.87	0.57
1:A:257:ASP:OD2	6:A:604:HOH:O	2.17	0.57
1:A:390:ILE:HD12	1:A:437:LEU:HD21	1.87	0.57
1:B:188:GLN:O	1:B:290:LYS:NZ	2.39	0.56
1:A:244:ALA:O	1:A:245:ALA:HB3	2.06	0.56
1:B:367:GLU:OE2	6:B:604:HOH:O	2.18	0.55
1:B:183:GLU:HG2	1:B:296:LEU:HD22	1.90	0.54
1:B:112:ARG:HG3	6:B:915:HOH:O	2.07	0.54
1:B:290:LYS:HD3	1:B:291:LEU:O	2.07	0.53
1:A:297:THR:OG1	6:A:605:HOH:O	2.19	0.51
1:B:141:ARG:NE	6:B:619:HOH:O	2.42	0.51
1:B:394:ARG:HB3	1:B:446:GLU:OE2	2.11	0.51
1:B:85:VAL:HG12	1:B:87:VAL:HG23	1.94	0.50
1:B:282:ASP:O	1:B:285:HIS:O	2.28	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:50:ARG:O	1:B:50:ARG:HG3	2.11	0.49
1:A:16:LEU:HD13	1:A:323[A]:MET:HG3	1.93	0.49
1:A:221:MET:HE3	1:A:223:ILE:HG23	1.95	0.48
1:B:58:GLU:OE2	1:B:410[B]:ARG:NH1	2.47	0.47
1:B:290:LYS:HD3	1:B:291:LEU:N	2.29	0.47
1:B:7[B]:ARG:HG3	1:B:30:TRP:CE2	2.49	0.47
1:B:335:GLU:HG3	6:B:815:HOH:O	2.15	0.46
1:B:257:ASP:C	1:B:257:ASP:OD1	2.54	0.46
1:A:153:ARG:O	1:A:154:LEU:HD23	2.17	0.45
1:A:214:ILE:HD12	1:A:222:CYS:O	2.16	0.45
1:B:51:SER:OG	1:B:109:ASN:ND2	2.50	0.45
1:A:112:ARG:HG3	6:A:925:HOH:O	2.17	0.45
1:A:424:GLU:HG2	6:A:906:HOH:O	2.17	0.45
1:B:28:ASN:ND2	6:B:608:HOH:O	2.35	0.45
1:B:52:ILE:HD11	1:B:110:ARG:HH21	1.81	0.44
1:A:335:GLU:HG3	6:A:858:HOH:O	2.16	0.44
1:A:192:TYR:HA	1:A:236:LEU:O	2.17	0.44
1:B:71:ARG:NH1	6:B:610:HOH:O	2.38	0.44
1:A:243:PRO:HD3	6:A:734:HOH:O	2.18	0.44
1:A:161:LEU:O	1:A:305:ALA:HA	2.19	0.43
1:B:156:LYS:HE3	4:B:503:PGE:H32	2.00	0.43
1:A:244:ALA:O	1:A:245:ALA:CB	2.67	0.43
1:A:52:ILE:HD11	1:A:194[A]:GLU:OE1	2.19	0.43
1:A:71:ARG:NH2	1:A:75:ALA:HB2	2.34	0.43
1:A:156:LYS:NZ	4:A:503:PGE:H32	2.34	0.42
1:A:434:HIS:ND1	1:A:439:SER:OG	2.38	0.42
1:A:154:LEU:HD11	6:A:926:HOH:O	2.18	0.42
1:A:373:SER:HA	6:A:787:HOH:O	2.19	0.42
1:B:192:TYR:HA	1:B:236:LEU:O	2.20	0.41
1:B:256:VAL:O	1:B:257:ASP:OD1	2.39	0.41
1:A:224:ALA:HB2	1:A:234:VAL:HG22	2.03	0.41
1:B:285:HIS:O	1:B:286:HIS:HB2	2.21	0.41
1:A:183:GLU:HG2	1:A:296:LEU:HD22	2.02	0.41
1:A:182:GLY:O	1:A:297:THR:HG23	2.21	0.40
1:B:97:TYR:HB3	1:B:408:PHE:CZ	2.56	0.40
1:B:156:LYS:NZ	4:B:503:PGE:H52	2.36	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:641:HOH:O	6:A:786:HOH:O[2_252]	1.90	0.30
6:A:641:HOH:O	6:A:680:HOH:O[2_252]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	451/460 (98%)	441 (98%)	9 (2%)	1 (0%)	47	28
1	B	446/460 (97%)	433 (97%)	12 (3%)	1 (0%)	47	28
All	All	897/920 (98%)	874 (97%)	21 (2%)	2 (0%)	47	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	286	HIS
1	A	246	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/371 (99%)	364 (99%)	4 (1%)	73	57
1	B	366/371 (99%)	361 (99%)	5 (1%)	67	46
All	All	734/742 (99%)	725 (99%)	9 (1%)	69	53

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	ARG
1	A	71	ARG
1	A	281	GLN
1	A	385	ARG
1	B	51	SER
1	B	157	ARG
1	B	257	ASP
1	B	290	LYS
1	B	389	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	B	501	-	51,58,58	1.33	6 (11%)	60,89,89	2.07	7 (11%)
2	FAD	A	501	-	51,58,58	1.19	3 (5%)	60,89,89	2.20	7 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	E0T	B	502	-	9,17,17	0.66	0	13,24,24	1.72	3 (23%)
4	PGE	A	503	-	9,9,9	0.33	0	8,8,8	0.49	0
4	PGE	B	503	-	9,9,9	0.29	0	8,8,8	0.58	0
3	E0T	A	502	-	9,17,17	1.12	1 (11%)	13,24,24	1.77	5 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	501	-	-	2/30/50/50	0/6/6/6
2	FAD	A	501	-	-	2/30/50/50	0/6/6/6
3	E0T	B	502	-	-	0/1/8/8	0/2/2/2
4	PGE	A	503	-	-	2/7/7/7	-
4	PGE	B	503	-	-	5/7/7/7	-
3	E0T	A	502	-	-	0/1/8/8	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	FAD	C4X-C10	5.26	1.44	1.38
2	A	501	FAD	C4X-C10	4.55	1.43	1.38
2	B	501	FAD	C10-N1	-3.84	1.28	1.33
2	A	501	FAD	C4-N3	3.39	1.38	1.33
2	B	501	FAD	C5X-N5	3.04	1.40	1.35
2	B	501	FAD	C4-N3	2.87	1.38	1.33
3	A	502	E0T	C12-C14	-2.50	1.36	1.39
2	B	501	FAD	C9A-N10	2.18	1.41	1.38
2	B	501	FAD	C4-C4X	2.18	1.45	1.41
2	A	501	FAD	C4-C4X	2.11	1.45	1.41

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	FAD	C4-N3-C2	12.59	125.77	115.14
2	B	501	FAD	C4-N3-C2	11.66	124.99	115.14
2	B	501	FAD	C4X-C4-N3	-6.83	114.09	123.43
2	A	501	FAD	C4X-C4-N3	-6.77	114.17	123.43
2	A	501	FAD	C10-C4X-N5	4.63	124.46	121.26
2	B	501	FAD	C10-C4X-N5	4.42	124.31	121.26

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	FAD	C4-C4X-C10	-4.20	117.17	119.95
3	A	502	E0T	C08-C14-CL2	3.96	123.11	119.66
3	B	502	E0T	C08-C14-CL2	3.78	122.95	119.66
2	B	501	FAD	C4-C4X-C10	-3.69	117.51	119.95
2	A	501	FAD	C4X-C10-N10	-3.64	116.56	120.30
2	B	501	FAD	C4X-C10-N10	-3.17	117.05	120.30
3	B	502	E0T	C09-C08-C07	-3.09	116.51	120.42
2	B	501	FAD	C1'-N10-C9A	2.93	120.60	118.29
2	A	501	FAD	C5A-C6A-N6A	2.68	124.42	120.35
2	A	501	FAD	C1'-N10-C9A	2.60	120.34	118.29
3	A	502	E0T	C11-C12-C14	2.60	123.42	120.58
3	A	502	E0T	C14-C12-CL1	-2.30	118.28	120.52
3	A	502	E0T	C12-C14-CL2	-2.22	117.17	120.02
3	A	502	E0T	C09-C08-C07	-2.18	117.66	120.42
2	B	501	FAD	C5A-C6A-N6A	2.18	123.67	120.35
3	B	502	E0T	C11-C12-C14	2.06	122.83	120.58

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	503	PGE	O2-C3-C4-O3
4	B	503	PGE	O2-C3-C4-O3
4	B	503	PGE	O1-C1-C2-O2
4	B	503	PGE	C3-C4-O3-C5
4	B	503	PGE	C6-C5-O3-C4
4	B	503	PGE	C1-C2-O2-C3
2	A	501	FAD	O4B-C4B-C5B-O5B
2	B	501	FAD	O4B-C4B-C5B-O5B
2	B	501	FAD	C1'-C2'-C3'-O3'
2	A	501	FAD	C1'-C2'-C3'-O3'
4	A	503	PGE	O1-C1-C2-O2


There are no ring outliers.


2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	503	PGE	1	0
4	B	503	PGE	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	445/460 (96%)	-0.12	16 (3%) 42 43	7, 18, 46, 59	0
1	B	447/460 (97%)	-0.17	10 (2%) 62 63	10, 20, 44, 58	0
All	All	892/920 (96%)	-0.14	26 (2%) 51 52	7, 19, 45, 59	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	245	ALA	6.1
1	B	50	ARG	5.0
1	B	48	ARG	4.8
1	A	89	GLY	4.1
1	A	248	ALA	3.6
1	A	379	PRO	3.5
1	A	246	GLN	3.5
1	B	457	SER	3.5
1	A	48	ARG	3.0
1	A	99	ARG	2.9
1	A	49	GLY	2.9
1	A	381	TYR	2.9
1	B	49	GLY	2.7
1	A	385	ARG	2.7
1	B	393	GLN	2.7
1	A	438	THR	2.6
1	B	256	VAL	2.6
1	A	437	LEU	2.5
1	A	91	PRO	2.4
1	A	50	ARG	2.3
1	A	285	HIS	2.3
1	B	248	ALA	2.3
1	B	99	ARG	2.3
1	B	377	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	433	ASN	2.1
1	B	444	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PGE	A	503	10/10	0.82	0.15	37,45,53,53	0
4	PGE	B	503	10/10	0.89	0.12	39,47,52,53	0
3	E0T	A	502	16/16	0.97	0.08	21,24,26,27	0
3	E0T	B	502	16/16	0.98	0.07	15,18,21,22	0
2	FAD	B	501	53/53	0.99	0.07	8,11,23,23	0
2	FAD	A	501	53/53	0.99	0.08	5,9,16,17	0
5	CA	A	504	1/1	0.99	0.15	35,35,35,35	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around E0T A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around E0T B 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.