



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 13, 2020 – 10:25 pm BST

PDB ID : 6FQX
Title : Plasmodium falciparum 6-phosphogluconate dehydrogenase in its apo form, in complex with its cofactor NADP+ and in complex with its substrate 6-phosphogluconate
Authors : Fritz-Wolf, K.; Haeussler, K.; Reichmann, M.; Rahlfs, S.; Becker, K.
Deposited on : 2018-02-15
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	468	
1	B	468	
1	C	468	
1	D	468	
1	E	468	
1	F	468	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	468	 % 73% 25% •
1	H	468	 3% 66% 33% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PEG	A	504	-	-	-	X
3	PEG	C	502	-	-	-	X
5	AE3	D	502	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 29994 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 6-phosphogluconate dehydrogenase, decarboxylating.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	468	3715	2362	627	702	24	0	0	0
1	B	468	3715	2362	627	702	24	0	0	0
1	C	468	3715	2362	627	702	24	0	0	0
1	D	468	3715	2362	627	702	24	0	0	0
1	E	468	3715	2362	627	702	24	0	0	0
1	F	468	3715	2362	627	702	24	0	0	0
1	G	468	3715	2362	627	702	24	0	0	0
1	H	468	3715	2362	627	702	24	0	0	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	E	1	Total	C	O	0	0
			6	3	3		
2	E	1	Total	C	O	0	0
			6	3	3		
2	F	1	Total	C	O	0	0
			6	3	3		
2	F	1	Total	C	O	0	0
			6	3	3		
2	F	1	Total	C	O	0	0
			6	3	3		
2	H	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 7 4 3	0	0
3	A	1	Total C O 7 4 3	0	0
3	B	1	Total C O 7 4 3	0	0
3	C	1	Total C O 7 4 3	0	0
3	E	1	Total C O 7 4 3	0	0
3	E	1	Total C O 7 4 3	0	0
3	F	1	Total C O 7 4 3	0	0
3	H	1	Total C O 7 4 3	0	0
3	H	1	Total C O 7 4 3	0	0

- Molecule 4 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		
4	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is 2-(2-ETHOXYETHOXY)ETHANOL (three-letter code: AE3) (formula: $C_6H_{14}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			9	6	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	12	Total O 12 12	0	0
6	B	21	Total O 21 21	0	0
6	C	11	Total O 11 11	0	0
6	D	10	Total O 10 10	0	0
6	E	14	Total O 14 14	0	0
6	F	13	Total O 13 13	0	0
6	G	12	Total O 12 12	0	0
6	H	5	Total O 5 5	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 6-phosphogluconate dehydrogenase, decarboxylating




- Molecule 1: 6-phosphogluconate dehydrogenase, decarboxylating




- Molecule 1: 6-phosphogluconate dehydrogenase, decarboxylating






- Molecule 1: 6-phosphogluconate dehydrogenase, decarboxylating




- Molecule 1: 6-phosphogluconate dehydrogenase, decarboxylating




- Molecule 1: 6-phosphogluconate dehydrogenase, decarboxylating







- Molecule 1: 6-phosphogluconate dehydrogenase, decarboxylating



- Molecule 1: 6-phosphogluconate dehydrogenase, decarboxylating





4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	134.45Å 134.45Å 406.05Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.53 – 2.80 47.69 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.6 (46.53-2.80) 92.5 (47.69-2.80)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.75 (at 2.81Å)	Xtrriage
Refinement program	PHENIX (dev_2645: ???)	Depositor
R, R_{free}	0.188 , 0.242 0.188 , 0.242	Depositor DCC
R_{free} test set	2023 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å ²)	63.5	Xtrriage
Anisotropy	0.377	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 56.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.047 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	29994	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, AE3, PEG, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/3781	0.72	9/5098 (0.2%)
1	B	0.53	1/3781 (0.0%)	0.69	4/5098 (0.1%)
1	C	0.49	0/3781	0.67	2/5098 (0.0%)
1	D	0.47	0/3781	0.65	0/5098
1	E	0.57	4/3781 (0.1%)	0.75	8/5098 (0.2%)
1	F	0.50	1/3781 (0.0%)	0.69	3/5098 (0.1%)
1	G	0.50	1/3781 (0.0%)	0.77	9/5098 (0.2%)
1	H	0.50	0/3781	0.80	13/5098 (0.3%)
All	All	0.51	7/30248 (0.0%)	0.72	48/40784 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	461	GLU	CG-CD	-7.02	1.41	1.51
1	E	372	PHE	CD2-CE2	6.29	1.51	1.39
1	E	372	PHE	CE2-CZ	6.22	1.49	1.37
1	E	59	GLU	CB-CG	-5.84	1.41	1.52
1	E	91	PHE	CE1-CZ	-5.66	1.26	1.37

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	460	ARG	NE-CZ-NH1	10.45	125.53	120.30
1	G	460	ARG	NE-CZ-NH2	-9.15	115.72	120.30
1	H	264	LYS	CB-CG-CD	-8.69	89.01	111.60
1	H	220	LYS	CA-CB-CG	8.41	131.91	113.40
1	E	59	GLU	CA-CB-CG	8.16	131.34	113.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	H	47	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3715	0	3733	87	0
1	B	3715	0	3733	80	0
1	C	3715	0	3733	85	0
1	D	3715	0	3733	78	0
1	E	3715	0	3733	108	0
1	F	3715	0	3733	80	0
1	G	3715	0	3733	134	0
1	H	3715	0	3731	135	0
2	A	12	0	16	1	0
2	B	24	0	32	1	0
2	C	6	0	8	0	0
2	D	6	0	8	1	0
2	E	12	0	16	1	0
2	F	18	0	24	1	0
2	H	6	0	8	0	0
3	A	14	0	20	0	0
3	B	7	0	10	0	0
3	C	7	0	10	0	0
3	E	14	0	20	2	0
3	F	7	0	10	0	0
3	H	14	0	20	0	0
4	A	10	0	14	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	10	0	14	0	0
5	D	9	0	14	4	0
6	A	12	0	0	0	0
6	B	21	0	0	1	0
6	C	11	0	0	0	0
6	D	10	0	0	2	0
6	E	14	0	0	0	0
6	F	13	0	0	1	0
6	G	12	0	0	0	0
6	H	5	0	0	1	0
All	All	29994	0	30106	712	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 712 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:264:LYS:HE2	1:H:271:THR:O	1.39	1.20
1:A:58:GLU:HA	1:A:61:ILE:HD13	1.31	1.12
1:G:308:ILE:HD12	1:G:309:LEU:N	1.64	1.12
1:F:12:MET:HE2	1:F:73:LEU:HD13	1.41	1.00
1:G:436:GLN:HG3	1:H:297:THR:HG22	1.44	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	466/468 (100%)	453 (97%)	13 (3%)	0	100 100
1	B	466/468 (100%)	452 (97%)	14 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	466/468 (100%)	453 (97%)	12 (3%)	1 (0%)	47	78
1	D	466/468 (100%)	454 (97%)	12 (3%)	0	100	100
1	E	466/468 (100%)	449 (96%)	14 (3%)	3 (1%)	25	56
1	F	466/468 (100%)	456 (98%)	10 (2%)	0	100	100
1	G	466/468 (100%)	454 (97%)	12 (3%)	0	100	100
1	H	466/468 (100%)	450 (97%)	16 (3%)	0	100	100
All	All	3728/3744 (100%)	3621 (97%)	103 (3%)	4 (0%)	51	81

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	304	ASN
1	E	304	ASN
1	E	352	LYS
1	E	370	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	405/406 (100%)	397 (98%)	8 (2%)	55	84
1	B	405/406 (100%)	398 (98%)	7 (2%)	60	87
1	C	405/406 (100%)	399 (98%)	6 (2%)	65	89
1	D	405/406 (100%)	400 (99%)	5 (1%)	71	92
1	E	405/406 (100%)	399 (98%)	6 (2%)	65	89
1	F	405/406 (100%)	400 (99%)	5 (1%)	71	92
1	G	405/406 (100%)	393 (97%)	12 (3%)	41	75
1	H	405/406 (100%)	396 (98%)	9 (2%)	52	83
All	All	3240/3248 (100%)	3182 (98%)	58 (2%)	59	86

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	12	MET
1	F	151	CYS
1	H	219	ASN
1	E	75	LYS
1	E	372	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	186	ASN
1	F	247	ASN
1	F	383	ASN
1	E	63	ASN
1	F	354	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	AE3	D	502	-	8,8,8	0.83	0	7,7,7	0.69	0
3	PEG	E	504	-	6,6,6	0.89	0	5,5,5	0.50	0
2	GOL	F	503	-	5,5,5	1.52	1 (20%)	5,5,5	0.58	0
4	PGE	A	505	-	9,9,9	0.86	0	8,8,8	0.30	0
3	PEG	F	504	-	6,6,6	0.73	0	5,5,5	0.49	0
3	PEG	E	503	-	6,6,6	0.87	0	5,5,5	0.45	0
2	GOL	D	501	-	5,5,5	0.82	0	5,5,5	1.02	0
2	GOL	E	501	-	5,5,5	1.10	0	5,5,5	0.87	0
2	GOL	A	502	-	5,5,5	0.78	0	5,5,5	0.98	0
4	PGE	B	506	-	9,9,9	0.97	0	8,8,8	0.59	0
3	PEG	A	503	-	6,6,6	0.87	0	5,5,5	0.44	0
2	GOL	B	502	-	5,5,5	1.25	1 (20%)	5,5,5	0.91	0
3	PEG	C	502	-	6,6,6	0.86	0	5,5,5	0.48	0
2	GOL	F	501	-	5,5,5	0.85	0	5,5,5	1.15	0
2	GOL	H	501	-	5,5,5	0.88	0	5,5,5	1.03	0
2	GOL	E	502	-	5,5,5	0.94	0	5,5,5	1.20	1 (20%)
2	GOL	B	503	-	5,5,5	0.89	0	5,5,5	1.02	0
2	GOL	B	504	-	5,5,5	0.96	0	5,5,5	1.01	0
2	GOL	F	502	-	5,5,5	0.78	0	5,5,5	1.14	0
2	GOL	B	501	-	5,5,5	0.84	0	5,5,5	1.43	1 (20%)
2	GOL	A	501	-	5,5,5	1.30	1 (20%)	5,5,5	1.00	0
2	GOL	C	501	-	5,5,5	0.75	0	5,5,5	1.13	0
3	PEG	B	505	-	6,6,6	0.85	0	5,5,5	0.46	0
3	PEG	H	503	-	6,6,6	0.86	0	5,5,5	0.32	0
3	PEG	A	504	-	6,6,6	0.80	0	5,5,5	0.23	0
3	PEG	H	502	-	6,6,6	1.00	0	5,5,5	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	AE3	D	502	-	-	4/6/6/6	-
3	PEG	E	504	-	-	1/4/4/4	-
2	GOL	F	503	-	-	2/4/4/4	-
4	PGE	A	505	-	-	4/7/7/7	-
3	PEG	F	504	-	-	1/4/4/4	-
3	PEG	E	503	-	-	2/4/4/4	-
2	GOL	D	501	-	-	0/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	E	501	-	-	2/4/4/4	-
2	GOL	A	502	-	-	2/4/4/4	-
4	PGE	B	506	-	-	2/7/7/7	-
3	PEG	A	503	-	-	1/4/4/4	-
2	GOL	B	502	-	-	0/4/4/4	-
3	PEG	C	502	-	-	2/4/4/4	-
2	GOL	F	501	-	-	0/4/4/4	-
2	GOL	H	501	-	-	0/4/4/4	-
2	GOL	E	502	-	-	0/4/4/4	-
2	GOL	B	503	-	-	0/4/4/4	-
2	GOL	B	504	-	-	2/4/4/4	-
2	GOL	F	502	-	-	4/4/4/4	-
2	GOL	B	501	-	-	2/4/4/4	-
2	GOL	A	501	-	-	4/4/4/4	-
2	GOL	C	501	-	-	4/4/4/4	-
3	PEG	B	505	-	-	0/4/4/4	-
3	PEG	H	503	-	-	3/4/4/4	-
3	PEG	A	504	-	-	1/4/4/4	-
3	PEG	H	502	-	-	4/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	503	GOL	C3-C2	2.53	1.62	1.51
2	A	501	GOL	C3-C2	2.18	1.60	1.51
2	B	502	GOL	C3-C2	2.11	1.60	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	GOL	C3-C2-C1	-2.44	102.21	111.70
2	E	502	GOL	C3-C2-C1	-2.02	103.84	111.70

There are no chirality outliers.

5 of 47 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	501	GOL	O1-C1-C2-C3
2	A	502	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	504	GOL	O1-C1-C2-C3
2	F	502	GOL	O1-C1-C2-O2
2	F	502	GOL	O1-C1-C2-C3

There are no ring outliers.

9 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	502	AE3	4	0
3	E	504	PEG	1	0
4	A	505	PGE	1	0
3	E	503	PEG	1	0
2	D	501	GOL	1	0
2	E	501	GOL	1	0
2	F	502	GOL	1	0
2	B	501	GOL	1	0
2	A	501	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	468/468 (100%)	0.04	6 (1%) 77 72	52, 70, 95, 110	0
1	B	468/468 (100%)	0.01	6 (1%) 77 72	51, 66, 87, 116	0
1	C	468/468 (100%)	0.00	11 (2%) 59 49	53, 68, 90, 124	0
1	D	468/468 (100%)	0.07	13 (2%) 53 43	53, 69, 91, 123	0
1	E	468/468 (100%)	-0.11	4 (0%) 84 80	51, 67, 87, 109	0
1	F	468/468 (100%)	-0.09	3 (0%) 89 86	52, 68, 90, 102	0
1	G	468/468 (100%)	0.02	4 (0%) 84 80	57, 76, 97, 123	0
1	H	468/468 (100%)	0.13	14 (2%) 50 40	59, 80, 102, 126	0
All	All	3744/3744 (100%)	0.01	61 (1%) 72 66	51, 70, 94, 126	0

The worst 5 of 61 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	309	LEU	5.1
1	G	308	ILE	4.5
1	B	313	ASN	4.2
1	C	310	ILE	4.2
1	C	307	ASN	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	B	502	6/6	0.53	0.30	93,101,106,111	0
3	PEG	E	503	7/7	0.65	0.38	76,82,99,103	0
5	AE3	D	502	9/9	0.70	0.41	71,76,81,85	0
2	GOL	A	501	6/6	0.70	0.36	86,90,93,94	0
2	GOL	F	503	6/6	0.71	0.38	67,75,76,84	0
3	PEG	H	503	7/7	0.73	0.26	70,71,86,87	0
3	PEG	C	502	7/7	0.74	0.45	79,80,86,88	0
3	PEG	A	504	7/7	0.74	0.42	93,94,103,105	0
2	GOL	E	502	6/6	0.81	0.24	72,74,86,93	0
4	PGE	B	506	10/10	0.81	0.28	61,76,81,83	0
3	PEG	H	502	7/7	0.81	0.47	79,82,87,88	0
3	PEG	A	503	7/7	0.82	0.24	71,76,81,87	0
3	PEG	B	505	7/7	0.83	0.16	82,84,88,99	0
2	GOL	H	501	6/6	0.83	0.31	82,87,97,99	0
3	PEG	E	504	7/7	0.84	0.44	71,78,81,85	0
2	GOL	B	504	6/6	0.86	0.27	76,78,83,86	0
4	PGE	A	505	10/10	0.87	0.33	69,73,80,81	0
2	GOL	E	501	6/6	0.87	0.20	78,84,86,89	0
3	PEG	F	504	7/7	0.91	0.35	75,77,87,89	0
2	GOL	A	502	6/6	0.91	0.24	62,64,71,75	0
2	GOL	B	503	6/6	0.93	0.23	75,81,85,85	0
2	GOL	F	501	6/6	0.95	0.24	64,66,69,70	0
2	GOL	B	501	6/6	0.96	0.33	63,66,68,74	0
2	GOL	F	502	6/6	0.96	0.24	65,69,72,72	0
2	GOL	D	501	6/6	0.97	0.30	64,67,70,70	0
2	GOL	C	501	6/6	0.98	0.29	63,65,69,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.