



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 10:01 pm BST

PDB ID : 4FSX
Title : crystal structure of Se-substituted Zea mays ZMET2 in complex with SAH
Authors : Du, J.; Patel, D.J.
Deposited on : 2012-06-27
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	784	<p>%</p> <p>21% 47% 17% • 13%</p>
1	B	784	<p>%</p> <p>19% 48% 18% • 14%</p>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 10631 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DNA (cytosine-5)-methyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	680	5331	3401	904	993	18	15	0	0	0
1	B	675	5248	3346	893	976	18	15	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	129	SER	-	EXPRESSION TAG	UNP Q9AXT8
B	129	SER	-	EXPRESSION TAG	UNP Q9AXT8

- Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: $C_{14}H_{20}N_6O_5S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	26	14	6	5	1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	B	1	26	14	6	5	1	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (cytosine-5)-methyltransferase 1




K785
L766
S767
S768
P771
L772
V773
P774
D775
Y776
M777
K784
S785
L786
P787
F788
G789
R791
L792
W793
W794
D795
E796
T797
F798
P799
T800
W801
V802
T803

R828
L829
Q830
D834
Y835
Y836
R837
L838
F839
I842
S140
G141
K843
E844
R845
I846
D847
G848
V849
I847
G850
S851
A852
W853
P856
V857
L861
F862
W863
D795
E796
T797
F798
P799
T800
W801
V802
T803
E874
G875
S876
D877
P878
L879
Y880
Q881
I812
I813
P883
P884
T814
S885
F886
T887
S888
V820
L821
G141
V185
I886
C187
S825
G141

GLN
ALA
ARG
ALA
SER
PRG
VAL
GLY
THR
PRO
ALA
GLY
GLU
VAL
VAL
GLN

● Molecule 1: DNA (cytosine-5)-methyltransferase 1



SER
ALA
GLY
ASP
H133
E134
P135
I136
F137
A201
G139
S140
P141
V142
A143
A144
I847
D145
E146
A147
G850
S149
G213
W150
D115
W151
K152
P152
K152
R154
Y155
F220
G156
R157
SER
SER
THR
ALA
ALA
LYS
LYS
PRO
ASP
GLU
GLU
GLU
GLU
L170
S235
K171
G174
H175
Y176
I239
S240
R177
S178
A179
K180
V181
D182
N183
G184
V184
V185
I886
C187
D191

V192
Y193
K194
K195
G197
I198
M199
E200
A201
Q202
S140
Y203
R206
I207
T208
E209
D210
F211
E212
G213
T214
D215
W151
K152
P152
K152
R154
Y155
F220
G156
R157
SER
SER
THR
ALA
ALA
LYS
LYS
PRO
ASP
GLU
GLU
GLU
GLU
L170
S235
K171
G174
H175
Y176
I239
S240
R177
S178
A179
K180
V181
D182
N183
G184
V184
V185
I886
C187
D191

L253
S254
E255
E256
L262
D263
C264
L265
R266
S267
Q268
V269
K270
I271
T272
R273
W274
D275
F276
M277
D279
P280
K281
A282
R283
D284
A284
Q285
L286
L287
E288
S289
C290
G352
D291
L292
K353
Y293
Y294
D295
M296
S297
Y298
S299
V300
A301
Y302
S303
T304
F305
A306
N307
T308
SER
SER
SER
SER
GLU
GLY
GLN
SER
GLY
SER
GLY
SER
GLY
D439
F379

SER
ASP
THR
ALA
SER
GLY
ILE
SER
SER
ASP
ASP
VAL
ASP
LEU
GLU
THR
THR
SER
SER
SER
SER
PRO
R339
T342
L343
L344
D345
L346
Y347
S348
G349
C350
G351
G352
D291
L292
K353
Y293
Y294
D295
M296
S297
Y298
S299
V300
A301
Y302
S303
T304
F305
A306
N307
T308
SER
SER
SER
SER
GLU
GLY
GLN
SER
GLY
SER
GLY
D439
F379

A390
G391
Q392
S393
L394
K395
Y396
M397
H398
P399
Q390
T391
S392
V393
R394
S395
E396
R399
E400
F401
L402
A403
L404
L405
K406
E407
W408
A409
Y410
L411
C412
R413
K414
Y415
VAL
GLN
ASP
VAL
ASP
SER
ASN
LEU
ALA
SER
SER
GLU
ASP
PRO
LEU
ALA
SER
SER
SER
GLU
ASP
PRO
LEU
LEU
ASP
LYS
D439
E440

F441
V442
E443
E444
L445
K446
V447
G448
H449
C450
Y451
G452
K268
S391
S454
D455
R456
E457
N458
G459
L460
Y461
F462
V463
L464
O465
K466
E467
W468
A409
Y469
E472
E473
D474
Y475
N476
E477
P478
L479
L482
S483
D484
C485
P486
Q487
K488
L489
R490
A491
F492
V493
Q494
G495
G496
L497
K498
R499
S500
L501
L502

P503
L504
P505
G506
D507
V508
D509
S510
F511
C512
S513
G514
P515
P516
C517
F518
G519
ILE
SER
GLY
PHE
GLY
ASN
ARG
TVR
ARG
ASN
K534
D535
E536
Y475
K537
N538
K539
Q540
N541
V542
T543
F544
N545
D546
I547
V548
A549
Y550
L551
K552
P553
K554
L555
V556
G557
L557
N558
E559
N560
V561
V562

D563
I564
L565
G570
Y571
L572
G573
K574
F575
Y576
A577
S578
C579
L580
V581
V582
N583
K584
Y585
Q586
R587
S588
L589
G590
M591
N592
C596
L599
P600
Q601
F602
R603
M604
R605
V606
F607
L608
G609
G610
A611
S614
M615
V616
L617
P618
K619
Y620
P621
L622
P623
T624
L625
V626
E626
V629

A633
P634
M635
A636
F637
S638
Q639
C640
M641
V642
A643
Y644
D645
E646
T647
Q648
K649
P650
S651
L652
R653
K654
A655
L656
L657
L658
G659
D660
G661
A662
P663
D664
L665
P666
F667
V668
Q669
M670
H671
Q672
P673
N674
D675
V676
M677
E678
Y679
G680
G681
S682
P683
T684
E685
L622
G686
P687
H688
Q688
Y689
I691
R692

L693
S694
R695
K696
D697
M698
L699
D700
P701
F702
W703
F704
G705
H706
Q707
D710
K713
L714
L715
D716
P717
Q718
P719
W720
R721
E722
N723
N724
D725
T726
Y727
A661
I662
S663
D664
L665
P666
F667
V668
Q669
M670
H671
Q672
P673
N674
D675
V676
M677
E678
Y679
G680
G681
S682
P683
T684
E685
L622
G686
P687
H688
Q688
Y689
I691
R692

L772
V773
P774
D775
Y776
A777
M778
S779
P841
I842
K843
E844
K845
L846
I847
Q848
A852
V855
P856
A860
L861
G862
Y863
L864
G865
G866
Q867
A868
R804
E806
P807
H808
N809
Q810
V811
I812
I813
H814
P815
T816
Q817
R818
R819
L820
L821
L822
I823
R824
A751
E825
N826
SER
PRG
L828
S767
S768
P771

P833
D834
Y835
Y836
R837
L838
F839
G840
P841
I842
K843
E844
K845
L846
I847
Q848
A852
V855
P856
A860
L861
G862
Y863
L864
G865
G866
Q867
A868
P868
L879
W880
Q881
L882
P883
P884
S885
THR
THR
SER
VAL
GLY
GLY
ARG
THR
ALA
ALA
GLN
ALA
ARG
ALA
SER
PRO
VAL
GLY
THR
PRO

ALA
GLY
GLU
VAL
VAL
GLU
GLN

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	64.86Å 88.95Å 113.49Å 93.47° 95.53° 110.41°	Depositor
Resolution (Å)	40.25 – 3.20 48.87 – 3.19	Depositor EDS
% Data completeness (in resolution range)	97.6 (40.25-3.20) 98.9 (48.87-3.19)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	0.17	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 3.19Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, R_{free}	0.240 , 0.263 0.230 , 0.254	Depositor DCC
R_{free} test set	1946 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	59.9	Xtrriage
Anisotropy	0.609	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 69.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	10631	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/5445	0.79	5/7358 (0.1%)
1	B	0.63	1/5362 (0.0%)	0.82	12/7252 (0.2%)
All	All	0.63	1/10807 (0.0%)	0.80	17/14610 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	1
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	579	CYS	CB-SG	-5.50	1.72	1.81

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	535	ASP	CB-CG-OD1	-9.04	110.17	118.30
1	A	632	GLY	N-CA-C	-8.60	91.61	113.10
1	B	535	ASP	N-CA-C	8.56	134.12	111.00
1	B	146	GLU	N-CA-C	-6.83	92.56	111.00
1	B	746	GLY	N-CA-C	-6.81	96.08	113.10

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	145	ASP	Peptide
1	A	455	ASP	Peptide
1	A	507	ASP	Peptide
1	B	504	LEU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5331	0	5166	987	0
1	B	5248	0	5023	1014	0
2	A	26	0	19	6	0
2	B	26	0	19	6	0
All	All	10631	0	10227	1998	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 96.

The worst 5 of 1998 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:695:ARG:HG2	1:B:835:TYR:CD1	1.50	1.46
1:B:656:LEU:C	1:B:657:LEU:HD12	1.33	1.45
1:A:134:GLU:HB3	1:A:135:PRO:CD	1.43	1.38
1:B:883:PRO:CG	1:B:884:PRO:HD2	1.53	1.37
1:B:695:ARG:HG2	1:B:835:TYR:CE1	1.64	1.32

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	670/784 (86%)	591 (88%)	62 (9%)	17 (2%)	5	32
1	B	665/784 (85%)	607 (91%)	43 (6%)	15 (2%)	6	34
All	All	1335/1568 (85%)	1198 (90%)	105 (8%)	32 (2%)	6	34

5 of 32 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	142	VAL
1	A	634	PRO
1	A	646	GLU
1	B	140	SER
1	B	279	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	564/652 (86%)	413 (73%)	151 (27%)	0	2
1	B	544/652 (83%)	384 (71%)	160 (29%)	0	1
All	All	1108/1304 (85%)	797 (72%)	311 (28%)	0	1

5 of 311 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	813	ILE
1	B	234	ASN
1	B	732	GLN
1	A	839	PHE
1	B	149	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	814	HIS
1	B	183	ASN
1	B	810	GLN
1	A	817	GLN
1	B	175	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	SAH	B	1000	-	21,28,28	1.70	5 (23%)	20,40,40	1.93	5 (25%)
2	SAH	A	1000	-	21,28,28	1.70	5 (23%)	20,40,40	1.93	5 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAH	B	1000	-	-	3/7/31/31	0/3/3/3
2	SAH	A	1000	-	-	5/7/31/31	0/3/3/3

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1000	SAH	C2-N3	5.02	1.40	1.32
2	B	1000	SAH	C2-N3	4.97	1.40	1.32
2	B	1000	SAH	C2-N1	3.27	1.40	1.33
2	A	1000	SAH	C2-N1	3.26	1.40	1.33
2	B	1000	SAH	C5-C4	-2.68	1.33	1.40

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	SAH	N3-C2-N1	-6.76	118.11	128.68
2	B	1000	SAH	N3-C2-N1	-6.72	118.18	128.68
2	B	1000	SAH	C5-C6-N6	-2.69	116.26	120.35
2	A	1000	SAH	C5-C6-N6	-2.69	116.27	120.35
2	B	1000	SAH	C5'-C4'-C3'	-2.22	109.50	115.06

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1000	SAH	O4'-C4'-C5'-SD
2	A	1000	SAH	N-CA-CB-CG
2	A	1000	SAH	C-CA-CB-CG
2	A	1000	SAH	O4'-C4'-C5'-SD
2	A	1000	SAH	C3'-C4'-C5'-SD


There are no ring outliers.


2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1000	SAH	6	0
2	A	1000	SAH	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	665/784 (84%)	-0.17	9 (1%) 75 63	21, 62, 129, 200	0
1	B	660/784 (84%)	-0.25	4 (0%) 89 83	17, 69, 134, 285	0
All	All	1325/1568 (84%)	-0.21	13 (0%) 82 72	17, 65, 132, 285	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	759	PRO	3.8
1	A	746	GLY	3.3
1	A	484	ASP	3.1
1	A	445	LYS	2.6
1	B	729	ARG	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SAH	A	1000	26/26	0.91	0.29	42,53,71,112	0
2	SAH	B	1000	26/26	0.95	0.19	31,55,87,93	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.