

Full wwPDB X-ray Structure Validation Report (i)

May 13, 2020 – 11:13 am BST

PDB ID	:	6FUZ
Title	:	Crystal structure of the TPR domain of KLC1 in complex with the C-terminal
		peptide of JIP1
Authors	:	Pernigo, S.; Dodding, M.P.; Steiner, R.A.
Deposited on	:	2018-02-28
Resolution	:	2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as 541 be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
$\operatorname{CCP4}$:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069(2.70-2.70)
Sidechain outliers	138945	3069(2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Qu	ality of chain		
			18%			
1	А	337	64%	14%	• 21%	6
	3.7	101	8%			
2	N	121	77%		16%	7%



$6 \mathrm{FUZ}$

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3020 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Kinesin light chain 1,Kinesin light chain 1,C-Jun-amino-termi nal kinase-interacting protein 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	967	Total	С	Ν	Ο	\mathbf{S}	0	0	0
L	Л	207	2140	1340	381	411	8	0	0	0

Chain	Residue	Modelled	Actual	Comment	Reference
A	201	GLY	ALA	conflict	UNP Q5UE59
А	202	SER	ALA	conflict	UNP Q5UE59
A	203	HIS	GLN	conflict	UNP Q5UE59
А	204	MET	GLN	conflict	UNP Q5UE59
A	671	THR	-	linker	UNP Q5UE59
A	672	GLY	-	linker	UNP Q5UE59
A	673	SER	-	linker	UNP Q5UE59
A	674	THR	-	linker	UNP Q5UE59
A	675	GLY	-	linker	UNP Q5UE59
А	676	SER	-	linker	UNP Q5UE59
A	677	THR	-	linker	UNP Q5UE59
А	678	GLY	-	linker	UNP Q5UE59
A	679	SER	-	linker	UNP Q5UE59
A	680	THR	-	linker	UNP Q5UE59
А	681	GLY	-	linker	UNP Q5UE59
A	682	SER	-	linker	UNP Q5UE59
A	683	THR	-	linker	UNP Q5UE59
A	684	GLY	-	linker	UNP Q5UE59
A	685	SER	-	linker	UNP Q5UE59
A	686	THR	-	linker	UNP Q5UE59
A	687	GLY	-	linker	UNP Q5UE59
А	688	SER	-	linker	UNP Q5UE59
А	689	THR	-	linker	UNP Q5UE59
A	690	GLY	-	linker	UNP Q5UE59
A	691	SER	-	linker	UNP Q5UE59
A	692	THR	-	linker	UNP Q5UE59

There are 34 discrepancies between the modelled and reference sequences:



00100100		ere ae page			
Chain	Residue	Modelled	Actual	Comment	Reference
А	693	GLY	-	linker	UNP Q5UE59
А	694	SER	-	linker	UNP Q5UE59
A	695	THR	-	linker	UNP Q5UE59
А	696	GLY	-	linker	UNP Q5UE59
А	697	SER	-	linker	UNP Q5UE59
А	698	THR	-	linker	UNP Q5UE59
A	699	GLY	-	linker	UNP Q5UE59
А	700	SER	-	linker	UNP Q5UE59

• Molecule 2 is a protein called nanobody.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	Ν	112	Total 861	C 541	N 147	O 168	$\frac{S}{5}$	0	0	0

• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	Ν	1	Total 6	С 3	O 3	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	7	Total O 7 7	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	Ν	6	Total O 6 6	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Kinesin light chain 1, Kinesin light chain 1, C-Jun-amino-terminal kinase-interacting protein 1







4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	107.47Å 90.36 Å 51.70 Å	Depositor
a, b, c, α , β , γ	90.00° 99.80° 90.00°	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	45.18 - 2.70	Depositor
Resolution (A)	45.18 - 2.70	EDS
% Data completeness	98.4 (45.18-2.70)	Depositor
(in resolution range)	98.4(45.18-2.70)	EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.53 (at 2.69 \text{\AA})$	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
D D .	0.233 , 0.257	Depositor
n, n_{free}	0.244 , 0.283	DCC
R_{free} test set	669 reflections $(5.06%)$	wwPDB-VP
Wilson B-factor (Å ²)	84.9	Xtriage
Anisotropy	0.271	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	$0.35 \;, 80.4$	EDS
L-test for $twinning^2$	$ < L >=0.49, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3020	wwPDB-VP
Average B, all atoms $(Å^2)$	113.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal Chain		Bond	lengths	Bond angles		
	Cham	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.45	0/2174	0.66	0/2929	
2	Ν	0.39	0/880	0.66	0/1191	
All	All	0.43	0/3054	0.66	0/4120	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2140	0	2141	27	0
2	N	861	0	824	10	0
3	N	6	0	8	1	0
4	А	7	0	0	0	0
4	N	6	0	0	0	0
All	All	3020	0	2973	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:475:ARG:HH21	1:A:487:GLU:HG3	1.63	0.64
1:A:271:TYR:HB3	1:A:310:ARG:HD2	1.80	0.62
1:A:226:GLN:HB3	1:A:228:ARG:HG3	1.83	0.61
1:A:294:PRO:HG3	1:A:330:VAL:HG11	1.81	0.60
1:A:475:ARG:NH2	1:A:487:GLU:HG3	2.20	0.57
2:N:39:GLN:HB2	2:N:45:LEU:HD12	1.87	0.56
1:A:468:LYS:HE3	1:A:494:ARG:HH21	1.71	0.55
2:N:83:MET:HE1	2:N:111:VAL:HG11	1.89	0.55
2:N:83:MET:HE2	2:N:86:LEU:HD21	1.89	0.54
1:A:399:GLN:HG2	3:N:201:GOL:H32	1.89	0.54
1:A:370:GLN:OE1	2:N:33:TRP:HB2	2.10	0.52
1:A:406:GLU:HB3	1:A:410:ARG:HH12	1.78	0.48
1:A:336:PRO:HD3	1:A:372:LYS:HG2	1.94	0.48
1:A:343:ASN:HD21	1:A:386:ASN:ND2	2.11	0.47
1:A:251:HIS:CD2	1:A:253:ASP:H	2.32	0.47
2:N:37:VAL:HG12	2:N:47:TRP:HA	1.97	0.47
1:A:362:TYR:CE1	1:A:386:ASN:HB3	2.51	0.46
1:A:350:GLN:HG3	1:A:393:LYS:HE2	1.98	0.45
1:A:340:LYS:HG3	1:A:710:LEU:HB2	1.98	0.45
1:A:216:LEU:O	1:A:220:VAL:HG23	2.17	0.45
2:N:102:GLY:HA3	2:N:105:ARG:HD3	1.98	0.44
1:A:312:LYS:HG2	1:A:315:GLU:HB2	1.99	0.44
1:A:298:ALA:HA	1:A:327:ARG:HH11	1.82	0.44
1:A:331:LEU:HD13	1:A:337:ASP:HB2	2.00	0.44
1:A:340:LYS:HA	1:A:710:LEU:HD12	1.99	0.44
1:A:327:ARG:HH22	1:A:340:LYS:HD2	1.83	0.43
2:N:50:THR:HB	2:N:59:TYR:HB2	2.00	0.43
1:A:425:PRO:HD2	1:A:428:MET:HG3	2.00	0.43
1:A:275:ALA:HB2	1:A:306:LEU:HD21	2.00	0.43
1:A:217:HIS:CE1	1:A:260:ILE:HD11	2.54	0.42
1:A:421:ASP:HA	1:A:424:LYS:HE2	2.01	0.42
2:N:95:TYR:HB3	2:N:105:ARG:HB2	2.02	0.42
2:N:22:CYS:HB3	2:N:79:LEU:HB3	2.02	0.41
1:A:475:ARG:HG3	1:A:475:ARG:HH11	1.86	0.41
1:A:375:PRO:HG3	2:N:32:TYR:CE1	2.56	0.40

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Favoured Allowed		Allowed	Outliers	Perce	entiles
1	А	261/337~(77%)	253~(97%)	7(3%)	1 (0%)	34	60
2	Ν	$110/121 \ (91\%)$	106~(96%)	3(3%)	1 (1%)	17	40
All	All	371/458 (81%)	359(97%)	10 (3%)	2(0%)	29	54

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	417	GLY
2	Ν	41	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	226/275 (82%)	211~(93%)	15 (7%)	16 38
2	Ν	90/99~(91%)	88 (98%)	2(2%)	52 79
All	All	316/374~(84%)	299~(95%)	17~(5%)	22 47

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	216	LEU
1	А	226	GLN
1	А	236	CYS
1	А	272	LYS



Mol	Chain	Res	Type
1	А	312	LYS
1	А	315	GLU
1	А	350	GLN
1	А	357	GLU
1	А	376	ASP
1	А	393	LYS
1	А	416	PHE
1	А	426	ILE
1	А	433	ARG
1	А	465	THR
1	А	467	LEU
2	N	12	VAL
2	N	112	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	276	ASN
1	А	343	ASN
1	А	386	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tune	Chain	nin Dog Lin		B	Bond lengths			Bond angles		
	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
3	GOL	Ν	201	-	5, 5, 5	0.31	0	5,5,5	0.30	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	Ν	201	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	N	201	GOL	O1-C1-C2-C3
3	N	201	GOL	C1-C2-C3-O3
3	N	201	GOL	O2-C2-C3-O3
3	N	201	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	N	201	GOL	1	0

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< RSRZ >	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	267/337~(79%)	1.29	59~(22%) 0 0	79, 112, 153, 180	0
2	Ν	112/121~(92%)	0.71	10 (8%) 9 7	82, 104, 124, 132	0
All	All	379/458~(82%)	1.12	69 (18%) 1 1	79, 108, 149, 180	0

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	229	TYR	6.9
1	А	247	SER	5.4
1	А	271	TYR	5.3
1	А	433	ARG	4.7
2	Ν	11	LEU	4.6
1	А	235	LEU	4.5
1	А	410	ARG	4.0
1	А	223	TYR	3.8
1	А	228	ARG	3.7
1	А	306	LEU	3.7
1	А	254	VAL	3.7
1	А	434	GLU	3.5
1	А	243	LEU	3.5
1	А	331	LEU	3.5
1	А	484	GLU	3.4
1	А	392	LEU	3.4
1	А	216	LEU	3.3
1	А	291	ARG	3.2
1	А	480	PHE	3.1
1	A	470	LEU	3.0
1	A	479	LYS	2.9
1	A	$\overline{365}$	ALA	2.8
2	N	86	LEU	2.8
2	Ν	59	TYR	2.8



6FI	UΖ
-----	----

Mol	Iol Chain Res		Type	RSRZ
1	А	288	THR	2.8
1	А	366	LEU	2.7
1	А	426	ILE	2.7
2	N	112	THR	2.7
1	А	387	LEU	2.7
1	А	481	GLU	2.7
1	А	249	HIS	2.6
1	А	487	GLU	2.6
1	А	430	ALA	2.5
1	А	482	ALA	2.5
1	А	240	LEU	2.5
2	N	18	LEU	2.5
1	А	471	GLY	2.5
1	А	477	GLN	2.5
1	A	322	ARG	2.5
1	А	224	ALA	2.5
1	А	310	ARG	2.4
1	А	495	LYS	2.4
1	А	375	PRO	2.4
1	А	219	LEU	2.4
1	А	257	MET	2.4
1	А	286	GLU	2.3
1	А	362	TYR	2.3
1	А	278	LEU	2.3
1	А	358	VAL	2.3
1	А	373	LEU	2.3
1	А	490	ALA	2.3
1	А	220	VAL	2.3
1	А	217	HIS	2.2
1	А	232	ALA	2.2
1	A	296	VAL	2.2
2	N	45	LEU	2.2
2	N	94	TYR	2.1
1	A	332	GLY	2.1
1	A	491	MET	2.1
1	A	289	LEU	2.1
1	A	486	LEU	2.1
1	A	221	ILE	2.1
1	A	251	HIS	2.1
1	A	330	VAL	2.1
2	N	111	VAL	2.0
1	A	346	ALA	2.0



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	Ν	76	LYS	2.0
2	Ν	57	ILE	2.0
1	А	389	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	$Q{<}0.9$
3	GOL	N	201	6/6	0.74	0.20	$101,\!102,\!103,\!103$	0

6.5 Other polymers (i)

There are no such residues in this entry.

