



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 19, 2023 – 11:28 PM EDT

PDB ID : 2FX7
Title : Crystal structure of hiv-1 neutralizing human fab 4e10 in complex with a 16-residue peptide encompassing the 4e10 epitope on gp41
Authors : Cardoso, R.M.F.; Brunel, F.M.; Ferguson, S.; Burton, D.R.; Dawson, P.E.; Wilson, I.A.
Deposited on : 2006-02-03
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	214	
2	H	228	
3	P	16	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	L	701	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3829 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab 4E10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	214	1639	1017	284	334	4	0	0	0

- Molecule 2 is a protein called Fab 4E10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	228	1690	1069	289	327	5	0	0	0

- Molecule 3 is a protein called Fragment of HIV glycoprotein (GP41).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	P	16	157	109	25	23	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
P	684	LYS	ILE	engineered mutation	UNP P05880
P	685	LYS	PHE	engineered mutation	UNP P05880
P	686	LYS	ILE	engineered mutation	UNP P05880

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	L	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	L	173	Total O 173 173	0	0
5	H	155	Total O 155 155	0	0
5	P	9	Total O 9 9	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab 4E10



- Molecule 2: Fab 4E10



- Molecule 3: Fragment of HIV glycoprotein (GP41)



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	157.94Å 44.66Å 85.26Å 90.00° 113.09° 90.00°	Depositor
Resolution (Å)	50.00 – 1.76 36.32 – 1.76	Depositor EDS
% Data completeness (in resolution range)	98.7 (50.00-1.76) 99.0 (36.32-1.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.53 (at 1.76Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.202 , 0.222 0.196 , 0.199	Depositor DCC
R_{free} test set	2738 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	19.3	Xtrriage
Anisotropy	0.181	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 51.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3829	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	0.31	0/1672	0.63	0/2266
2	H	0.30	0/1733	0.64	0/2366
3	P	0.35	0/164	0.40	0/220
All	All	0.31	0/3569	0.62	0/4852

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1639	0	1589	17	0
2	H	1690	0	1679	27	0
3	P	157	0	155	7	0
4	L	6	0	4	0	0
5	H	155	0	0	1	0
5	L	173	0	0	1	0
5	P	9	0	0	1	2
All	All	3829	0	3427	48	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:62:ARG:HB2	2:H:62:ARG:HH11	1.34	0.93
2:H:38:ARG:HG3	2:H:43:ARG:NH2	1.97	0.80
1:L:198:HIS:CD2	1:L:200:GLY:H	2.02	0.78
3:P:681:TYR:O	3:P:685:LYS:HG3	1.94	0.68
1:L:108:ARG:HD3	1:L:109:THR:O	1.96	0.66
2:H:100(E):LYS:HD2	3:P:677:ASN:HD21	1.60	0.65
3:P:678:TRP:CH2	3:P:682:ILE:HD11	2.35	0.62
2:H:212:HIS:HD2	2:H:215:SER:OG	1.82	0.62
2:H:119:PRO:HB3	2:H:147:TYR:HB3	1.82	0.61
2:H:30:SER:HA	2:H:52(A):PRO:HB2	1.84	0.60
1:L:198:HIS:HD2	1:L:200:GLY:H	1.45	0.60
2:H:149:PRO:O	2:H:212:HIS:HE1	1.85	0.59
3:P:678:TRP:CZ3	3:P:682:ILE:HD11	2.37	0.59
2:H:52:ILE:HB	2:H:56:ILE:CG2	2.34	0.58
2:H:40:ALA:HB3	2:H:43:ARG:HD3	1.86	0.56
1:L:185:ASP:HA	1:L:188:LYS:HD3	1.86	0.56
1:L:122:ASP:O	1:L:126:LYS:HG3	2.05	0.55
2:H:6:GLN:HE21	2:H:104:GLY:HA3	1.71	0.55
1:L:37:GLN:HB2	1:L:47:LEU:HD11	1.87	0.55
2:H:40:ALA:H	2:H:43:ARG:NE	2.06	0.53
1:L:18:ARG:HD2	5:L:854:HOH:O	2.08	0.53
1:L:2:ILE:HG12	1:L:27:GLN:CG	2.40	0.52
2:H:40:ALA:O	2:H:43:ARG:HG2	2.10	0.51
1:L:210:ASN:HB2	1:L:213:GLU:OE2	2.11	0.50
1:L:183:LYS:O	1:L:187:GLU:HG3	2.12	0.50
2:H:40:ALA:N	2:H:43:ARG:HG2	2.28	0.48
1:L:124:GLN:HG2	1:L:129:THR:O	2.14	0.47
2:H:1:GLN:HG3	2:H:100(B):TRP:CH2	2.50	0.47
1:L:160:GLN:HE22	2:H:179:GLN:HA	1.80	0.47
2:H:38:ARG:HG3	2:H:43:ARG:HH21	1.73	0.46
1:L:141:PRO:O	1:L:198:HIS:HE1	1.99	0.46
2:H:6:GLN:HE22	2:H:91:TYR:HA	1.80	0.46
2:H:62:ARG:HD3	5:H:368:HOH:O	2.15	0.45
2:H:187:LEU:C	2:H:187:LEU:HD12	2.38	0.45
3:P:678:TRP:HZ2	5:P:331:HOH:O	2.00	0.45
2:H:62:ARG:HH11	2:H:62:ARG:CB	2.16	0.44
2:H:47:TRP:CZ2	2:H:49:GLY:HA2	2.52	0.44
3:P:678:TRP:CZ3	3:P:682:ILE:CD1	3.00	0.44
2:H:100(E):LYS:HD2	3:P:677:ASN:ND2	2.30	0.44
2:H:100(J):PHE:N	2:H:100(J):PHE:CD2	2.84	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:66:ARG:HD2	2:H:82(A):ASN:O	2.18	0.42
1:L:55:PRO:HG2	1:L:58:VAL:CG2	2.50	0.42
1:L:108:ARG:CD	1:L:109:THR:O	2.67	0.42
2:H:211:ASN:HD21	2:H:218:LYS:HE2	1.85	0.42
2:H:62:ARG:HB2	2:H:62:ARG:NH1	2.16	0.41
1:L:146:VAL:HG13	1:L:194:CYS:SG	2.61	0.40
1:L:129:THR:HG22	1:L:130:ALA:N	2.36	0.40
2:H:52:ILE:HB	2:H:56:ILE:HG22	2.01	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:P:91:HOH:O	5:P:219:HOH:O[2_657]	0.59	1.61
5:P:187:HOH:O	5:P:330:HOH:O[2_657]	0.67	1.53

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	212/214 (99%)	207 (98%)	5 (2%)	0	100	100
2	H	226/228 (99%)	218 (96%)	8 (4%)	0	100	100
3	P	14/16 (88%)	14 (100%)	0	0	100	100
All	All	452/458 (99%)	439 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	184/184 (100%)	182 (99%)	2 (1%)	73	60
2	H	187/187 (100%)	184 (98%)	3 (2%)	62	45
3	P	16/16 (100%)	16 (100%)	0	100	100
All	All	387/387 (100%)	382 (99%)	5 (1%)	69	54

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	105	GLU
1	L	108	ARG
2	H	62	ARG
2	H	100(C)	LEU
2	H	151	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	31	ASN
1	L	42	GLN
1	L	93	GLN
1	L	100	GLN
1	L	147	GLN
1	L	160	GLN
1	L	198	HIS
1	L	210	ASN
2	H	1	GLN
2	H	6	GLN
2	H	58	ASN
2	H	82(A)	ASN
2	H	211	ASN
2	H	212	HIS
3	P	677	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	L	701	-	5,5,5	4.49	5 (100%)	5,5,5	5.76	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	L	701	-	-	3/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	L	701	GOL	C3-C2	-7.31	1.21	1.51
4	L	701	GOL	O1-C1	4.78	1.62	1.42
4	L	701	GOL	O3-C3	3.37	1.56	1.42
4	L	701	GOL	C1-C2	-2.66	1.40	1.51
4	L	701	GOL	O2-C2	-2.51	1.35	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	L	701	GOL	O3-C3-C2	10.31	159.63	110.20
4	L	701	GOL	O2-C2-C3	6.81	139.12	109.12
4	L	701	GOL	O1-C1-C2	3.48	126.88	110.20

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	L	701	GOL	O1-C1-C2-C3
4	L	701	GOL	C1-C2-C3-O3
4	L	701	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	214/214 (100%)	-0.07	3 (1%) 75 82	13, 21, 32, 52	0
2	H	228/228 (100%)	0.96	24 (10%) 6 8	14, 20, 51, 65	0
3	P	16/16 (100%)	1.48	5 (31%) 0 0	18, 29, 62, 66	0
All	All	458/458 (100%)	0.49	32 (6%) 16 21	13, 20, 45, 66	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	100	TRP	15.6
2	H	100(B)	TRP	14.4
2	H	133	THR	13.8
2	H	99	GLY	11.6
2	H	134	SER	11.1
2	H	100(A)	GLY	11.0
2	H	100(C)	LEU	10.7
2	H	128	SER	9.1
2	H	130	SER	9.0
2	H	135	GLY	8.3
2	H	129	LYS	7.5
2	H	1	GLN	5.4
2	H	200	THR	5.0
3	P	686	LYS	4.8
1	L	213	GLU	4.5
2	H	28	SER	4.2
2	H	228	LYS	4.1
3	P	684	LYS	4.1
2	H	27	GLY	3.9
2	H	26	GLY	3.8
2	H	25	SER	3.7
1	L	212	GLY	3.7
2	H	127	SER	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	P	678	TRP	3.1
2	H	199	GLY	3.0
3	P	681	TYR	3.0
2	H	2	VAL	2.7
1	L	194	CYS	2.7
3	P	683	LYS	2.4
2	H	98	THR	2.4
2	H	3	GLN	2.2
2	H	136	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	L	701	6/6	0.67	0.22	23,30,32,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.